

Reply to TR Hill and I Kyriazakis

Article

Accepted Version

Guo, J., Jackson, K. G. ORCID: <https://orcid.org/0000-0002-0070-3203>, Givens, D. I. and Lovegrove, J. A. ORCID: <https://orcid.org/0000-0001-7633-9455> (2018) Reply to TR Hill and I Kyriazakis. *Journal of Nutrition*, 148 (4). p. 665. ISSN 1541-6100 doi: <https://doi.org/10.1093/jn/nxy010> Available at <https://centaur.reading.ac.uk/80930/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1093/jn/nxy010>

Publisher: American Society for Nutrition

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Reply to the Letter to the Editor for “A 25-hydroxycholecalciferol-fortified dairy drink is more effective at raising a marker of postprandial vitamin D status than cholecalciferol in men with suboptimal vitamin D status.” (Manuscript doi: 10.3945/jn.117.254789) by

Jing Guo, Kim G Jackson, Che Suhaili binti Che Taha, Yue Li, David I Givens, and Julie A Lovegrove

¹ From the Institute for Food, Nutrition and Health (JG, KGJ, DIG, JAL); Hugh Sinclair Unit of Human Nutrition (JG, KGJ, YL, DIG, JAL); Institute for Cardiovascular and Metabolic Research (JG, KGJ, DIG, JAL); 2 School of Psychology and Clinical Language Sciences (CSBCT), University of Reading, Reading, RG6 6AP, United Kingdom.

² Corresponding author: Julie A. Lovegrove, Hugh Sinclair Unit of Human Nutrition, Department of Food & Nutritional Sciences, Whiteknights, PO Box 226, University of Reading, Reading, RG6 6AP, United Kingdom. E-mail: j.a.lovegrove@reading.ac.uk.

³ Author names for indexing: Guo, Jackson, Che Taha, Li, Givens and Lovegrove.

⁴ Word count: 352

⁵ Figures: 0

⁶ Tables: 0

⁷ OSM submitted: 0

⁸ Supported by the Barham Benevolent Foundation.

⁹ Abbreviations used: NA.

¹⁰ Running title: Dairy drink fortification with vitamin D isoforms

¹¹ Author names for indexing: Guo, Jackson, Che Taha, Li, Givens, Lovegrove.

¹² Author disclosures: JG, KGJ, CSCT, YL, DIG and JAL, no conflicts of interest.

We thank Drs Thomas R Hill and Ilias Kyriazakis for their comments on our paper. We agree that a clean label approach for food vitamin D enrichment is favoured by the consumer and the low levels of vitamin D₃ and 25(OH) D₃ naturally present in animal derived foods, such as eggs and milk, can be significantly increased by supplemental additions of vitamin D₃ and 25(OH) D₃ to the animals' diets (biofortification) (1). However, although a statistically significant increase in vitamin D₃ and 25(OH) D₃ has been reported after biofortification at supplemental quantities in line with EU legislation (2), these changes are quantitatively trivial and would not contribute to increase in dietary vitamin D₃ intake and human vitamin D status as stated by Drs Hill and Kyriazakis (1). We confirmed this in a recent study (3) in which dairy cows' diets were supplemented either with 0.075mg/kg vitamin D₃ (control), the maximum permitted dose of vitamin D₃ (0.1mg/kg) recommended by the EU (2), or with 0.03mg/kg vitamin D₃ plus 25(OH) D₃ (0.075 mg/kg) for 8 weeks feeding from calving to early lactation. The vitamin D₃ and 25(OH) D₃ concentrations in milk from both treatments were not significantly different to the control milk or to themselves (3). For a typical milk serving of 200 ml would contribute 0.02 to 0.66 µg vitamin D (3), which well below the current UK vitamin D recommended intake of 10 µg/day (4). The authors believe that without changes to the permitted dietary supplementation levels in dairy diets, milk fortification with vitamin D, may be a more feasible strategy to increase dietary vitamin D₃ intake and ultimately increase population vitamin D status, than biofortification.

Our current finding that a dairy drink fortified with 25(OH) D₃ was more effective at raising plasma 25(OH) D₃ concentrations than dairy drink fortified with vitamin D₃ in men with suboptimal vitamin D status supported previous studies (5, 6), which demonstrates the value of 25(OH) D₃ food fortification. However this would require changes in the EU legislation before the potential advantage of this form of vitamin D can be realised for food fortification in the EU.

References

1. Guo J, Lovegrove JA, Givens DI. 25(OH) D₃ enriched or fortified foods are more efficient at tackling inadequate vitamin D status than vitamin D₃. Proc Nutr Soc. 2017 [Accepted and in press].
2. Commission Implementing Regulation (EU) No 2017/1492 of 21 August 2017 concerning the authorisation of cholecalciferol as a feed additive for all animal species. Official Journal of the European Union L216/19.
3. Guo J, Jones AK, Givens DI, Lovegrove JA, Kliem KE. Effect of dietary vitamin D₃ and 25(OH) D₃ supplementation on plasma and milk 25(OH) D₃ concentration in dairy cows. J Dairy Sci [under review].
4. SACN. 2015. Draft Vitamin D and Health Report. Accessed Oct. 20, 2016. <https://www.gov.uk/government/consultations/consultation-on-draft-sacn-vitamin-d-andhealth-report>.
5. Cashman KD, Seamans KM, Lucey AJ, Stocklin E, Weber P, Kiely M, Hill TR. Relative effectiveness of oral 25-hydroxyvitamin D₃ and vitamin D₃ in raising wintertime serum 25-hydroxyvitamin D in older adults. Am J Clin Nutr 2012;95: 1350–1356.
6. Jetter A, Egli A, Dawson-Hughes B, Staehelin HB, Stoecklin E, Goessl R, Henschkowski J, Bischoff-Ferrari HA. Pharmacokinetics of oral vitamin D-3 and calcifediol. Bone 2014;59:14–9.