

Measuring the unmeasurable? A method to quantify adoption of integrated pest management practices in temperate arable farming systems

Article

Accepted Version

Creissen, H., Jones, P., Tranter, R., Girling, R., Jess, S., Burnett, F., Gaffney, M., Thorne, F. and Kildea, S. (2019) Measuring the unmeasurable? A method to quantify adoption of integrated pest management practices in temperate arable farming systems. *Pest Management Science*, 75 (12). pp. 3144-3152. ISSN 1526-4998 doi: <https://doi.org/10.1002/ps.5428> Available at <https://centaur.reading.ac.uk/81514/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1002/ps.5428>

Publisher: Wiley

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online



Measuring the unmeasurable? A method to quantify adoption of Integrated Pest Management practices in temperate arable farming systems

Journal:	<i>Pest Management Science</i>
Manuscript ID	PM-19-0057.R1
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Creissen, Henry; Scotland's Rural College, Agriculture Department Jones, Philip; University of Reading, Centre for Agricultural Strategy Tranter, Richard; University of Reading, Centre for Agricultural Strategy Girling, Robbie; University of Reading, Centre for Agri-Environmental Research Jess, Stephen; Agri-Food and Biosciences Institute for Northern Ireland, Applied Plant Science Burnett, Fiona; Scotland's Rural College, Agriculture Department Gaffney, Michael; Teagasc, Horticultural Development Department Thorne, Fiona; Teagasc, Agricultural Economics and Farm Surveys Department Kildea, Steven; Teagasc Crops Research Centre, Crop Production and Protection
Key Words:	Integrated Pest Management, IPM scoring system, IPM metric, arable farming, farmer survey, sustainable agriculture, National Action Plan, Sustainable Use Directive, IPM uptake, IPM adoption

SCHOLARONE™
Manuscripts

1
2
3 1 Title: **Measuring the unmeasurable? A method to quantify adoption of**
4
5
6 2 **Integrated Pest Management practices in temperate arable farming systems**
7
8

9
10 3 Running title: **Quantifying adoption of Integrated Pest Management in arable**
11
12 4 **farming**
13
14

15
16 5 **Henry E. Creissen ^{a, b *}, Philip J. Jones ^c, Richard B. Tranter ^c, Robbie D. Girling ^d,**
17
18 6 **Stephen Jess ^e, Fiona J. Burnett ^a, Michael Gaffney ^f, Fiona S. Thorne ^g, Steven Kildea ^b**
19
20

21
22 7 ^a Agriculture Department, Central Faculty, Scotland's Rural College, Edinburgh EH9 3JG, UK.
23
24

25 8 ^b Crop Science Department, Oak Park Crops Research Centre, Carlow, Ireland.
26
27

28 9 ^c Centre for Agricultural Strategy, School of Agriculture, Policy and Development, University of
29
30 10 Reading, Reading RG6 6AR, UK.
31
32

33 11 ^d Centre for Agri-Environmental Research, School of Agriculture, Policy and Development, University
34
35 12 of Reading, Reading RG6 6AR, UK.
36
37

38 13 ^e Agri-Food Sciences Division, Agri-Food and Biosciences Institute, 18a Newforge Lane, Belfast BT9
39
40 14 5PX, UK.
41
42

43 15 ^f Horticultural Development Department, Teagasc, Ashtown, Dublin 15, Ireland.
44
45

46 16 ^g Agricultural Economics and Farm Surveys Department, Teagasc, Ashtown, Dublin 15, Ireland.
47
48

49 17 *Corresponding author Email: henry.creissen@sruc.ac.uk Phone: +44 131 535 4119
50
51

52 18
53
54
55
56
57
58
59
60

19 **Abstract**

20 **BACKGROUND:** The impetus to adopt integrated pest management (IPM) practices has re-
21 emerged in the last decade, mainly as a result of legislative and environmental drivers.
22 However a significant deficit exists in the ability to practically monitor and measure IPM
23 adoption across arable farms; therefore the aim of the project reported here was to establish a
24 universal metric for quantifying adoption of IPM in temperate arable farming. This was
25 achieved by: (a) identifying a set of key activities that contribute to IPM; (b) weighting these
26 in terms of their importance to the achievement of IPM using panels of expert stakeholders in
27 order to create the metric (scoring system from 0-100 indicating level of IPM practiced); (c)
28 surveying arable farmers in the UK and Ireland about their pest management practices; and (d)
29 measuring level of farmer adoption of IPM using the new metric.

30 **RESULTS.** This new metric was found to be based on a consistent conception of IPM between
31 countries and professional groups. The survey results showed that, while level of adoption of
32 IPM practices varied over the sample, all farmers had adopted IPM to some extent (minimum
33 27.2 points, mean score of 65.1), but only 13 of 225 farmers (5.8%) had adopted more than
34 85% of what is theoretically possible, as measured by the new metric.

35 **CONCLUSION.** We believe that this new metric would be a viable and cost-effective system
36 to use to facilitate the benchmarking and monitoring of national IPM programmes in temperate
37 zone countries with large scale arable farming systems.

38 **Keywords:** Integrated Pest Management, IPM metric, IPM score, arable farming, farmer
39 survey, sustainable agriculture.

1. Introduction

Significant increases in crop production over the last century have resulted primarily from advancements in crop agronomy, including crop protection and nutrition, plant breeding and mechanisation of husbandry practices. These advances have largely been predicated on intensive use of inorganic chemical inputs, including fertilisers for plant nutrition and pesticides (collectively fungicides, herbicides, insecticides, molluscicides and nematicides) for crop protection. At the present time, commercial agriculture globally remains dependent on continued use of these synthetic crop protection products to prevent significant crop losses from pests.¹ Furthermore, prophylactic application of pesticides has become common across many intensive crop production systems.² However, input-intensive agriculture can and does result in unwanted consequences, including, adverse impacts on human and environmental health, development of pesticide resistance, all of which potentially reduce the sustainability of these systems.^{3,4} With increasing awareness of these risks, there is now widespread acknowledgement for the need to move towards more sustainable methods of agricultural production. One such method, which was first proposed in the 1950s,⁵ but that has gained significant traction and political support in recent years, is integrated pest management (IPM). IPM is regarded by many as a necessity for ensuring the optimum control of pests in an economically and environmentally sustainable manner.⁵⁻¹⁰ Whilst the precise definition of IPM can vary between studies and stakeholders,¹¹⁻¹³ it can broadly be categorised as an approach that considers the crop, the production system, the target pest(s) and their potential risks to production, as a whole system. IPM simultaneously employs multiple pest-control solutions, targeting different parts of this system, as a means to minimise the use of pesticides and ensure the long-term sustainability of pest control measures^{10,14,15}.

1
2
3 63 Whilst IPM can be readily understood in terms of such generalised statements and objectives,
4
5 64 the diversity of pest control practices that exist across all scales of an individual production
6
7
8 65 system makes identifying a definitive set of IPM practices extremely difficult. European Union
9
10 66 Directive 2009/128/EC, on the sustainable use of pesticides, which requires each member state
11
12 67 to encourage the use of IPM, identifies eight principles of IPM and a number of specific crop
13
14 68 management activities within each.⁷ These eight principles (Table 1) have been further
15
16 69 expanded upon by Barzman et al.¹⁰ to provide the basis from which IPM can be approached by
17
18 70 all those involved in crop production. However, as Barzman et al.¹⁰ concede, even with this
19
20 71 level of specification, it is difficult to provide a definitive checklist of IPM practices, or even
21
22 72 recommendations for approaches to implementing the eight principles. However, most
23
24 73 commentators would agree that the over-arching principle must be preventing or suppressing
25
26 74 the pest as opposed to intervening after the pest has become established, and that the
27
28 75 implementation of each of the eight principles should involve a continual process of
29
30 76 management plan redesign, implementation and evaluation.^{10,15} The perceived difficulties
31
32 77 associated with quantifying adoption of IPM practices has influenced some countries, such as
33
34 78 Denmark, to rely on pesticide usage as a proxy. In Denmark a pesticide tax system, which is
35
36 79 based on the wider impacts of pesticide use, is employed in an attempt to encourage adoption
37
38 80 of IPM practices. However, approaches to encourage adoption of IPM that rely heavily on a
39
40 81 single measure, such as pesticide usage, do not account for differences in the need for pesticides
41
42 82 between different cropping systems experiencing different pest challenges.

43
44
45 83 In arable farming there are a range of fairly ubiquitous crop management practices that are
46
47 84 consistent with these eight principles of IPM, but which are simply understood as good
48
49 85 husbandry. It is reasonable to assume, therefore, that most arable farmers may already be
50
51 86 practicing some form of IPM^{16,17} even when they do not appreciate that fact. Arable famers
52
53 87 tend to adopt IPM practices, in part, or incrementally over time by assessing the impacts of

1
2
3 88 individual components and slowly building up to a fully integrated approach to pest control at
4
5 89 which point complementarities between components occur.^{18,19} The practice of IPM, therefore,
6
7
8 90 can be seen as a continuum, with some farmers further than others along that continuum to
9
10 91 complete adoption.²⁰ Previous attempts to establish a metric of IPM, which would allow for an
11
12 92 assessment of where individual farmers lie on this continuum, have failed, largely due to the
13
14 93 lack of an objective approach to assessing the relative importance of different IPM components
15
16 94 and a lack of involvement of IPM practitioners in further pursuing the development of such a
17
18
19 95 metric.²¹⁻²³

20
21
22
23 96 Before informed efforts to increase the adoption of IPM in arable production systems can be
24
25 97 successfully implemented, accurate information on the current level of IPM practiced across a
26
27 98 diversity of systems is required. To achieve this, it is a necessity to be able to place both
28
29 99 individual farms and farm system typologies on some agreed metric of IPM. Such a metric
30
31
32 100 requires two attributes: first, it must capture a core of IPM management activities, based on a
33
34 101 consensus about what these are; and, second, the metric, must be able to use information on
35
36 102 adoption of these activities to create a continuum of degree of IPM adoption. Hence, the IPM
37
38 103 metric must be defined in terms of low-order, specific, actionable management activities. As
39
40 104 such, it must be a compound measure, capturing multiple IPM management activities
41
42
43 105 simultaneously.

44
45
46
47 106 The over-arching goal of this study therefore was to design and test a compound metric of IPM
48
49 107 with sufficient flexibility to be applied to a variety of farm situations and with sufficient
50
51 108 resolution to capture the continuum of degree of IPM adoption in a meaningful way. To reach
52
53 109 this goal, the achievement of a number of sub-objectives are necessary: identify the main IPM
54
55 110 activities that can be carried out on temperate arable farms; use stakeholder views to weight
56
57
58 111 these activities based on their relative contribution to achieving IPM; construct a composite
59
60

1
2
3 112 IPM measure based on these activities; test the efficacy of the IPM measure on a representative
4
5 113 sample of arable farms in England, Scotland, Northern Ireland and Ireland; validate the outputs
6
7
8 114 of the measure; and identify potential drivers of IPM involvement from among the
9
10 115 sociodemographic data collected from each participating farm business. Finally, some
11
12 116 conclusions are drawn of prime relevance to both practice and policy.
13
14
15

16 117 **2. Methods**

17
18
19
20 118 The study used a multi-stage process to achieve the objectives set out above, as outlined in
21
22 119 Figure 1; Tasks 1-3: design, optimisation and piloting of the data collection instrument; Task
23
24 120 4: data collection; Task 5-6: developing the IPM metric; Task 7: use, validation and secondary
25
26 121 analysis of the IPM metric.
27
28
29

30 122 **2.1. Design, piloting and optimisation of the farmer survey (Tasks 1-3)**

31
32
33 123 Following a review of the IPM literature, including the general principles of IPM as outlined
34
35 124 by the EU's Sustainable Use Directive 2009/128/EC (Table 1), a list of IPM practices
36
37 125 associated with temperate arable agriculture was identified. This list was then used to inform
38
39 126 the design of a farmer questionnaire to record level of involvement with IPM. The farmer
40
41 127 questionnaire consisted of three types of question, first, questions which captured information
42
43 128 on farmer engagement with specific IPM activities; second, farm and farmer sociodemographic
44
45 129 information; and finally, information in farmer attitudes towards, and perceptions of, IPM.
46
47 130 Questions were a mix of: multiple choice, 5-point rating scales, and some open-ended
48
49 131 questions, as appropriate to the type of information being elicited. The draft questionnaire was
50
51 132 tested via two rounds of piloting with farmers, agronomists and arable researchers. Following
52
53 133 the pilot, the number of questions was reduced from 44 to 22 by removing questions that proved
54
55 134 too complicated to answer fully, and combining questions to reduce repetition in the survey.
56
57
58
59
60

1
2
3 135 The final questionnaire contained a total of 22 questions; nine questions relating to individual
4
5 136 IPM activities, based on the eight principles of IPM (Table 1), and a further five questions
6
7
8 137 relating to perceptions of IPM. The remaining eight questions collected sociodemographic
9
10 138 information (a copy of the questionnaire is available in Supplementary Materials; Appendix 1).
11
12 139 On average the questionnaire took 10-15 minutes to complete. To protect against any biases
13
14 140 that farmers may have concerning IPM, whether these be positive or negative, the survey was
15
16 141 described as addressing best pest management practice in arable farming generally, rather than
17
18 142 IPM specifically.

143 **2.2. Farmer survey sampling strategy (Task 4)**

144 Arable farmers were selected for interview at random using national datasets as a sampling
145 frame in each of the four study countries. Each national research partner was set a target of
146 collecting 50 completed responses to ensure a sufficient number of responses for robust
147 statistical analysis within each country. All responses were collected between 2016 and 2017.
148 Data collection was by face-to-face interviews in England, Northern Ireland and Ireland, these
149 being carried out by experienced farm data recorders, while data collection in Scotland was by
150 means of a postal questionnaire.

151 **2.3. Developing a metric for the adoption of IPM on temperate arable farms (Tasks 5-6)**

152 The raw data collected from the survey on levels of adoption of each of the activities
153 contributing to IPM contained no indication of the relative importance of these individual
154 activities towards IPM. This weighting information was derived from a panel of industry
155 stakeholders, all of whom are actively involved in the practice of pest management in arable
156 crops, using a two-stage Delphi style approach. The Delphi technique uses data from a panel
157 of informed people and builds this data, using an iterative process, towards a consensus. The

1
2
3 158 strength of the technique lies in the fact that at each iteration the stakeholders have the
4
5 159 opportunity to amend their original judgements in light of the data and arguments supplied by
6
7
8 160 others.²⁴ In the first stage, a consultation with 11 stakeholders in Ireland was held in the form
9
10 161 of a workshop (see Table 2 for details). At the workshop, stakeholders were given a guidance
11
12 162 document (Supplementary material; Appendix 2) and a copy of the farmer questionnaire
13
14 163 (Supplementary material; Appendix 1) in addition to a verbal explanation of the project and
15
16
17 164 the aims and structure of the meeting.

18
19
20 165 The stakeholders were then asked to weight each of the six pre-selected questions relating to
21
22 166 adoption of IPM in the farmer questionnaire, on the basis of the importance of the pest
23
24 167 management activity that it captured, for IPM as a whole. The weighting process was
25
26
27 168 undertaken in two parts. First, where questions had sub-components, i.e. the question captured
28
29 169 multiple activities of a certain type, stakeholders were asked to provide ranks on a 1-5 scale for
30
31
32 170 these. The ranks were generated through an open discussion of the relative importance of each
33
34 171 sub-component to the question as a whole. Discussion continued until a consensus was reached
35
36 172 around a rank score. Second, each question was then awarded a weight based on its importance
37
38
39 173 to IPM. This involved allocating a total of 100 points over all six questions to decide on the
40
41 174 percentage contribution each question made to the overall IPM score. All six questions were
42
43 175 then combined, after applying the appropriate question weights, and divided by five (each
44
45 176 measure in the composite represents a 5-point scale) to form a composite Likert-type rating
46
47
48 177 scale²⁵ with a 100-point range representing level of IPM uptake, i.e. the IPM score.

49
50
51 178 The provisional set of weights derived from this workshop were then presented to a larger
52
53 179 stakeholder panel (see Table 2 and Table 3), by email and postal surveys across the study
54
55 180 countries in late 2017. Stakeholders were targeted in attempt to gather responses from those
56
57
58 181 actively involved in the practice of IPM. A total of 174 surveys were distributed and 46
59
60

1
2
3 182 responses collected. The group of stakeholders were informed of the original provisional set of
4
5 183 weights and then asked to provide their own estimate for each weight. To ensure consistency
6
7
8 184 of approach between participants in these surveys, each was sent a guidance document,
9
10 185 providing instructions for completing the questionnaire (Supplementary material; Appendix 3).
11
12 186 A total of 46 responses were collected and weights collected used on a one vote per stakeholder
13
14 187 basis. All data collected for this study, i.e. from the stakeholder workshop, and stakeholder and
15
16 188 farmer surveys were transcribed into electronic datasets and checked for errors (survey data
17
18 189 input file can be found in Supplementary material; Appendix 4). All statistical analyses of the
19
20 190 survey data were undertaken using the data analytics package SAS version 9.4.²⁷
21
22
23
24

25 191 **2.4. Validation and secondary analysis of the IPM metric (Task 7)**

26
27
28 192 A probability-probability plot was used to determine whether the distribution of the composite
29
30 193 IPM variable was normal or otherwise before other statistical operations were performed.
31
32 194 Cronbach's alpha was used to test for internal consistency in the composite IPM measure, by
33
34 195 measuring the inter-correlation between items, where an Alpha score of >0.7 is assumed to
35
36 196 indicate that the component questions cohere, i.e. they co-vary together.
37
38
39
40

41 197 To test the extent to which there is a common understanding of IPM between countries and
42
43 198 subgroups of stakeholders, the weights awarded by these different stakeholder groups were
44
45 199 compared. For the purposes of the analysis four different classes of stakeholder were identified:
46
47 200 Farmers, Independent agronomists (defined as those who do not directly benefit financially
48
49 201 from pesticide sales), Merchant agronomists, and Others. As degrees of freedom in some of
50
51 202 these groups were low, t-test comparisons were performed with aggregated stakeholder groups,
52
53 203 i.e. Farmer + Independent agronomist + Merchant agronomist compared with 'Other', where
54
55 204 'Other' represents researchers, agricultural college educators and policy makers. The rationale
56
57 205 for dividing the stakeholders into these two groups is that the first group comprises those that
58
59
60

206 have a commercial interest in pest management, whilst the ‘Other’ group are unlikely to have
207 such commercial interest.

208 To test for country understanding of IPM, an ANOVA was carried out to determine whether
209 stakeholders from the different countries applied different weights to the questions. For the
210 purposes of this analysis, Northern Ireland was combined with the data for Ireland due to low
211 observation numbers for Northern Ireland and the authors’ perception of cultural and
212 agricultural similarities between the countries.

213 **3. Results**

214 **3.1. The farmer survey sample**

215 A total of 225 responses were collected for the farmer survey: Northern Ireland (71), Ireland
216 (58), England (53), Scotland (43) (Table 4). The majority of respondents to the survey were
217 owners of the farm businesses. Farms varied considerably in size between countries, with the
218 largest farms found in Scotland, with an average size of 362 ha, and the smallest in the Ireland,
219 with an average farm size of 101 ha. 67% of the land on farms in the England sample was
220 arable land, with the remainder being improved grassland. Land cover on farms in the other
221 countries was much more heterogeneous, with smaller percentages of arable land and more
222 grassland (Table 4).

223 **3.2. The weighting of the components of the IPM metric**

224 The final set of weights provided by stakeholders for each question is outlined in Table 5,
225 whilst the final weights for sub-elements within each question are available in Appendix 3 of
226 the supplementary material. Overall, the weights awarded by stakeholders at the workshop
227 differed from the final stakeholder weights by between 1.6% and 17.8% for all questions, with

1
2
3 228 the exception of question 5, where the variation was 75.4%. However, there was an inverse
4
5 229 relationship between the absolute size of the weights and the percentage variation, with the
6
7
8 230 bigger weights showing the smallest variation (Table 5). Question 8, which focussed on
9
10 231 activities designed to prevent weeds, disease and insects/molluscs, was judged to account for
11
12 232 47% of the achievable IPM score, with factors influencing pest management plans (Question
13
14
15 233 9) coming in second, with a relative contribution of 15% to the IPM score.
16
17

18 234 **3.3. Validation of the IPM metric**

19
20
21 235 Farmers IPM scores were relatively normally distributed (Supplementary materials; Appendix
22
23 236 5) with a range of 27.2 – 91.3, a mean of 65.1 and a standard deviation of 13.8 (Coefficient of
24
25 237 Variation 21%), i.e. exhibiting a normal bell-shaped curve, although the distribution is
26
27
28 238 somewhat skewed towards higher IPM scores (Figure 2), suggesting that the majority of
29
30 239 farmers are already implementing at least some measures that would be seen as characteristic
31
32
33 240 of IPM. However, while all farmers are practicing some of level of IPM, only 13 out of 225
34
35 241 farmers (5.8%) scored more than 85 on a possible scale of 100. Any responses containing a
36
37 242 high amount of unanswered questions, leading to a score of less than 20, would have been
38
39 243 removed from the survey but none of our respondents fell into that category.
40
41
42
43
44 244

45 46 47 245 **3.4. The coherence of the IPM metric**

48
49
50 246 Cronbach's Alpha was used to test the coherence of the questions combined, after weighting,
51
52 247 to create the IPM metric. There are no hard and fast rules about what Alpha value is required
53
54 248 to show adequate coherence in the sets of measures used to form composites, but it is widely
55
56
57 249 held that the higher the value the better (although extremely high values might be suggestive
58
59 250 of redundancy among the measures). A precedent has become established that Cronbach's
60

1
2
3 251 Alpha values in excess of 0.7 are classed as good or better, so this convention and threshold
4
5 252 has been followed here.²⁶ When undertaking this testing of the internal consistency of the
6
7 253 composite IPM measure, the three components of Question 8, relating to the choice of pest
8
9 254 prevention measures, were separately tested for coherence. Each of these three questions, a)
10
11 255 measures for prevention of weeds; b) measures to prevent diseases; and c) measures to prevent
12
13 256 insect pests/nematodes/molluscs, was itself a composite measure, made up of a number of sub-
14
15 257 components. With the exception of Question 8b – ‘What measures are used to control diseases’,
16
17 258 all questions had a Cronbach’s Alpha >0.7 (Table 6) strongly suggesting that these composite
18
19 259 questions were also coherent. Question 8b had a Cronbach’s Alpha of 0.68. While obviously
20
21 260 falling below the 0.7 threshold set above, the literature suggests that this value is still
22
23 261 acceptable.²⁶ Overall these results suggest a high degree of internal consistency in the
24
25 262 composite IPM measure.

263 **3.5 A consensus as to what constitutes IPM practice across groups and countries**

264 For four of the six questions, there were no significant differences in the weights attributed by
265 the two stakeholder consultation classes (Table 7). This lack of significant difference between
266 the weights awarded by the two stakeholder classes indicates that there was a consensus on
267 what constitutes IPM practice. For two questions, there were significant differences between
268 the two stakeholder classes; Q5 – ‘What influences your choice of cereal variety?’, and Q14 –
269 ‘Membership of an agronomy/crop discussion group’. However, these two questions only
270 account for, a combined weight of, 14.6% of the overall IPM score. In both instances, those
271 stakeholders with commercial interest in pest management (farmers and agronomists) weighted
272 the questions higher in importance than those stakeholders that are commonly considered to
273 have no commercial interest (researchers, regulators, educators), with Q5 seeing a 69%
274 increase in weight and Q14 a 78% increase.

1
2
3 275 No statistically significant differences amongst countries in the weights applied by stakeholders
4
5 276 to any of the IPM questions were identified (Table 8). In the case of Q4 – ‘Why do you typically
6
7 277 use an arable rotation?’ differences were close to being significant ($P=0.07$), but this is a
8
9 278 question with a relatively small weighting. The national differences in weight on this question
10
11 279 occurred between England (12.1%) and Scotland (9.9%), and also the island of Ireland (14.0%)
12
13
14 280 and Scotland.

18 281 **4. Discussion and conclusion**

22 282 IPM is a knowledge-intensive process in which scientifically proven measures are selected for
23
24 283 use, based on the specific set of biotic threats affecting the crop and the financially viable
25
26 284 approaches available to the grower, to reduce risk associated with these threats.^{10,28} As such
27
28 285 IPM does not necessarily rely upon individual control mechanisms in isolation but seeks to use
29
30 286 a complexity of inter-related strategies. It is this complexity that makes capturing levels of IPM
31
32 287 practiced at the farm scale difficult. This difficulty can be further compounded by unintentional
33
34 288 perspective bias, for example relating to what does and does not constitute an IPM activity,
35
36 289 imposed by observers via both the methods used to collect such data and the assessment
37
38 290 process.²³ Such bias can result in the exclusion of activities which legitimately contribute to
39
40 291 IPM being from the survey and other IPM activities being attributed irrational weights.

46 292 The development of a metric to assess the extent of adoption of IPM described here differs in
47
48 293 approach from those currently in existence. This is because the generation of the weighting
49
50 294 system for the various elements of IPM was, in this case, rigorous, and involved a number of
51
52 295 IPM practitioners from various professional backgrounds. Many previous attempts to develop
53
54 296 such a metric have not been able to garner widespread support due to the fact that the process
55
56 297 of determining which activities to include in the metric and the weights attributable to these
57
58 298 has remained solely in the hands of researchers, with little or no reference to industry

1
2
3 299 stakeholders such as farmers and professional agronomists.²¹⁻²³ The carefully controlled
4
5 300 approach to developing a metric for IPM reported here, together with the observed clear within-
6
7 301 sector and between-country consensus about what constitutes IPM suggests that this IPM
8
9 302 metric has potential for use in an international context.

10
11
12
13 303 Currently, gaps exist between farmer perception of the value of IPM and their actual practice.
14
15 304 Whereas farmer attitudes towards IPM are often positive, the practicalities and perceived
16
17 305 financial implications associated with IPM adoption can act as barriers.^{29,30} Gaps may also exist
18
19 306 between actual and perceived practices i.e. farmers may believe they are practicing IPM when
20
21 307 in reality they are not, and vice versa.^{16,31} Such a phenomenon may have contributed to
22
23 308 differences between weightings awarded to certain questions at the workshop, and then later at
24
25 309 the stakeholder survey. Whilst the perceived importance of the majority of questions as
26
27 310 contributors to IPM were not viewed differently amongst the different stakeholder groups,
28
29 311 differences did exist for perceived importance of some of the lesser contributors, i.e. factors
30
31 312 influencing variety choice and membership of discussion groups. Likewise, with the exception
32
33 313 of the question relating to cereal rotations (Question 4), stakeholders from the different
34
35 314 countries ranked questions in equal importance. Stakeholders from the island of Ireland and
36
37 315 England considered the question on rotations to be relatively more important for IPM than did
38
39 316 stakeholders from Scotland. This could be due to the dominance of spring barley in the Scottish
40
41 317 arable sector. With a single, premium crop dominating the market, alternative suitable cropping
42
43 318 options are potentially reduced and, thus, growing different crops in rotation may not be
44
45 319 considered a viable option. Regarding the other differences between the two stakeholder
46
47 320 groups, those stakeholders who have a commercial interest in pest management weighted
48
49 321 discussion group membership as being more important for IPM than the stakeholder group who
50
51 322 are unlikely to have a commercial interest in pest management. This indicates that they
52
53 323 recognise a greater value in this form of knowledge exchange which may lead to an increase
54
55
56
57
58
59
60

1
2
3 324 in adoption of IPM practices. There were also differences in the weights awarded to the
4
5 325 question on varietal selection, with stakeholders who have a commercial interest in pest
6
7 326 management weighting the question more important for IPM. Selecting varieties based on their
8
9
10 327 disease resistance rating, in particular, has long been promoted as a major tool for disease
11
12 328 management. Scottish barley growers have claimed to select and grow disease resistant
13
14 329 varieties, yet on consulting the Agriculture and Horticulture Development Board (AHDB)
15
16 330 disease resistance ratings, the varieties grown are often rated much weaker than others for the
17
18 331 major disease threats such as rhynchosporium.³¹ This finding was mirrored in the survey of UK
19
20 332 growers undertaken by ADAS consulting limited.¹⁶ The lower weighting awarded to this
21
22 333 question at our expert workshop could have been, in part, due to discussions involving
23
24 334 stakeholders in Ireland who may consider themselves restricted in their choice of varieties due
25
26 335 to a lack of suitability of commercial cultivars to Irish growing conditions owing to an absence
27
28 336 of cereal breeding programmes in Ireland. The fact that these differences were not observed
29
30 337 when investigating differences in weighting between the stakeholders from different countries
31
32 338 may suggest that this may, in fact, be an artefact of the consultation method. As such only the
33
34 339 aggregate score from the survey panel of stakeholders was used in the creation of the IPM
35
36 340 metric.

37
38
39
40
41
42
43 341 By combining the targeted survey of arable farms with a stratified sampling method and the
44
45 342 consensual development of a metric to capture IPM in arable production systems, it is
46
47 343 anticipated that current levels of IPM adoption, and perception of it, in both the UK and Ireland
48
49 344 can be determined and if the survey were to be repeated changes in adoption could be tracked.
50
51 345 If the barriers or, indeed, the limitations of IPM in such systems are to be identified, this is a
52
53 346 key step in the process. Although all respondents were considered to be practicing IPM (of the
54
55 347 225 respondents all scored >20 of a total score of 100), a wide range of scores within a broadly
56
57 348 normal distribution was recorded. This distribution opens the possibility of identifying such

1
2
3 349 barriers/limitations to further adoption. This process could be further enhanced by including,
4
5 350 in subsequent analysis of drivers and barriers, various questions relating to IPM perception
6
7 351 and/or socioeconomic data. As the data set obtained for both Ireland and England contained
8
9 352 official national farm business survey statistics, it may also be possible to delve further into
10
11 353 financial components of the farm enterprise that may directly or indirectly influence IPM
12
13 354 practice.
14
15
16
17

18 355 The applicability of the metric to arable farming in other temperate zone countries is as yet,
19
20 356 unknown. However, it is foreseeable that the metric and the phenomena it captures will be
21
22 357 relevant elsewhere. Using the approach reported here, modification of the metric, by re-
23
24 358 weighting questions based on expert opinion, according to the challenges and opportunities for
25
26 359 IPM in each country, may render the metric widely applicable. This would result in a locally-
27
28 360 weighted IPM metric approach. Furthermore, the process by which the survey and metric were
29
30 361 developed can be easily adapted to cover additional crops and cropping systems requiring
31
32 362 different approaches to pest control.
33
34
35
36
37

38 363 Despite a considerable body of legislation relating to pesticide practice and use, both nationally
39
40 364 and at EU level, there has been, to date, no agreed upon metric available that would allow the
41
42 365 measurement of the effectiveness of IPM at reducing pesticide usage or increasing adoption of
43
44 366 sustainable crop protection methods. The study reported here provides a novel and useful
45
46 367 metric to assess the extent of adoption of IPM practices and the possible development of a
47
48 368 sustainable plant protection system for arable cropping in temperate climates.
49
50
51
52

53 369

54
55
56 370 **Acknowledgements**
57
58
59
60

1
2
3 371 We acknowledge the financial and logistical support of the: Scottish Government Strategic
4
5 372 Research Programme; Rural Business Research (England); Department of Agriculture, Food
6
7 373 and the Marine (Ireland) (RSF 14/S/879); and the Department of Agriculture, Environment and
8
9 374 Rural Affairs (Northern Ireland) in enabling the research reported here. We thank all who
10
11 375 contributed to the design of the survey and all involved in data collection processes including
12
13 376 the Teagasc National Farm Survey recorders, Farm Business Survey recorders, Science and
14
15 377 Advice for Scottish Agriculture, Census and Single Farm Payment data recorders. We also
16
17 378 thank the stakeholders involved in developing the IPM metric and all the farmers who
18
19 379 participated in the survey.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **380** **References**
4
5

- 6
7 381 1 Oerke E, Crop losses to pests. *The Journal of Agricultural Science* **144**:31-43 (2006).
8
9 382 2 Cooper J, Dobson H, The benefits of pesticides to mankind and the environment. *Crop Prot.*
10
11 383 **26**:1337–1348 (2007).
12
13 384 3 Goulson D, Thomson J, Croombs A, Rapid rise in toxic load for bees revealed by analysis of
14
15 385 pesticide use in Great Britain. *PeerJ* **6**:e5255 (2018).
16
17 386 4 Lamichhane JR, Dachbrodt-Saaydeh S, Kudsk P, Messean A, Toward a reduced reliance on
18
19 387 conventional pesticides in European agriculture. *Plant Dis.* **100**:10-24 (2016).
20
21 388 5 Stern V, Smith R, van den Bosch R, Hagen K, The integrated control concept. *Hilgardia* **29**:81–
22
23 389 101 (1959).
24
25 390 6 European Union, Directive 2009/127/EC of the European Parliament and of the Council of 21
26
27 391 October 2009 amending Directive 2006/42/EC with regard to machinery for pesticide application.
28
29 392 *Official Journal of the European Union* **52**: 29–33 (2009a).
30
31 393 7 European Union, Directive 2009/128/EC of the European parliament and of the council of 21
32
33 394 October 2009 establishing a framework for community action to achieve the sustainable use of
34
35 395 pesticides. *Official Journal of the European Union* **52**:71–86 (2009b).
36
37 396 8 European Union, Regulation (EC) No 1107/2009 of the European parliament and of the council
38
39 397 of 21 October 2009 concerning the placing of plant protection products on the market and repealing
40
41 398 council directives 79/117/EEC and 91/414/EEC. *Official Journal of the European Union* **52**:1–50
42
43 399 (2009c).
44
45 400 9 European Union, Regulation (EC) No 1185/2009 of the European Parliament and of the Council
46
47 401 of 25 November 2009 concerning statistics on pesticides. *Official Journal of the European Union*
48
49 402 **52**:1–22. (2009d).
50
51 403 10 Barzman M, Barberi P, Birch ANE, Boonekamp P, Dachbrodt-Saaydeh S, Eight principles of
52
53 404 integrated pest management. *Agron. Sustainable Dev.* **35**:1199-1215 (2015).
54
55 405 11 Prokopy R., Kogan. M., 2003. Integrated pest management, in Encyclopedia of Insects, ed. by
56
57 406 Resh VH and Carde RT. Academic Press, San Diego, pp. 589–595 (2003).
58
59
60

- 1
2
3 407 12 Kogan M, Bajwa WI, Integrated pest management: A global reality? *An. Soc. Entomol. Bras.*
4
5 408 **28**:1–25 (1999).
6
7 409 13 Jeger M, Bottlenecks in IPM. *Crop Prot.* **19**:787–792 (2000).
8
9 410 14 Kogan M, Integrated pest management: Historical perspectives and contemporary
10 411 developments. *Annu. Rev. Entomol.* **43**:243–270 (1998).
11
12 412 15 Ehler LE, Integrated pest management (IPM): Definition, historical development and
13 413 implementation, and the other IPM. *Pest Manage. Sci.* **62**:787–789 (2006).
14
15 414 16 ADAS, The Awareness, Use and Promotion of Integrated Crop & Pest Management Amongst
16 415 Farmers and Growers, a Survey on Behalf of DEFRA and the CPA (2002).
17
18 416 17 Doonan H, Adoption of Integrated Pest Management in the UK arable sector. *Outlooks Pest*
19 417 *Manage.* **28**:29-32 (2017).
20
21 418 18 Lamine C, Transition pathways towards a robust ecologization of agriculture and the need for
22 419 system redesign cases from organic farming and IPM. *Journal of Rural Studies* **27**:209–219 (2011).
23
24 420 19 Zepeda JF, Barreto-Triana N, Falck-Zepeda J, Barreto-Triana N, Baquero-Haeberlin I, Espitia-
25 421 Malagón E, Fierro-Guzmán H, López N, An exploration of the potential benefits of integrated
26 422 pestmanagement systems and the use of insect resistant potatoes to control the Guatemalan tuber
27 423 moth (*Tecia solanivora* Povolny) in Ventaquemada, Colombia. International Food Policy Research
28 424 Institute, Washington, DC (2006).
29
30 425 20 Benbrook CM, Groth E, Halloran JM, Hansen M, Marquardt S, Pest management at the
31 426 crossroads. Consumers Union, Yonkers, NY, 272 pp (1996).
32
33 427 21 Shennan C, Cecchetti CL, Goldman GB, Zalom FG, Profiles of California farmers by degree
34 428 of IPM use as indicated by self descriptions in a phone survey. *Agric., Ecosyst. Environ.* **84**:267–
35 429 275 (2001).
36
37 430 22 Robertson MJ, Zehnder GW, Hamming MD, Adoption of integrated pest management practices
38 431 in South Carolina cotton growers. *Journal of Extension* **43** (2005).
39
40 432 23 Puente M, Darnall N, Forkner RE, Assessing Integrated Pest Management Adoption:
41 433 Measurement Problems and Policy Implications. *Environ. Manage.* **48**:1013–1023 (2011).
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 434 24 Hsu CC, Sandford BA, The Delphi technique: making sense of consensus. *Practical Assessment.*
4
5 435 *Research and Evaluation* **12**: 1-8 (2007).
6
7 436 25 Likert R, A technique for the measurement of attitudes. *Archives of Psychology* **140**: 1-55 (1932).
8
9 437 26 Taber KS, The use of Cronbach's alpha when developing and reporting research instruments in
10
11 438 science education. *Research in Science Education*, 48(6), 1273–1296. (2018).
12
13 439 27 SAS Institute Inc., Cary, North Carolina, USA.
14
15 440 28 Byerlee D, Modern varieties, productivity, and sustainability—recent experience and emerging
16
17 441 challenges. *World Development* **24**:697–718 (1996).
18
19 442 29 Sherman J, Gent D, Concepts of sustainability, motivations for pest management approaches,
20
21 443 and implications for communicating change. *Plant Dis.* **98**:1024-1035 (2014).
22
23 444 30 Royle DJ, Shaw MW, The costs and benefits of disease forecasting in farming practice. in:
24
25 445 Control of Plant Diseases: Costs and Benefits. eds. by Clifford BC and Lester E, Blackwell
26
27 446 Scientific Publications, London. Pp. 233-236 (1988).
28
29 447 31 Stetkiewicz S, Bruce A, Burnett FJ, Ennos RA, Topp CFE, Perception vs practice: Farmer
30
31 448 attitudes towards and adoption of IPM in Scottish spring barley. *Crop Prot.* **112**:96-102 (2018)
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1: The eight principles of IPM and their components as defined by the European Union, (2009b) and expanded by Barzmann et al. (2015). Also, the alignment of questions from the study questionnaire with each principle.

Principle	Description	Components	Survey Questions¹
1.	Prevention and suppression	Crop rotation, cultivation techniques, varietal resistance, phytosanitary measures, beneficial organisms	3, 5, 7, 8 (a, b, c), 9
2.	Monitoring	Field monitoring, forecasting, seeking expert advice	8 (a, b, c), 9, 14
3.	Informed decision making	Protection measures based on expert advice, action thresholds	8 (a, c) , 9,
4.	Non-chemical methods	Preference for biological and physical control methods over chemical.	3, 5, 7, 8 (a, b, c)
5.	Pesticide selection	Using pesticide that minimise negative effects on human health and the environment	8 (a, b, c)
6.	Reduced pesticide use	Reduced doses, reduced application frequency considering the risk for development of pesticide resistance	8 (a, b, c)
7.	Anti-resistance management	Alternation/mixing pesticides containing multiple modes of action.	9
8.	Evaluation	Assessment of the efficacy of control treatments used to inform future management decisions.	9

¹Only questions used to generate the IPM score are highlighted. The full questionnaire can be found in Supplementary materials.

Table 2: Number of participants in the initial stakeholder workshop and subsequent stakeholder consultation panel, their principal occupation and their stakeholder class.

¹Stakeholder workshop held on June 27th 2017 at Teagasc Oak Park, Ireland.

Principal Occupation	Stakeholder Workshop¹	Stakeholder Panel²	Stakeholder class for weighting analysis
Farmer	2	18	Farmer
Independent Agronomist	2	11	Independent Agronomist
Merchant agronomist	1	8	Merchant agronomist
Researcher	5	0	Other
Agricultural college lecturer	0	9	Other
Agricultural Regulator	1	0	Other
Total	11	46	

²Stakeholder panel completing the survey from which the final weighting were derived.

Table 3: Number of stakeholder panel participants from each country involved in construction of the IPM metric.

Country	Participants
England	11
Ireland	12
Northern Ireland	4
Scotland	19

Table 4: Overview of respondents to the survey and means of data collection in each of the participating countries.

	England	Scotland	Northern Ireland	Republic of Ireland
Method of data collection	Farm Business Survey recorders	Postal	Census and Single Farm Payment data recorders	Teagasc National Farm Survey recorders
Sample size	53	43	71	58
Percent owned	54.6	83.7	97.2	68.6
Farm size (ha)	202.19	361.5	109.2	101.07
Of which arable (ha)	135.32	198.2	59.1	63.6

Table 5: Relative contribution of each question (% weight) awarded by stakeholders at the workshop (n=11), the survey panel (n=46), and the final combined set.

Specific question within the survey	Weights produced at workshop	Final set of weights (produced by survey panel)	Variation from the workshop weights (%)
Q3. Proportion of land in continuous cereal production	10	11.46	14.6
Q4. Why do you typically use an arable rotation?	10	11.78	17.8
Q5. What influences your choice of cereals variety?	5	8.77	75.4
Q8. What preventive measures are used to control weeds, disease and insects/molluscs?	55	46.93	14.7
Q9. What factors do you consider when deciding on your pest management plan?	15	15.24	1.6
Q14. Membership of an agronomy / crop discussion group?	5	5.82	16.4
Total	100	100	

Table 6: Correlation of component questions with overall IPM score and Cronbach’s Alpha test.

Specific question within the survey	Correlation with total (standardised scores)	Alpha ¹
Q3. What proportion of land on your farm is in continuous cereals production?	0.412456	0.720713
Q4. Why do you typically use an arable rotation?	0.395048	0.724010
Q5. What influences your choice of crop variety?	0.407890	0.721580
Q8a. What preventive measures are used to control weeds	0.500479	0.703677
Q8b. What preventive measures are used to control diseases	0.602404	0.683182
Q8c. What preventive measures are used to control insects	0.471783	0.709298
Q9. What factors do you consider when deciding on your pest management plan?	0.362842	0.730049
Q14. Membership of an agronomy / crop discussion group?	0.343871	0.733569

¹ High Alpha scores (>0.7) for a specific question indicate a high correlation of that question with the overall score.

Table 7: Impact of stakeholder occupations on the specific weighting for each of the identified questions relating to IPM practice.

Specific question within the survey	Difference between groups ¹	T value	Variances	Pr>T
Q3. What proportion of land on your farm is in continuous cereals production?	-0.9535	-0.40	Equal	0.6943
Q4. Why do you typically use an arable rotation?	1.5240	0.76	Equal	0.4535
Q5. What influences your choice of crop variety?	2.9339	1.71	Unequal	0.0130
Q8. What preventive measures are used to control weeds, disease and insects/molluscs?	-4.6727	-1.16	Equal	0.2522
Q9. What factors do you consider when deciding on your pest management plan?	0.3784	0.25	Equal	0.8070
Q14. Membership of an agronomy / crop discussion group?	1.3303	1.71	Unequal	0.0041

¹Groups as per Table 2.

Table 8: Impact of stakeholder country of origin on the specific weighting awarded for each of the identified questions relating to IPM practice.

Specific question within the survey	F value	Pr > F	R-Square
Q3. What proportion of land on your farm is in continuous cereals production?	0.90	0.4128	0.040318
Q4. Why do you typically use an arable rotation?	2.76	0.0743	0.113889
Q5. What influences your choice of crop variety?	0.78	0.4631	0.035174
Q8. What preventive measures are used to control weeds, disease and insects/molluscs?	1.70	0.1942	0.073393
Q9. What factors do you consider when deciding on your pest management plan?	0.35	0.7067	0.016016
Q14. Membership of an agronomy / crop discussion group?	0.06	0.9397	0.002890

For Peer Review

1
2
3 **Figure 1.** Overall approach used to develop and validate the IPM metric, divided into seven tasks.
4

5 **Figure 2.** Distribution of sample by IPM score.
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

1
2
3 Supplementary materials.
4

5 **Appendix 1: Best arable farming practice survey**
6
7
8

9 This survey is part of a wider study which aims to improve farmer awareness and adoption of
10 Integrated Pest Management (IPM) in the UK and Ireland. IPM is widely regarded as offering benefits
11 to farmers in their efforts to control pests, including reducing pollution risk and costs.
12
13

14 The study objectives will be achieved by:
15

- 16 1. collection of information (through this survey) on current IPM practices in arable farming;
- 17 2. analysis of this information to identify best practice; and
- 18 3. transfer of this knowledge back to farmers and advisors through various knowledge transfer
19 activities, including publications in trade magazines.
20
21
22

23 The findings will be reported at a group level only i.e. data collected from individual growers,
24 discussion groups, facilitators or advisors will not be published. This study is not concerned with, and
25 does not collect data relating to, farm support payments, Cross Compliance activities or Agri-
26 Environment schemes. The survey should take between 10-15 minutes to complete and you should
27 not need to consult farm records.
28
29

30 All data supplied will be treated in the strictest confidence, will be used solely for the purposes of this
31 study, and will not be passed on to third parties. If you submit a completed survey form you can
32 withdraw at any time. If you should wish to withdraw please contact me using the contact information
33 provided on this letter. Please retain this letter for your records. This survey has passed all ethical
34 clearance procedures at the co-ordinating institution.
35
36
37

38 Thank you for participating in our survey.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Best arable farming practice survey

- Please answer the questions as accurately as you can. Good data is needed to provide reliable advice back to farmers and advisors.
- Please note that the term '**pests**' relates to **diseases, weeds and invertebrate pests** (insects and molluscs). Similarly, '**pesticides**' refers to **fungicides, herbicides, insecticides and molluscicides**.
- Please read question instructions carefully as the type of response required may vary from question to question.
- Please complete the survey in full.

1. How familiar are you with Integrated Pest Management (IPM)? Please tick one answer only.

- Not at all familiar (if this answer, please move direct to Question 3.)
- Somewhat unfamiliar
- Moderately familiar
- Familiar
- Very familiar

2. Which of the following factors do you consider to be important components of IPM? Please tick one box in every row.

	Very unimportant	Not important	Neither important or unimportant	Fairly Important	Very important
Preventative measures (hygiene practices such as cleaning equipment, sourcing clean seed etc.)					
Biological control methods (growing competitive crops, beetle banks etc.)					
Cultural control methods (altering drilling dates to reduce disease, increasing seeding rate to control weeds, rotating crops etc.)					
Monitoring and surveillance of insect pest, weed and disease levels (crop walking, reacting to high disease/pest pressure alerts etc.)					
Minimum use of pesticides					

3. What proportion of your land is in continuous cereal production i.e. growing cereals on the same land for 5 or more consecutive years without growing a break crop (e.g. oilseed rape, beans, peas, grass)? Please circle the relevant proportion below.

None	1 – 25%	26 – 50%	51 – 75%	76 – 100%
------	---------	----------	----------	-----------

4. Why do you practice continuous cereal production? More than one answer may be provided.

- I don't practice it
- Land unsuitable for other crops
- Climate unsuitable for other crops
- No access to machinery/equipment/storage facilities required to grow non-cereal crops
- Greater risks associated with growing different crops
- End-market requirements

5. **If you typically use an arable rotation, why do you do this?** *Please skip this question if you do not practice a rotation. Otherwise please answer every row.*

	Strongly disagree	Disagree	Neither agree or disagree	Agree	Strongly agree
To control weeds					
To control disease					
To control insect pests					
To improve and maintain soil structure and fertility					
To spread financial risks					
Largely because it's necessary to comply with regulations					

6. **Please indicate your tillage practice.** *More than one answer may be provided.*

- Regular ploughing
 Direct drilling (no tillage)
 Strip tillage
 Reduced (minimum) tillage
 Rotational ploughing (every few years)

7. **What influences your choice of cereal or oilseed variety grown?** *Please tick the factors you consider most important for each crop you grow. More than one answer may be provided.*

	Important factors
Recommended lists (where available)	
Availability of seed	
Advisor recommendation	
End-market	
Disease resistance	
Weed competitiveness	
Insect pest resistance	
Abiotic stress resistance (e.g. drought/flooding/cold tolerance)	
Yield potential	
Quality potential	
Consistency of performance	

8. **Which preventative measures do you currently employ to control the introduction and spread of pests on your farm?** *Please tick all the boxes below that apply.*

a) *To control weeds*

- Stale seedbeds
 Full inversion ploughing to control low dormancy weeds
 Min-till to control high dormancy weeds
 Pre-emergence herbicide applications
 Spot spraying weeds (if necessary)
 Hand rogueing/hoeing weeds
 Manage headlands differently to remainder of field
 Fields with high weed levels are harvested last

- Crop inspections, please indicate frequency; once or twice per (*please circle one of the following options*):

week	month	season
------	-------	--------

b) *To control disease*

- Grow resistant varieties
 Use certified seed
 Test non-certified seed and treat if required
 Regularly test soils for soil borne pathogens
 Use seed treatments to control disease
 Crop inspections, please indicate frequency; once or twice per (*please circle one of the following options*):

week	month	season
------	-------	--------

c) *To control insects, nematodes and molluscs*

- Encourage beneficial insects through planting habitat
 Avoid broad spectrum insecticides
 Use seed treatments
 Preparations for control of molluscs e.g. avoiding direct drilling or preparation of fine seed bed
 Regularly monitor above ground pest populations
 Set action thresholds
 Regularly test soils for nematodes
 Regularly test soils for insect pests
 Frequently clean harvesting and storage equipment
 Crop inspections, please indicate frequency; once or twice per (*please circle one of the following options*):

week	month	season
------	-------	--------

9. What factors do you consider when deciding on a pest management plan at the start of the season?

Please tick those that apply.

- I don't use pest management plans
 Crop walking data from last season, used to assess the performance of various control measures
 Technical research on pesticide product efficacy
 Weed maps, created and monitored for changes between seasons
 Yield maps, used to identify areas requiring specific attention
 Cost-benefit analysis of management options
 End-market requirements
 Variety resistance
 Soil borne pathogens
 Position in rotation
 Anti-resistance strategies
 None of the above
 Other (please specify) _____

10. What factors influence your decision to adjust your spray programme (e.g. changes in timings, rates, products) throughout the season? Please answer every row by ticking in one of the columns.

	No/low influence	Moderate influence	High influence
Growth stage of the crop			
Calendar date			
Weather conditions and forecasts			
Crop monitoring information, such as CropMonitor (where available)			
Predictions of Decision Support Systems (where available)			
Observed levels of pest presence in the field			
Advisor recommendation			
Actions of/advice from other farmers in the area			
None of the above, I operate a fixed spraying programme			

11. Name the specific pests (weeds, diseases, insects or molluscs) which you see as being of the greatest concern to crop production on your farm. Start with 1 = greatest concern.

Current threat		Future threat (5+ years' time)	
1.		1.	
2.		2.	
3.		3.	

12. How valuable are the following sources of pest (weeds, diseases, insects or molluscs) management advice?

Please rank the top 3 sources of pest advice starting with 1 being the most valuable. After listing the top 3, please leave the rest blank.

	Rank top 3 sources
Open days/crop walks	
Farmer discussion groups	
Other farmers (not including discussion groups)	
Independent agronomist	
Chemical company representative	
Merchant agronomist	
Contractors	
Past experience	
Farming press	
Other (please specify)	

13. Which of the following statements best describes your relationship with your main crop protection advisor? Please tick one only.

- I rely on them and act on their suggestions
- I tell them what I want from them and they respond to meet my wishes
- We decide on the pest management strategy together
- I listen to their advice but will always consult other sources of information
- I listen to their advice but adjust if needed when in the field
- I don't use an advisor

14. Are you a member of an agronomy or crop discussion group? Please tick.

- 1
2
3 Yes
4 No

5 *If Yes, please specify*
6 _____
7 _____
8 _____
9 _____

10 **15. What is your position on the farm? More than one box may be ticked.**

- 11
12
13 Owner
14 Tenant
15 Farm worker
16 Contractor
17
18
19

20 **16. How much land do you farm or manage?**

21 Total area: _____ hectares:

22 Of which – arable _____ hectares

23 grassland _____ hectares

24 rough grazing _____ hectares

25 fallow _____ hectares

26 biodiversity scheme _____ hectares

27 How much of the area farmed is rented/leased: _____ hectares

28 How much of the area farmed is shared: _____ hectares

29
30
31 **17. Which description best fits your farming type? Please tick one box only.**

- 32 Predominantly arable
33 Predominantly livestock
34 Mixed
35 Horticulture
36
37
38
39

40 **18. Your age. Please circle.**

41 20 - 30	42 31 – 40	43 41 - 50	44 51 – 60	45 61 and over
------------	------------	------------	------------	----------------

46
47
48 **19. Which qualifications have you achieved? Please tick one only.**

- 49 Certificate in farming (e.g. BASIS)
50 O levels/GCSEs/Junior certificate
51 A levels/Leaving certificate
52 Diploma
53 Bachelor's Degree
54 Higher degree (Master's degree/PhD)
55
56
57
58
59
60

1
2
3 **20. Does the main decision-maker of the farm business have an off-farm income source? Please**
4 *tick.*

5
6
7 Yes

8 No

9
10
11 **21. Is there a successor identified to take over the farm business? Please tick.**

12
13 Yes

14 No

15 Uncertain at present

16
17
18 **22. Are you involved in a scheme that promotes biodiversity e.g. Countryside Stewardship,**
19 **LEAF, AEOS, GLAS? Please tick.**

20
21 Yes

22 No

23 **Thank you very much for your help in our research.**

24
25 **The information you have provided will be treated in the strictest confidence by us.**

1
2
3 **Appendix 2: Guidance for stakeholders attending the workshop designed to create an initial**
4 **weighting system for IPM adoption.**
5

6 **Best arable farming practice survey**
7

8 This survey is part of a wider study which aims to improve farmer awareness and adoption of
9 Integrated Pest Management (IPM) in the UK and Ireland. IPM is widely regarded as offering benefits
10 to farmers in their efforts to control pests, including reducing pollution risk and costs.
11
12

13 The study objectives will be achieved by:

- 14 1. collection of information (through this survey) on current IPM practices in arable farming;
- 15 2. analysis of this information to identify best practice; and
- 16 3. transfer of this knowledge back to farmers and advisors through various knowledge transfer
17 activities, including publications in trade magazines.
18
19

20
21 The findings will be reported at a group level only i.e. data collected from individual growers,
22 discussion groups, facilitators or advisors will not be published. This study is not concerned with, and
23 does not collect data relating to, farm support payments, Cross Compliance activities or Agri-
24 Environment schemes. The survey should take between 10-15 minutes to complete and you should
25 not need to consult farm records.
26
27

28 All data supplied will be treated in the strictest confidence, will be used solely for the purposes of this
29 study, and will not be passed on to third parties. If you submit a completed survey form you can
30 withdraw at any time. If you should wish to withdraw please contact me using the contact information
31 provided on this letter. Please retain this letter for your records. This survey has passed all ethical
32 clearance procedures at the co-ordinating institution.
33
34

35
36 **Scoring system**
37

38 Today we'll be collectively creating a point scoring system to quantify IPM adoption. The approach
39 will involve weighting questions according to how important we decide they are for IPM. The
40 method will involve allocating a total of 100 points over the questions we've pre-selected to
41 contribute towards the point scoring system. These questions (Q3, 5, 7, 8, 9, 14) are highlighted in the
42 survey.
43
44

45
46 - For Q3, 5, 7, 9, and 14 we will discuss the importance of the options, come to a consensus and rate
47 them on a 1-5 scale.
48
49

50
51 - Q8 is probably the most important question in the set as it captures the IPM control measures
52 actually being undertaken. For this question we will weight the individual control measures for
53 importance using a weighting scale that really allows individual measures to stand out. So, the
54 weighting for Q8 options will involve allocating 100 points over all the measures.
55
56
57

58 Thank you for your contribution to this project.
59
60


Appendix 3: Stakeholder guidance letter for consultation on scoring system for assessing farmers' IPM adoption

Best arable farming practice study

In early 2017, researchers from four institutions surveyed arable farmers in the UK and Ireland. This survey collected information on current Integrated Pest Management (IPM) practices in arable farming. To facilitate our analysis of the responses we would like you to help us identify what best IPM practice is.

At a recent workshop, stakeholders were asked to score each survey question (and options within questions) relating to IPM practice. By creating an IPM scoring system, we can apply these scores to the survey responses and, in this way, rate each respondent in terms of their IPM activity. We need your help in validating the proposed scoring system. Below is a worked example showing what we want you to do.

At the workshop, agronomic and IPM issues relating to key survey questions were discussed in detail to capture the contribution (%) of each to the overall IPM score. Separate scores are applied on a 1 to 5 scale (from 1 being not at all important to 5 being very important) to capture the relative importance of specific management activities within each question. Initial scores for each possible answer were suggested, and were then agreed upon during the discussions. If a clear consensus was not reached, a vote decided the final score. Survey participants' responses to each question will determine what share (%) of the maximum contribution for the question makes to the IPM score.



1
2
3 To validate the proposed scoring system, we are consulting stakeholders to ask them to modify our
4 scores if they wish. We would appreciate it if you could take the time to consider the questions
5 attached, together with the scores given to each at the workshop. Please examine each question and all
6 possible answers and adjust the scores as you see fit. Once you are finished, please send the document
7 back to us.
8
9

10 This is part of a wider study aimed at improving farmer awareness and rates of adoption of IPM. Once
11 we have identified best practice, we will communicate with farmers and advisors through various
12 knowledge transfer activities. Thank you for your contribution to this project. Your responses will
13 remain entirely confidential and individual answers will not be identified in any report we may
14 produce.
15
16

17
18 Dr Henry Creissen Henry.creissen@sruc.ac.uk
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Stakeholder consultation on scoring system for assessing farmers' IPM adoption

NB: In this file the final weights for sub-elements within each question have been inserted in the columns labelled 'Your score', which were of course blank when the file was distributed to the stakeholder panel.

Instructions:

- The original scores from the stakeholder workshop are shown in red. Please insert your score in the blue columns next to the workshop scores. Please indicate the relative contribution of each question to the final IPM score which should total to 100% for all questions combined.
- Please rate each option within each question on a 1 to 5 scale with 1 being not at all important for IPM, to 5 being very important for IPM.

Key questions:

3. Proportion of land in continuous cereal production i.e. cereals on the same land for 5 or more years without a break crop (e.g. OSR, beans, grass)?

Option	Workshop score	Your score
None	5	4.93
1 – 25%	5	4.46
26 – 50%	3	2.98
51 – 75%	2	1.98
77 – 100%	1	1.13

Contribution (%) of this question to the overall IPM score:
Workshop suggestion = 10%. Your suggestion = 11.51%

5. If you typically use an arable rotation, why do you do this?

Option	Strongly disagree		Disagree		Neither agree or disagree		Agree		Strongly agree	
	Workshop score	Your score	Workshop score	Your score	Workshop score	Your score	Workshop score	Your score	Workshop score	Your score
To control weeds	1	0.87	2	1.61	3	2.39	4	3.26	5	4.13
To control disease	1	0.95	2	1.63	3	2.39	4	3.29	5	4.13
To control insect pests	1	0.92	2	1.63	3	2.39	4	3.18	5	4.03
To improve and maintain soil structure and fertility	1	0.97	2	1.63	3	2.45	4	3.29	5	4.15
To spread financial risks	5	3.34	4	2.89	3	2.32	2	1.47	1	1.18
Largely because it's necessary to comply with regulations	5	3.53	4	2.92	3	2.24	2	1.55	1	1.18

Contribution (%) of this question to the overall IPM score:
Workshop suggestion = 10%. Your suggestion = 11.84 %

7. Influences on choice of cereal or oilseed variety grown?

Option	Workshop score	Your score
Recommended lists (where available)	5	4.41
Availability of seed	1	1.67
Advisor recommendation	2	3.09
End-market	1	2.48
Disease resistance	5	4.59
Weed competitiveness	2	2.59
Insect pest resistance	2	2.78
Abiotic stress resistance (e.g. drought/flooding/cold tolerance)	2	2.22
Yield potential	1	2.65
Quality potential	2	2.80
Consistency of performance	3	3.52

Contribution (%) of this question to the overall IPM score:
Workshop suggestion = 5%. Your suggestion = 8.80 %

8. Preventative measures currently employed to control the introduction and spread of pests on their farms?

a) To control weeds

Option	Workshop score	Your score
Stale seedbeds	5	4.57
Full inversion ploughing to control low dormancy weeds	5	4.24
Min-till to control high dormancy weeds	5	4.52
Pre-emergence herbicide applications	3	3.65
Spot spraying weeds (if necessary)	5	4.26
Hand rogueing/hoeing weeds	5	4.39
Manage headlands differently to remainder of field	5	4.33
Fields with high weed levels are harvested last	4	3.57
Weekly crop inspections	5	4.52
Monthly crop inspections	3	2.72
A single crop inspection per season	1	1.09

b) *To control disease*

Option	Workshop score	Your score
Grow resistant varieties	5	4.85
Use certified seed	5	4.20
Test non-certified seed and treat if required	5	4.70
Regularly test soils for soil borne pathogens	2	2.63
Use seed treatments to control disease	5	4.63
Weekly crop inspections	5	4.67
Monthly crop inspections	4	3.24
A single crop inspection per season	1	1.02

c) *To control insects, nematodes and molluscs*

Option	Workshop score	Your score
Encourage beneficial insects through planting habitat	4	3.93
Avoid broad spectrum insecticides	5	4.57
Use seed treatments	5	4.70
Preparations for control of molluscs e.g. avoiding direct drilling or preparation of fine seed bed	5	4.50
Regularly monitor above ground pest populations	5	4.87
Set action thresholds	5	4.61
Regularly test soils for nematodes	3	3.04
Regularly test soils for insect pests	1	1.63
Frequently clean harvesting and storage equipment	5	4.89
Weekly crop inspections	5	4.61
Monthly crop inspections	4	3.15
A single crop inspection per season	1	1.02

Contribution (%) of this question to the overall IPM score:
 Workshop suggestion = 55%. Your suggestion = 47.13%

9. Factors considered when deciding on a pest management plan at the start of the season.

Option	Workshop score	Your score
I don't use pest management plans	0	0.48
Crop walking data from last season, used to assess the performance of various control measures	5	4.52
Technical research on pesticide product efficacy	4	4.35
Weed maps, created and monitored for changes between seasons	4	3.98
Yield maps, used to identify areas requiring specific attention	4	3.96
Cost-benefit analysis of management options	2	2.80
End-market requirements	2	3.02
Variety resistance	4	4.46
Soil borne pathogens	5	4.50
Position in rotation	5	4.83
Anti-resistance strategies	5	4.89
None of the above	-5	-1.11

Contribution (%) of this question to the overall IPM score:
Workshop suggestion = 15%. Your suggestion = 15.30%

14. Member of an agronomy or crop discussion group.

Option	Workshop score	Your score
Yes	3	3.39
No	1	0.87

Contribution (%) of this question to the overall IPM score:
Workshop suggestion = 5%. Your suggestion = 5.85 %

Appendix 4:

For survey data input file see separate Excel file.


Appendix 5: Normal probability-probability plot of the data

Figure shows a Normal probability-probability plot indicating that the distribution of scores over the IPM metric key variables was close to normal.