

Oat bran, but not its isolated bioactive β -glucans or polyphenols, have a bifidogenic effect in an in vitro fermentation model of the gut microbiota

Article

Accepted Version

Kristek, A., Wiese, M., Heuer, P., Kosik, O., Schar, M. Y., Soycan, G., Alsharif, S., Kuhnle, G. G. C. ORCID: <https://orcid.org/0000-0002-8081-8931>, Walton, G. ORCID: <https://orcid.org/0000-0001-5426-5635> and Spencer, J. P. E. ORCID: <https://orcid.org/0000-0003-2931-7274> (2019) Oat bran, but not its isolated bioactive β -glucans or polyphenols, have a bifidogenic effect in an in vitro fermentation model of the gut microbiota. *British Journal of Nutrition*, 121 (5). pp. 549-559. ISSN 0007-1145 doi: 10.1017/S0007114518003501 Available at <https://centaur.reading.ac.uk/82885/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1017/S0007114518003501>

Publisher: Cambridge University Press

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Oat bran, but not its isolated bioactive β -glucans or polyphenols, have a bifidogenic effect in an *in vitro* fermentation model of the gut microbiota¹⁻⁴

Angelika Kristek¹, Maria Wiese², Patrizia Heuer¹, Ondrej Kosik³, Manuel Y Schär¹, Gulten Soycan¹, Sarah Alsharif¹, Gunter G. C. Kuhnle¹, Gemma Walton¹ and Jeremy P E Spencer^{1, 4}

¹Department of Food and Nutritional Sciences, School of Chemistry, Food and Pharmacy, The University of Reading, RG6 6AP, Reading, United Kingdom

²Department of Food Science, University of Copenhagen, Denmark

³Department of Plant Sciences, Rothamsted Research, Harpenden, Herts AL5 2JQ, UK

Keywords: oat, polyphenols, β -glucan, bioactive, gut microbiota, 16S rRNA gene sequencing, short chain fatty acids, prebiotic, bifidogenic

PubMed indexing: Kristek, Wiese, Heuer, Kosik, Schär, Soycan, Alsharif, Kuhnle, Walton and Spencer

⁴Corresponding Author: Jeremy Paul Edward Spencer, Department of Food and Nutritional Sciences, School of Chemistry, Food and Pharmacy, University of Reading, UK, Tel.: +44 118 378 8724; Fax: +44 118 931 0080; E-mail: j.p.e.spencer@reading.ac.uk

Abstract

Wholegrain oats are known to modulate the human gut microbiota and have prebiotic properties (increase the growth of some health promoting bacterial genera within the colon). Research to date mainly attributes these effects to the fibre content; however, oat is also a rich dietary source of polyphenols, which may contribute to the positive modulation of gut microbiota. *In vitro* anaerobic batch-culture experiments were performed over 24 h to evaluate the impact of two different doses (1 and 3 % w/v) of oat bran, matched concentrations of β -glucan extract or polyphenol mix, on the human faecal microbiota composition using 16S RNA gene sequencing and short chain fatty acid analysis (SCFA). Supplementation with oats increased the abundance of Proteobacteria ($p < 0.01$) at 10 h, Bacteroidetes ($p < 0.05$) at 24h and concentrations of acetic and propionic acid increased at 10 and 24 h compared to negative control. Fermentation of the 1 % w/v oat bran resulted in significant increase in SCFA production at 24 h (86 ± 27 mM vs. 28 ± 5 mM; $p < 0.05$) and a bifidogenic effect, increasing the relative abundance of *Bifidobacterium unassigned* at 10h and *Bifidobacterium adolescentis* ($p < 0.05$) at 10 and 24 h compared to negative control. Considering the β -glucan treatment induced an increase in the phylum Bacteroidetes at 24 h, it explains the Bacteroidetes effects of oats as a food matrix. The polyphenol mix induced an increase in Enterobacteriaceae family at 24 h. In conclusion, in this study, we found that oats increased bifidobacteria, acetic acid and propionic acid, and this is mediated by the synergy of all oat compounds within the complex food matrix, rather than its main bioactive β -glucan or polyphenols. Thus oats as a whole food led to the greatest impact on the microbiota.

Keywords: oats, β -glucan, polyphenols, gut microbiota, 16S rRNA gene sequencing, short chain fatty acids, prebiotic, bifidogenic

1. Introduction

A large body of evidence from prospective and intervention studies suggests that a diet rich in oats could significantly reduce the risk of bowel disease [1, 2], cardiovascular disease [3-5] and lowers high blood cholesterol levels [4-7]. Recent reports suggest that oats may act as a prebiotic, modulating the gut microbiota and impacting on metabolic disease risk [8, 9]. To date, it is believed that the protective effect of whole grain oats is mainly due to the presence of dietary fibre, in particular, soluble β -glucan [6, 10-12], and resistant starch [13]. β -glucan is known to lower cholesterol and bile acid absorption through formation of viscous gels in the upper gut [14] and/or directly binding of cholesterol or bile acids. Resistant starch leads to the production of short chain fatty acids (SCFA) in the colon and may also increase the growth of some health promoting bacterial genera within, such as *Bifidobacterium* and *Lactobacillus* [15-18]. Species of these genera influence the cholesterol metabolism through increasing bile-salt hydrolase enzyme activity and the deconjugation of bile acids [19].

Oats, however, are also rich dietary sources of polyphenols, including avenanthramides and phenolic acids [20]. These are likely to contribute to the health effects of a diet rich in oats [21-23] but have not yet been examined in detail. Phenolic acids are found in three different forms within the oat food matrix: as soluble free acids, as soluble conjugates esterified to low molecular weight components such as sugars, and as insoluble bound acids esterified to high molecular weight components including lignin, cell wall polysaccharides and storage proteins [20]. A large proportion of oat polyphenols are bound via an ester bond, and hence are poorly absorbed in the upper intestine and reach the colon [24], where they may beneficially modulate the microbiota [8]. There are no esterases in human tissues that break these ester links [25]; therefore, the main catalytic site is in the colon through cinnamoyl esterase activity of human

67 faecal microbiota, such as *Escherichia coli* (three isolates), *Bifidobacterium lactis* and
 68 *Lactobacillus gasseri* (two strains) [26, 27].

69 Gut microbiota (GM) alterations by whole grain oats have been observed in a number of *in*
 70 *vitro* fermentation models [9, 28-32] and animal studies [13, 33-35]. The fermentation of oats
 71 has led to increased *Bifidobacterium*, *Lactobacillus* [8, 9, 30, 31, 33, 34] and *Bacteroides* [32]
 72 populations and, at the same time decreased *Clostridium* [30, 33].

73 The soluble β -glucans have been one of the most commonly studied components of oats [28,
 74 36, 37]. However, oat-derived isolated β -glucan only impacted on *Bacteroides* growth, not on
 75 *Bifidobacterium* in 24h pH-controlled anaerobic batch culture fermenters [28]. Furthermore,
 76 Crittenden *et al.* [38] reported that β -glucan was fermented by *Bacteroides* spp. but not by
 77 *Lactobacillus* or *Bifidobacterium*. The viscosity and molecular weight of β -glucan may
 78 strongly influence its ability to lead to SCFA production and act as a prebiotic [39].
 79 Additionally, these studies have used the relatively high concentration of 1% v/w of β -glucan
 80 dose [31, 40] which is much higher than the physiological β -glucan concentration delivered
 81 from eating oats. To date, none of the oat or β -glucan *in-vitro* fermentation studies assessed the
 82 polyphenol content of their treatment [9, 28, 32]. Thus, little information exists on the ability
 83 of oat-derived avenanthramides, hydroxycinnamic and hydroxybenzoic acids to influence the
 84 GM.

85 The main purpose of the present *in vitro* study was to examine if a physiologically relevant
 86 dose of polyphenol mix (avenanthramide, hydroxycinnamic acids and benzoic acid
 87 derivatives), β -glucan extract or digested oat bran as whole food matrix beneficially modulates
 88 the GM. Secondly, the study aimed to assess the impact of different doses of digested oat bran
 89 on the GM, notably using an oat bran that was well characterised in terms of physicochemical
 90 properties.

91

92

93 2. Materials and Methods

94 2.1. Reagents

95 All chemicals and reagents were purchased from Sigma-Aldrich Chemical Co Ltd. (Pool,
96 Dorset, UK) or Fisher (Loughborough, Leics, UK) unless stated otherwise. Mixed-linkage β -
97 glucan kit was supplied by Megazyme Co (Wicklow, Ireland). The anaerobic jar (AnaerojarTM
98 2,5L) and gas-generating kit (AnaeroGen TM) were obtained from Oxoid Ltd (UK), the
99 dialysis tube from Spectrum (VWR International). Media and instruments were autoclaved at
100 121°C for 15 min. HPLC column and guard cartridges were obtained from Phenomenex
101 (Cheshire, UK). PowerSoil[®]DNA Isolation Kit was purchased from Mo Bio Laboratories, Inc
102 (USA), the primers for the 16S rRNA gene amplification from Integrated DNA Technologies,
103 BVBA (Belgium), AccuPrimeTM SuperMix II from Life Technologies (CA, USA), AMPure
104 XP beads from Beckman Coulter Genomic (CA, USA).

105 2.2. Oats and Controls

106 The oat bran was purchased from White's (Tandragee, Northern Ireland). The oat
107 macronutrient composition was analysed by Campden BRI laboratories (Total carbohydrate,
108 Available carbohydrate, Total dietary fibre- AOAC 991.43 method, Total fat – BS 4401, and
109 protein – AOAC 981.10 method), whereas the detailed polyphenol content was measured in
110 our laboratory at the University of Reading based on a previous method [41]. The beta-glucan
111 method used was employed specifically to quantify 1,3:1,4- β -D-glucan. Synergy1 the
112 oligofructose-enriched inulin was supplied by Beneo (Belgium) and 94%- β -glucan extract
113 from Megazyme Co (Wicklow, Ireland). Polyphenols were purchased from Sigma-Aldrich
114 Chemical Co Ltd. (Pool, Dorset, UK).

115 2.3. *In vitro* digestion of oat bran (from mouth to small intestine)

116 The method employed was adapted from Mills *et al.* [42]. Oat bran was digested *in vitro* in
117 three phases: the oral phase, the gastric phase and the small intestinal phase. 60 g of oat bran
118 was mixed with 150 ml of sterile and distilled water and homogenised, transferred into a 500
119 ml Duran bottle, and microwaved for 1 minute. In the oral phase, 20 mg of α -amylase was
120 dissolved in 6.25 ml CaCl_2 (1 mM, pH 7.0) and added to the solution, then incubated at 37°C
121 for 30 minutes on a shaker set at 120 *xg*. After incubation, the pH was adjusted to 2.0 with 6
122 M HCl solution and the gastric phase introduced by adding 2.7 g pepsin in 25 ml HCl (0.1 M)
123 and further incubated for 2 hours under the same conditions. In the small intestinal phase, 560
124 mg pancreatin and 3.5 g bile were mixed with 125 ml NaHCO_3 (0.5 M) and dispensed into the
125 mix. The pH was adjusted to 7.0 then incubated for 3 hours at 37 °C with shaking. Finally, the
126 sample solution was transferred to a seamless semi-permeable 100-500 Dalton molecular
127 weight cut-off regenerated cellulose dialysis tubing and dialysed against NaCl (0.01 M at 5°C)
128 to remove low molecular mass digestion products. After 15 hours, the dialysis fluid was
129 changed and the process continued for an additional 2 hours. The digested oat bran mix was
130 collected and transferred into several 250 ml clear plastic containers, frozen to -80° and freeze-
131 dried for 5 days to remove all fluid content.

132 2.4. Extraction and analyses of polyphenols from undigested and digested oat bran

133 Polyphenols were extracted from undigested (raw) and digested (after *in vitro* digestion) oat
134 bran in two separate fractions (i.e. free and conjugated or bound) using the method of Schar *et*
135 *al.* [41]. The phenolic acids and avenanthramides in oat extracts were identified and quantified
136 using a high-performance liquid chromatography (HPLC) Agilent 1100 series (Agilent
137 Technologies Ltd) equipped with a quaternary pump, autosampler, column thermostat, sample
138 thermostat and photo diode array detector. Compound separation was achieved by a Kinetex

139 biphenyl column (100 Å 250 x 4.6 mm length, 5 µm particle size; Phenomenex) and using a
140 gradient elution. Mobile phase A consisted of 0.1 % (v/v) formic acid in HPLC water (A),
141 while mobile phase B was 0.1 % (v/v) formic acid in methanol. The following optimised
142 gradient protocol was run: 0 min, 95% A, 5 % B; 20 min, 75 % A, 25 % B; 25 min, 74 % A,
143 26 % B; 30 min, 65 % A, 35 % B; 40 min, 64 % A, 36 % B; 53 min, 30 % A, 70 % B; 56 min,
144 5 % A, 95 % B; 61 min, 5 % A, 95 % B; 62 min, 95 % A, 5 % B; 65 min, 95 % A, 5 % B. The
145 flow rate of the mobile phase was 1.0 ml/min and the sample injection volume was 20 µl [43].
146 The absorbance was recorded at 254, 280 and 320 nm and quantification was based on 12 point
147 linear calibration curves (mean $R^2 > 0.99$) and as a ratio to the internal standard (i.e. 3,5-
148 dichloro-4-hydroxybenzoic acid) to account for losses during extraction ($R^2 \geq 0.99$).

149

150 2.4. Determination of the β -glucan content

151 The β -glucan content was analysed with the 1,3:1,4 mixed-linkage β -glucan kit. The assay uses
152 lichenase and β -glucosidase to metabolize β -glucan to β -gluco-oligosaccharides and
153 subsequently glucose. Glucose then reacts with GOPOD reagent and its absorbance was
154 measured at 510 nm by UltroSpec 1100 photo spectrometer (Scintech Instruments LLC, USA).
155 The final β -glucan content was calculated by the Megazyme Mega-Calc™ tool [44].

156 The molecular weight of the β -glucan was determined by size-exclusion HPLC. The
157 chromatography system consisted of three serially connected columns (Shodex SB-G, Shodex
158 SB-806M, Shodex SB-804 HQ) and an UV-MALLS-Viscometer-dRI detector. The column
159 temperature was set at 40°C, the mobile phase was MiliQ water (Millipore, Bedford, MA)
160 containing 0.02 % sodium azide, and the flow rate set at 0.5 ml/min. Samples were prepared at
161 a concentration of 10 mg/ml, heated at 60°C for 3 hours under constant shaking, syringe filtered
162 (0.45 µm PVD; Whatman, NY) and diluted to a concentration of 1 mg/ml before injection.

163 2.5. pH controlled faecal batch culture fermentation

164 Substrate ability to modulate gut microbiota was determined using anaerobic, stirred, pH and
 165 temperature controlled faecal batch cultures. Batch culture fermentation vessels (300 ml
 166 volume: one vessel per treatment) were sterilised and filled with 135 ml of sterile basal medium
 167 (g/l: 2 g peptone, 2 g yeast extract, 0.1 g NaCl, 0.04 g K₂HPO₄, 0.04 g KH₂PO₄, 0.01 g
 168 MgSO₄·7H₂O, 0.01 g CaCl₂·6H₂O, 2 g NaHCO₃, 2 ml Tween 80, 0.05 g Hemin (dissolved in
 169 a few drops of 1 M NaOH), 10 µl vitamin K, 0.5 g L-cysteine HCl, 0.5 g bile salts and 4ml of
 170 resazurin solution (0.02 %)). Before addition to the vessel, the medium was adjusted to pH 7.0
 171 and autoclaved. The sterile medium was gassed overnight with O₂-free N₂ (15 ml / min) to
 172 establish anaerobic conditions. To mimic the proximal region of the human large intestine, pH
 173 was held in the range of 6.7 - 6.9 by automatic addition of 0.5 M NaOH or 0.5 M HCl and
 174 controlled via pH meter controllers (Electrolab, UK) and the temperature was kept at 37 °C.
 175 Faecal samples were collected from three healthy female donors, who were between 25 and 40
 176 years old, with no history of bowel disorders, who had not received antibiotic treatment for at
 177 least 6 months before the study and had not consumed pre- or probiotic supplements one month
 178 before the study. All donors were informed of the study aims and procedures, and provided
 179 their verbal consent for stool samples to be used for the experiments in compliance with the
 180 ethics procedures required at the University of Reading. Samples were collected in anaerobic
 181 jars and used within one hour of collection. Samples were diluted 1:10 (w/v) with anaerobic
 182 phosphate buffer (0.1 M, pH 7.4) and homogenised in a stomacher for 2 mins; the batch culture
 183 systems were inoculated with 15 ml faecal slurry from an individual sample. The vessels were
 184 treated with the following substrate: 1 % w/v, 1.5 g digested oat bran (OAT1.5); 3 % w/v, 4.5
 185 g digested oat bran (digestibility of oats see McCane et al.[45]) (OAT4.5); 0.12 % w/v, 180 mg
 186 94 % β-glucan extract (BG); 0.01 % w/v, 1.7 mg polyphenol mix (same profile as 4.5 g digested
 187 oats) (POLY); 1 % w/v, 1.5 g Synergy1 (positive control, PC); an additional vessel was

prepared under the same conditions but without any substrate, negative control (NC) (Table 2). The Synergy 1 is an inulin enriched with oligofructose, has been proved their prebiotic effects [46]. The 3 % w/v oat was chosen as the highest dose since it would be the concentration reached by an average person consuming 60 g oat bran (i. e. assuming 30 g oat bran reaches the colon and colonic total volume is 1L) [45]. Based on this calculation, β -glucan extract and polyphenol mix treatments were matched to the dose present in 3 % w/v oat bran. Treatment with OAT1.5 was added to the experiment to monitor the impact of the oat dose on the prebiotic effect. Five millilitre samples were collected at 0, 5, 10, 24h, of which 1 ml aliquots were centrifuged at 13,000 $\times g$ for 10 min. Supernatants and pellets were stored separately at -20°C until analysis.

2.6. High throughput sequencing of the gut microbiota

The GM compositions and dynamics were determined using tag-encoded 16S rRNA gene MiSeq-based (Illumina, CA, USA) high throughput sequencing. DNA was extracted from 2 ml of faecal samples from fermentation vessel using power soil DNA isolation kit. The bead-beating was performed in 3 cycles of 15 second each at 6.5 pulse / s. (The FastPrep-24™ 5G Instrument, MP Biomedicals). DNA concentrations and purity were determined using Nanodrop 1000 (ThermoScientific, USA). The V3 region of the 16S rRNA gene was amplified using primers compatible with the Nextera Index Kit NXt_338_F: 5'-TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGACWCCTACGGGWGGCAGCAG-3' and NXt_518_R: 5'-GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGATTACCGCGGCTGCTGG-3' [47]. PCR reactions and library preparation were conducted as described in [48]. Briefly, PCR reactions containing 12 μ l AccuPrime SuperMix II, 0.5 μ l of each primer (10 μ M), 5 μ l of

212 genomic DNA (~10 ng / μ l), and nuclease-free water to a total volume of 20 μ l were run on a
213 SureCycler 8800 (Agilent, CA, USA). Applied cycling conditions were: denaturation at 95 °C
214 for 2 min; 35 cycles of 95 °C for 15 s, 55 °C for 15 s and 68 °C for 40 s; followed by final
215 elongation at 68°C for 5 min. To incorporate primers with adapters and indices, PCR reactions
216 contained 12 μ l Phusion High-Fidelity PCR Master Mix, 2 μ l P5 and P7 primer, 2 μ l PCR
217 product and nuclease-free water for a total volume of 25 μ l.

218 Cycling conditions applied were: 98°C for 1 min; 12 cycles of 98°C for 10 s, 55°C for 20 s and
219 72°C for 20 s; elongation at 72°C for 5 min. The amplified fragments with adapters and tags
220 were purified using AMPure XP beads. Prior to library pooling, clean constructs were
221 quantified using a Qubit fluorometer (Invitrogen, Carlsbad, CA, USA) and mixed in
222 approximately equal concentrations to ensure even representation of reads per sample. 180 bp
223 pair-ended MiSeq (Illumina) sequencing was then performed according to the instructions of
224 the manufacturer. The raw dataset containing pair-ended reads with corresponding quality
225 scores was trimmed using CLC Genomic Workbench (CLC bio, Aarhus, Denmark). Trimming
226 settings were set to a low-quality limit of 0.01, with no ambiguous nucleotides allowed, and
227 trimming off the primer sequences. Merging overlapped reads was performed using the "Merge
228 overlapping pairs" tool using default settings. The Quantitative Insight Into Microbial Ecology
229 (QIIME) tool (version. 1.8.0; Open source software) was used for further analysis [49].

230 Purging the dataset from chimeric reads was performed using USEARCH, while the usearch61
231 method was used for Operational Taxonomic Units (OTUs) selection [50]. The Greengenes
232 (version 12.10) 16S rRNA gene database and EzTaxon were used as reference [51, 52].

233 High throughput sequencing yielded 12465219 sequences free from chimeric reads, providing
234 an average \pm SD of 178202 ± 80036 sequences per sample (Range 292-447040), five baseline
235 samples and one 10h sample were excluded due to low reads (292-509).

236 2.7. Short-chain fatty acid analysis

237 The defrosted supernatant samples were filtered (0.22 μ m, Millipore) and spiked with 50 μ l
 238 internal standard (2-ethyl butyric acid, 100 mM). SCFAs were measured in an ion exclusion
 239 HPLC system (Agilent 1100 Series) equipped with DAD detector (G-1315B), autosampler
 240 (G1316A) and Aminex HPX-8711 column (300 x 7.8mm) heated to 84°C. Samples (20 μ l)
 241 were injected in duplicate, and UV absorption was measured at 214 nm. The mobile phase was
 242 2.5 mM aqueous sulphuric acid run at a flow rate of 0.6 ml/min. Peaks were integrated using
 243 Agilent ChemStation software (Agilent Technologies, Oxford, UK) and single point internal
 244 standard method. Peak identity and quantification were determined using a mixture of
 245 standards of acetic, butyric, iso-butyric, propionic, lactic, formic, valeric and isovaleric acids.
 246 Quantification was based on the 10-point calibration curve of analytical standards ($R^2 \geq 0.99$).

247 2.8. Statistics

248 For calculation of alpha and beta diversity measurement of the sequencing data, the d- and e-
 249 values were set to 9800 reads per sample (85 % of the sequence number of the most indigent
 250 sample). Alpha diversity measures expressed with an observed species with Chao1, the
 251 sequence similarity 97 % OTUs value were computed for rarefied OTU tables using the alpha
 252 rarefaction workflow. Differences in alpha diversity were determined using a t-test-based
 253 approach employing the non-parametric (Monte Carlo) method (999 permutations)
 254 implemented in the compare alpha diversity workflow with QIIME.

255 The relative distribution of the genera registered was calculated and summarised at the genus
 256 level OTU tables, followed by Principal Coordinates Analysis (PCoA) plots generated with the
 257 Jackknifed beta diversity workflow based on 10 distance matrices calculated using 10
 258 subsampled OTU tables with QIIME. The *p*-value and the conservative FDR-corrected *p*-value

for multiple comparisons are reported. 3D plots were constructed from the three primary PCs from the PCoA of the MiSeq analysis to visualise group differences in the composition of the GM.

For the rest of the data analysis, GraphPad Prism statistics software package version 7 was used. One-way ANOVA was used to determine differences between fermentation treatments (OAT1.5, OAT4.5, POLY, BG) at the same time point (0, 5, 10 or 24 h), followed by the least significant difference (Bonferroni) post hoc test. A repeated measures ANOVA was used to explore the differences within the same treatment (OAT1.5, OAT4.5, POLY, BG) with all the time points (0, 5, 10 and 24 h) with Bonferroni as the post hoc test. In addition to these analyses, the p values were corrected using false discovery rate (FDR), $p \leq 0.05$ was considered statistically significant.

3. Results

3.1. The composition of undigested and digested oat bran

The macronutrient, fibre and phenolic content of undigested and digested oat bran are reported in Table 1. The *in vitro* oral, gastric and small intestine digestion reduced the carbohydrate content by 53 %, available carbohydrate by 62 %, while the dietary fibre and β -glucan content remained stable, losing only 2 % and 12 %, respectively. Free and conjugated, bound polyphenol compounds decreased by 48 % and 26 %, respectively.

3.2. Polyphenols and β -glucan content in oat bran and extract

The OAT4.5 contained 1.73 mg polyphenols (total amount of free + conjugated and bound), the POLY intervention were matched to OAT4.5 (Table 2). OAT1.5 and OAT4.5 delivered 56.67 mg and 170 mg β -glucan, respectively; the BG vessel to match OAT4.5 180 mg of 94% β -glucan extract was added (Table 2). The molecular weight of the β -glucan in the digested oat

bran and the extract were similar (mean \pm SD 335.5 \pm 1.0 % kDa and 387.3 \pm 1.0 % kDa; $p > 0.05$).

3.2. Changes in alpha and beta diversity

Independent of the donor, alpha diversity decreased with oats treatments over the 24h fermentation period (Chao1 2618 \pm 15 at 0h vs. 1577 \pm 22 at 24h for OAT1.5) but was constant over time in the negative control (Chao1 2639 \pm 87 at 0h vs. 2488 \pm 77 at 24h) and POLY (Chao1 2655 \pm 138 at 0h vs. 2241 \pm 320 at 24h).

PCoA analyses based on unweighted UniFrac distance matrices of all samples showed a clear donor effect (Figure 1.A). However, the donor effect was no longer significant when using weighted UniFrac distance matrix analysis (ANOSIM test, Donor 1. Vs. Donor 2 $p = 1$, Donor 1 vs Donor 3 $p = 1$, Donor 2. Vs. Donor 3 $p = 1$, Figure 1.B).

Across all samples, seven bacterial phyla were classified and one phylum designated as unassigned. Phyla composition was similar for all donors ($p < 0.06$), although at baseline the third donor had lower Tenericutes ($p < 0.01$), while the second donor had higher Actinobacteria ($p < 0.05$). At baseline, the bacterial communities, were dominated by Firmicutes (57-67 %) and Bacteroidetes (32-41 %), while the remaining five phyla including Actinobacteria (0.8-1.4 %), Verrucomicrobia, Cyanobacteria, Tenericutes and Proteobacteria (0.4-0.9 %) constituted < 1.5 % of the community (Figure 2.).

3.3. Compositional Shifts

Supplementation with OAT4.5 led to a lower abundance of Firmicutes ($p < 0.02$) at 5 to 24 h and to an increased abundance of Proteobacteria ($p < 0.01$) and Bacteroidetes at 24 h compared to NC. A similar trend was observed for OAT1.5 (p values)(Figure 2).

304 The OAT1.5 treatment had a significant effect on the relative abundance of Actinobacteria at
 305 10 h. The relative abundance of Actinobacteria differed between OAT1.5 and OAT4.5 at
 306 baseline (1.1 % and 0.8 %, respectively). Following 10 h fermentation, Actinobacteria were
 307 almost four times higher in OAT1.5 compared to OAT4.5 (4.9 % and 1.3 %, respectively),
 308 however, differences did not persist over 24 h (4.2 % vs. 3.4 %, respectively). The treatment
 309 of BG and POLY did not promote the growth of Actinobacteria, even though their dose was
 310 matched to the OAT4.5 treatment. The PC, Synergy1 resulted in the strongest bifidogenic
 311 effect and the relative abundance of Actinobacteria (baseline 1.1 % vs. 5 h at 31.1 %, 10 h at
 312 11.5 %, $p < 0.05$) (Figure 2.).

313 Focusing on changes over 24 h, the abundance of Bacterioidetes increased with BG (49.3 % vs.
 314 NC for 21 %), whereas Proteobacteria significantly decreased in comparison to the NC (14 %
 315 vs. 33.5 %). POLY treatment, the abundance of Proteobacteria increased over the course of
 316 fermentation (45 % vs. NC for 33.5 %) and decreased for Bacterioidetes ($p < 0.04$) (18.6 % vs.
 317 46.8 % for OAT4.5). The oats and BG treatments promoted the growth of Bacterioidetes (Figure
 318 2.).

319 At the lower taxonomic level and across all samples, 59 genera and 69 species were identified
 320 when setting the cut-off at a relative abundance above 0.1 % of the community (Table 3). 5 h,
 321 the OAT4.5 treatment led to a significantly lower abundance of *Ruminococcus* (0.1 %), and
 322 *Coprococcus* (0.1 %) genus and, at the same time, increased the proportion of unassigned
 323 *Enterobacteriaceae* (61.2 %) compared to NC (25.8 %) (Table 3). Most notably, OAT1.5
 324 increased the abundance of *Bifidobacterium unassigned* (1.5 % vs. 0.2 for NC, $p < 0.05$) at 10
 325 h and *Bifidobacterium adolescentis* at 10 h (1.9 % vs. 0.3 % for NC, $p < 0.04$) and at 24 h (2
 326 % vs. 0.4 for NC, $p < 0.02$) (Table 3), whereas OAT4.5 did not significantly impact the relative
 327 abundance of these species due to the high standard deviation among donors. Using EzTaxon

database, we were able to further identify the species designation of several OTUs at 24 h with BG and POLY treatments. At 24 h, the BG treatment had higher abundances in *Bacteroides ovatus* (13.5 %), *B. uniform* (12.9 %) and *B. xylaniosolvans* (3.7 %) species, while POLY led to the most abundant levels of *Enterobacteriaceae* species, *Hafnia alvei* (21.7 %). However, these changes mediated by BG and POLY treatments were not significantly different compared to NC.

3.4. Production of SCFA

OAT4.5 led to a high production of total SCFA compared to NC at 10 h (98.2 ± 22 mM vs 12.8 ± 4 mM respectively) and 24 h (151.5 ± 43 mM vs 28.1 ± 5 mM). The concentration of butyric acid was significantly increased after 24 h fermentation of OAT1.5 ($p < 0.05$), and PC at 10 h ($p < 0.05$) and 24 h ($p < 0.01$) compared to NC (Figure 3). Acetic acid was the most abundant SCFA. Both oat treatments resulted in a significant increase of acetic acid at 10 h ($p < 0.01$), and 24 h ($p < 0.01$) compared with baseline and NC (Figure 3.). At 24 h, OAT4.5, OAT1.5 and PC significantly increased the concentration of propionic acid compared to NC (48 ± 24 mM, 16.7 ± 3 mM, 21 ± 9 mM and 4.5 ± 0.3 mM, respectively $p < 0.01$, 0.05 and 0.05 , respectively). Lactic and formic acids increased upon fermentation of OAT4.5 after 10 h compared to baseline, but did not reach significance due to the large variation among donors. BG led to a small production of acetic and butyric acids (not significant) whereas POLY did not induce SCFA production. Overall, there was a dose-response effect between the two oats doses in total SCFA at 10 h (98.2 ± 22 mM vs 58 ± 19 mM respectively, $p < 0.05$) and 24 h (151.5 ± 43 mM vs 86.1 ± 27 mM, $p < 0.05$).

4. Discussion

This study aimed to identify the impact of different doses of oats and its isolated bioactive compounds (i.e. β -glucan or polyphenols) on the faecal gut microbiota using *in vitro* systems. Digested oat bran decreased alpha diversity and had a bifidogenic effect but isolated β -glucan or polyphenol mix given at a matched dose did not induce a similar effect. (Figure 2.). This is likely to be because these compounds alone did not provide enough energy for bacterial growth. We have demonstrated that OAT1.5 significantly increased the proliferation of *Bifidobacterium adolescentis* (Table 3). An increase in *Bifidobacterium* (genus level) was reported by Connolly *et al.* [9] and Kedia *et al.* [30] after fermentation of 1 % and 5 % oat grains in similar *in vitro* studies. *B. adolescentis* has a number of benefits including prevention of the development of diabetes by stimulating insulin secretion. Furthermore *B. adolescentis* has the ability to synthesise and secrete the neuroactive substance gamma-aminobutyric acid (GABA). GABA facilitates communication between bacteria and the human nervous system, enabling release of other neurotransmitters from specific epithelial intestinal cells [53, 54]. Our main findings suggest that oat bran could have beneficial effects on the host through increasing the relative abundance of *B. adolescentis*.

In accordance with the literature [55, 56], the compositional analysis of digested oat bran showed limited degradation of β -glucan in the stomach and the small intestine (Table 1). BG (180 mg – 387 kDa) did not introduce changes to Actinobacteria phyla (Figure 2.), which is supported by previous studies on β -glucan [28, 38]. However, BG tended to promote the growth of beneficial *Bacteroides uniformis*, *Bacteroides ovatus* and *Bacteroides xylaniosolvens* compared with NC (Table 3), which is in line with reports of Wang *et al.* [57]. Additionally, Hughes *et al.* [28] used a similar anaerobic batch culture system for fermentation of 0.5 g β -glucan and also showed an increase in a member of Bacteroidetes, specifically in the

375 *Bacteroides-Prevotella* group following 24 h fermentation. However, the dose of β -glucan used
 376 by Hughes *et al.*, was almost three times higher in concentration compared with what was
 377 found in 60 g oat bran (containing 0.17 g β -glucan), which would equate to consumption of
 378 150 g oat bran.

379 While human digestive enzymes cannot degrade plant cell wall polysaccharides, gut
 380 xylanolytic bacteria (e.g. *B. xylaniosolvens*, *B. uniformis* and *B. ovatus*) can, while producing
 381 SCFA with potential health-beneficial effects [58-60]. Certain strains of *B. uniformis*
 382 downregulate gene and protein expression of pro-inflammatory cytokines, notably iNOS and
 383 PPAR- γ , IFN- γ , resulting in reduced inflammatory status [61], suggesting that oat β -glucan
 384 could have beneficial effects on human health. More powered studies should be applied to
 385 confirm this effect.

386 *In vitro* upper gut digestion (mouth to small intestine) reduced polyphenols in the free -
 387 conjugated, bound fractions by 48 % and 26 %, respectively (Table 1). This is in agreement
 388 with findings by Dall'Asta *et al.* [62] on phenolic bioaccessibility in durum wheat aleurone
 389 fractions where caffeic and sinapic acids appeared as the most bioaccessible (83.3 % and 79.5
 390 %) while total ferulic acid and *p*-coumaric acid were less bioaccessible (29.5 % and 40.7%)
 391 with *in vitro* digestion. Moreover, several phenolic metabolites, including vanillic acid, 4 - and
 392 3 - hydroxyhippuric acids, sulfate-conjugates of benzoic and ferulic acids are derived from the
 393 hepatic and microbial metabolism of oat brans [41]. Schar *et al.* [41] found relevant
 394 concentrations in urinary excretion of 30 different phenolics, and amounted to a total excretion
 395 of 33.7 ± 7.3 μ mol, suggesting that a high proportion of oat phenolics are bioavailable with
 396 absorption occurring both in the small intestine and then in the large intestine within eight hours
 397 of consumption.

In the current study, POLY did not change the abundance of the Actinobacteria phyla, which is similar to the research conducted by Gwiazdowska *et al.* [63], which showed that 20 µg/ml polyphenols had an effect on *Bifidobacterium* after 1 h incubation, but no effect at 24 h. The effects of polyphenols on microbial composition, may also be related to the fact that there is no carbohydrate energy available for this fermentation. Therefore, at least part of the change may also be due to utilisation of protein as an energy source. However, in the current study a POLY- induced increase in Enterobacteriaceae phyla and *Hafnia alvei* species was detected [64]. Several studies connected this group with polyphenol-degrading metabolism [65, 66]. While Wen Gu *et al.* [67] and Hunter *et al.* [68] observed that Enterobacter strains could transform ferulic acid to vanillin via the non-oxidative decarboxylation, Kuntz *et al.* [69] believed that the *H. alvei* significantly attenuated the expression of adhesion molecules and cytokine secretion (IL-8 and IL-6), resulting in reduced inflammation [69]. The higher relative abundance of Enterobacteriaceae phyla and *H. alvei* species in POLY and digested oat bran treatments might be associated with the metabolic activity of these microbial groups towards plant-derived polyphenols and saccharides [70].

Previous oat fermentation studies have not shown significant increases in the growth of Enterobacteriaceae phyla and *H. alvei* species [30], most likely because the fluorescence *in situ* hybridisation (FISH) method used did not target Enterobacteriaceae or allow analysis at the species level. We have detected an increased presence of Proteobacteria phyla (Figure 2) in all 3 vessels including NC, which might be due to the artificial conditions of batch culture systems [71, 72]. Facultative anaerobic microbes are indeed less abundant in the human colon [73] but this batch culture model may not be able to achieve a strict anaerobic environment, causing an increase in Proteobacteria phyla [72].

Despite *in vitro* enzymatic digestion (mimicking the digestion in the upper intestine prior to colon fermentation), the available carbohydrate content of the oat bran remained high (62 %) (Table 1). Wholegrain oats are a source of starch (60 % of the total dry matter of the oat grain), consisting of 7 % rapidly digestible starch, 22 % slowly digestible and 25 % resistant starch [74] all of which contribute to the available carbohydrates. The findings of Englyst *et al.* suggest that human digestive enzymes do not break down dietary resistant starch [75]. Strikingly, 30 % resistant starch type 4 also led to species level alterations in a colon and increases in *B. adolescentis*. [76]. A previous study [40] reported that 1 % (w/v) arabinoxylan, a dietary fibre found in wholegrain including oats, has a bifidogenic effect using a similar 24 h batch culture fermentation *in vitro* model, yet oats generally have about 3.8 – 13.2 % of arabinoxylan [20]. This, in combination with our findings, suggests that interactions among a wide range of dietary polysaccharides may explain the bifidogenic effect of oats [77].

The current results indicate that digested oat bran treatments increase SCFA production dose-dependently, with a higher dose inducing higher acetate and propionate production but lower butyrate (Figure 3.). Lactic acid disappeared at 24 h, possibly because it can be further metabolised by bacteria to acetic and propionic acids [78]. Acetate and lactate are widely produced by different bacteria; however, bifidobacteria have a distinct pathway, transforming glucose to acetate and lactate [79]. SCFA can create acidic conditions in the human colon inhibiting the growth of pathogens and reducing harmful enzyme activities in the human intestine, while they also act as an energy source for gut epithelial cells [80, 81]. Furthermore, propionate derived from carbohydrate fermentation is a substrate for gluconeogenesis in humans, and also inhibits the utilisation of acetate for cholesterol synthesis in the colon and liver [82, 83]. Therefore oat bran may have a beneficial impact on human health through SCFA production.

In previous studies [84, 85] differences were observed between the activity of pure and of food matrix derived β -glucan on small intestine digestion and lipolysis. Purified β -glucan was observed to have less gel forming capacity and a lesser effect on lipolysis and cholesterol metabolism, than the matrix derived β -glucan. In addition, dietary fibre induced SCFA production has been shown to interact with phenolics [86], where butyric acid increased the ferulic acid absorption in the colon, potentially translating to higher concentrations of ferulic acid in systemic circulation *in vivo*. These findings indicate positive interactions of fibres and polyphenols within the food matrix on cholesterol metabolism and the bioavailability of phenolics, and also highlights the importance of considering the structure and physicochemical properties of foods, and not just the nutrient content.

In conclusion, our study has shown that oat bran as a complex food matrix beneficially increases the number of *B. adolescentis* and the amount of SCFA production. In contrast, a matched dose of the isolated bioactive compounds, β -glucan and other polyphenols did not show any effect on the abundance of Actinobacteria. However, by regulating the Bacteroides and Enterobacteriaceae families, they may also contribute to further health benefits.

ACKNOWLEDGMENTS

We kindly thank Prof Dennis Sandris Nielsen Department of Food Science, University of Copenhagen, Denmark for his support to conduct DNA extraction and amplification in his laboratory.

CONTRIBUTIONS

The authors' contributions were as follows: AK, MYS and GW designed the study; OK analysed the β -glucan molecular weight, AK and PH conducted the *in vitro* fermentation experiments; MW and AK conducted the 16S rRNA gene sequencing, AK and MW conducted the statistical analyses. AK, MW, MYS, GW, GK, GS, SA and JS interpreted the data and AK,

469 MW, MYS, GM and JS drafted the manuscript. AK had primary responsibility for the final
 470 content. All authors agreed on the final version of the manuscript.

471 The authors declare no conflict of interest

472 References

- 473 1. Haboubi, N.Y., S. Taylor, and S. Jones, *Coeliac disease and oats: a systematic review*.
 474 Postgrad Med J, 2006. 82(972): p. 672-8.
- 475 2. Thies, F., et al., *Oats and bowel disease: a systematic literature review*. Br J Nutr, 2014.
 476 112 Suppl 2: p. S31-43.
- 477 3. Ye, E.Q., et al., *Greater whole-grain intake is associated with lower risk of type 2*
 478 *diabetes, cardiovascular disease, and weight gain*. J Nutr, 2012. 142(7): p. 1304-13.
- 479 4. Thies, F., et al., *Oats and CVD risk markers: a systematic literature review*. Br J Nutr,
 480 2014. 112 Suppl 2: p. S19-30.
- 481 5. Ruxton, C.H.S. and E. Derbyshire, *A systematic review of the association between*
 482 *cardiovascular risk factors and regular consumption of oats*. British Food Journal,
 483 2008. 110(11): p. 1119-1132.
- 484 6. Whitehead, A., et al., *Cholesterol-lowering effects of oat beta-glucan: a meta-analysis*
 485 *of randomized controlled trials*. Am J Clin Nutr, 2014. 100(6): p. 1413-21.
- 486 7. Bao, L., et al., *Effect of oat intake on glycaemic control and insulin sensitivity: a meta-*
 487 *analysis of randomised controlled trials*. Br J Nutr, 2014. 112(3): p. 457-66.
- 488 8. Connolly, M.L., et al., *Hypocholesterolemic and Prebiotic Effects of a Whole-Grain*
 489 *Oat-Based Granola Breakfast Cereal in a Cardio-Metabolic "At Risk" Population*.
 490 Frontiers in Microbiology, 2016. 7: p. 1675.
- 491 9. Connolly, M.L., J.A. Lovegrove, and K.M. Tuohy, *In vitro fermentation characteristics*
 492 *of whole grain wheat flakes and the effect of toasting on prebiotic potential*. J Med
 493 Food, 2012. 15(1): p. 33-43.
- 494 10. Tosh, S.M., *Review of human studies investigating the post-prandial blood-glucose*
 495 *lowering ability of oat and barley food products*. Eur J Clin Nutr, 2013. 67(4): p. 310-
 496 7.
- 497 11. Tosh, S.M. and Y. Chu, *Systematic review of the effect of processing of whole-grain*
 498 *oat cereals on glycaemic response*. Br J Nutr, 2015. 114(8): p. 1256-62.
- 499 12. Tappy, L., E. Gugolz, and P. Wursch, *Effects of breakfast cereals containing various*
 500 *amounts of beta-glucan fibers on plasma glucose and insulin responses in NIDDM*
 501 *subjects*. Diabetes Care, 1996. 19(8): p. 831-4.
- 502 13. Drzikova, B., G. Dongowski, and E. Gebhardt, *Dietary fibre-rich oat-based products*
 503 *affect serum lipids, microbiota, formation of short-chain fatty acids and steroids in rats*.
 504 Br J Nutr, 2005. 94(6): p. 1012-25.

- 505 14. Zhang, Y., et al., *The effect of oat beta-glucan on in vitro glucose diffusion and glucose*
506 *transport in rat small intestine*. J Sci Food Agric, 2016. 96(2): p. 484-91.
- 507 15. Ovando-Martínez, M., et al., *Effect of hydrothermal treatment on physicochemical and*
508 *digestibility properties of oat starch*. Food Research International, 2013. 52(1): p. 17-
509 25.
- 510 16. Rasane, P., et al., *Nutritional advantages of oats and opportunities for its processing as*
511 *value added foods - a review*. J Food Sci Technol, 2015. 52(2): p. 662-75.
- 512 17. Gangopadhyay, N., et al., *A Review of Extraction and Analysis of Bioactives in Oat and*
513 *Barley and Scope for Use of Novel Food Processing Technologies*. Molecules, 2015.
514 20(6): p. 10884-909.
- 515 18. Gibson, G.R., et al., *Dietary modulation of the human colonic microbiota: updating the*
516 *concept of prebiotics*. Nutr Res Rev, 2004. 17(2): p. 259-75.
- 517 19. Oner, O., B. Aslim, and S.B. Aydas, *Mechanisms of cholesterol-lowering effects of*
518 *lactobacilli and bifidobacteria strains as potential probiotics with their bsh gene*
519 *analysis*. J Mol Microbiol Biotechnol, 2014. 24(1): p. 12-8.
- 520 20. Shewry, P.R., et al., *Phytochemical and fiber components in oat varieties in the*
521 *HEALTHGRAIN Diversity Screen*. J Agric Food Chem, 2008. 56(21): p. 9777-84.
- 522 21. Chen, C.Y., et al., *Avenanthramides and phenolic acids from oats are bioavailable and*
523 *act synergistically with vitamin C to enhance hamster and human LDL resistance to*
524 *oxidation*. J Nutr, 2004. 134(6): p. 1459-66.
- 525 22. Liu, S., et al., *Antioxidant Effects of Oats Avenanthramides on Human Serum*.
526 *Agricultural Sciences in China*, 2011. 10(8): p. 1301-1305.
- 527 23. Kern, S.M., et al., *Absorption of hydroxycinnamates in humans after high-bran cereal*
528 *consumption*. J Agric Food Chem, 2003. 51(20): p. 6050-5.
- 529 24. Del Rio, D., et al., *Dietary (poly)phenolics in human health: structures, bioavailability,*
530 *and evidence of protective effects against chronic diseases*. Antioxid Redox Signal,
531 2013. 18(14): p. 1818-92.
- 532 25. Andreasen, M.F., et al., *Esterase activity able to hydrolyze dietary antioxidant*
533 *hydroxycinnamates is distributed along the intestine of mammals*. J Agric Food Chem,
534 2001. 49(11): p. 5679-84.
- 535 26. Couteau, D., et al., *Isolation and characterization of human colonic bacteria able to*
536 *hydrolyse chlorogenic acid*. J Appl Microbiol, 2001. 90(6): p. 873-81.
- 537 27. Raimondi, S., et al., *Role of bifidobacteria in the hydrolysis of chlorogenic acid*.
538 *MicrobiologyOpen*, 2015. 4(1): p. 41-52.
- 539 28. Hughes, S.A., et al., *In vitro fermentation of oat and barley derived beta-glucans by*
540 *human faecal microbiota*. FEMS Microbiol Ecol, 2008. 64(3): p. 482-93.
- 541 29. Connolly, M.L., J.A. Lovegrove, and K.M. Tuohy, *In vitro evaluation of the microbiota*
542 *modulation abilities of different sized whole oat grain flakes*. Anaerobe, 2010. 16(5): p.
543 483-8.
- 544 30. Kedia, G., et al., *In vitro fermentation of oat bran obtained by debranning with a mixed*
545 *culture of human fecal bacteria*. Curr Microbiol, 2009. 58(4): p. 338-42.
- 546 31. Connolly, M.L., K.M. Tuohy, and J.A. Lovegrove, *Wholegrain oat-based cereals have*
547 *prebiotic potential and low glycaemic index*. Br J Nutr, 2012. 108(12): p. 2198-206.

- 548 32. Chappell, A.J., et al., *The effect of in vitro fermentation of oats (Avena sativa) and*
549 *barley (Hordeum vulgare) on the faecal gut microbiota.* Proceedings of the Nutrition
550 Society, 2015. 74(OCE1): p. null-null.
- 551 33. Zhou, A.L., et al., *Whole grain oats improve insulin sensitivity and plasma cholesterol*
552 *profile and modify gut microbiota composition in C57BL/6J mice.* J Nutr, 2015. 145(2):
553 p. 222-30.
- 554 34. Berger, K., et al., *Cereal byproducts have prebiotic potential in mice fed a high-fat diet.*
555 J Agric Food Chem, 2014. 62(32): p. 8169-78.
- 556 35. Immerstrand, T., et al., *Effects of oat bran, processed to different molecular weights of*
557 *beta-glucan, on plasma lipids and caecal formation of SCFA in mice.* Br J Nutr, 2010.
558 104(3): p. 364-73.
- 559 36. Cloetens, L., et al., *Role of dietary beta-glucans in the prevention of the metabolic*
560 *syndrome.* Nutr Rev, 2012. 70(8): p. 444-58.
- 561 37. Ibrugger, S., et al., *Extracted oat and barley beta-glucans do not affect cholesterol*
562 *metabolism in young healthy adults.* J Nutr, 2013. 143(10): p. 1579-85.
- 563 38. Crittenden, R., et al., *In vitro fermentation of cereal dietary fibre carbohydrates by*
564 *probiotic and intestinal bacteria.* Journal of the Science of Food and Agriculture, 2002.
565 82(8): p. 781-789.
- 566 39. Kim, H.J. and P.J. White, *Optimizing the molecular weight of oat beta-glucan for in*
567 *vitro bile acid binding and fermentation.* J Agric Food Chem, 2011. 59(18): p. 10322-
568 8.
- 569 40. Hughes, S.A., et al., *In vitro fermentation by human fecal microflora of wheat*
570 *arabinoxylans.* J Agric Food Chem, 2007. 55(11): p. 4589-95.
- 571 41. Schar, M.Y., et al., *Excretion of Avenanthramides, Phenolic Acids and their Major*
572 *Metabolites Following Intake of Oat Bran.* Mol Nutr Food Res, 2017.
- 573 42. Mills, D.J., et al., *Dietary glycated protein modulates the colonic microbiota towards*
574 *a more detrimental composition in ulcerative colitis patients and non-ulcerative colitis*
575 *subjects.* J Appl Microbiol, 2008. 105(3): p. 706-14.
- 576 43. Shewry, P.R., *HEALTHGRAIN Methods: Analysis of Bioactive Components in Small*
577 *Grain Cereals.* Vol. 290pp. 2010.
- 578 44. Zygmunt, L.C. and S.D. Paisley, *Enzymatic method for determination of (1-->3)(1--*
579 *>4)-beta-D-glucans in grains and cereals: collaborative study.* J AOAC Int, 1993.
580 76(5): p. 1069-82.
- 581 45. McCance, R.A. and E.M. Glaser, *The Energy Value of Oatmeal and the Digestibility*
582 *and Absorption of its Proteins, Fats and Calcium.* British Journal of Nutrition, 1948.
583 2(03): p. 221-228.
- 584 46. Stiverson, J., et al., *Prebiotic Oligosaccharides: Comparative Evaluation Using In*
585 *Vitro Cultures of Infants' Fecal Microbiomes.* Appl Environ Microbiol, 2014. 80(23):
586 p. 7388-97.
- 587 47. Ovreås, L., et al., *Distribution of bacterioplankton in meromictic Lake Saelenvannet,*
588 *as determined by denaturing gradient gel electrophoresis of PCR-amplified gene*
589 *fragments coding for 16S rRNA.* Applied and Environmental Microbiology, 1997.
590 63(9): p. 3367-3373.

- 591 48. Kristensen, K.H., et al., *Gut Microbiota in Children Hospitalized with Oedematous and*
592 *Non-Oedematous Severe Acute Malnutrition in Uganda*. PLoS Negl Trop Dis, 2016.
593 10(1): p. e0004369.
- 594 49. Caporaso, J.G., et al., *QIIME allows analysis of high-throughput community*
595 *sequencing data*. Nat Methods, 2010. 7(5): p. 335-6.
- 596 50. Edgar, R.C., et al., *UCHIME improves sensitivity and speed of chimera detection*.
597 Bioinformatics, 2011. 27(16): p. 2194-200.
- 598 51. McDonald, D., et al., *An improved Greengenes taxonomy with explicit ranks for*
599 *ecological and evolutionary analyses of bacteria and archaea*. Isme j, 2012. 6(3): p.
600 610-8.
- 601 52. Kim, O.S., et al., *Introducing EzTaxon-e: a prokaryotic 16S rRNA gene sequence*
602 *database with phylotypes that represent uncultured species*. Int J Syst Evol Microbiol,
603 2012. 62(Pt 3): p. 716-21.
- 604 53. Dyachkova, M.S., et al., *Draft Genome Sequences of Bifidobacterium angulatum*
605 *GT102 and Bifidobacterium adolescentis 150: Focusing on the Genes Potentially*
606 *Involved in the Gut-Brain Axis*. Genome Announc, 2015. 3(4).
- 607 54. Averina, O.V. and V.N. Danilenko, *Human intestinal microbiota: Role in development*
608 *and functioning of the nervous system*. Microbiology, 2017. 86(1): p. 1-18.
- 609 55. McCleary, B.V., et al., *Determination of insoluble, soluble, and total dietary fiber*
610 *(CODEX definition) by enzymatic-gravimetric method and liquid chromatography:*
611 *collaborative study*. J AOAC Int, 2012. 95(3): p. 824-44.
- 612 56. Knudsen, K.E.B., B.B. Jensen, and I. Hansen, *Digestion of polysaccharides and other*
613 *major components in the small and large intestine of pigs fed on diets consisting of oat*
614 *fractions rich in β -D-glucan*. British Journal of Nutrition, 2007. 70(2): p. 537-556.
- 615 57. Wang, Y., et al., *High Molecular Weight Barley β -Glucan Alters Gut Microbiota*
616 *Toward Reduced Cardiovascular Disease Risk*. Frontiers in Microbiology, 2016. 7: p.
617 129.
- 618 58. Ulsemer, P., et al., *Preliminary safety evaluation of a new Bacteroides xylanisolvens*
619 *isolate*. Appl Environ Microbiol, 2012. 78(2): p. 528-35.
- 620 59. Hosseini, E., et al., *Propionate as a health-promoting microbial metabolite in the*
621 *human gut*. Nutrition Reviews, 2011. 69(5): p. 245-258.
- 622 60. Beckmann, L., O. Simon, and W. Vahjen, *Isolation and identification of mixed linked*
623 *beta -glucan degrading bacteria in the intestine of broiler chickens and partial*
624 *characterization of respective 1,3-1,4-beta -glucanase activities*. J Basic Microbiol,
625 2006. 46(3): p. 175-85.
- 626 61. Fernández-Murga, M.L. and Y. Sanz, *Safety Assessment of Bacteroides uniformis*
627 *CECT 7771 Isolated from Stools of Healthy Breast-Fed Infants*. PLoS ONE, 2016.
628 11(1): p. e0145503.
- 629 62. Dall'Asta, M., et al., *In Vitro Bioaccessibility of Phenolic Acids from a Commercial*
630 *Aleurone-Enriched Bread Compared to a Whole Grain Bread*. Nutrients, 2016. 8(1): p.
631 42.
- 632 63. Gwiazdowska, D., et al., *The impact of polyphenols on Bifidobacterium growth*. Acta
633 Biochim Pol, 2015. 62(4): p. 895-901.

- 634 64. Lindsay, R.F. and F.G. Priest, *Decarboxylation of Substituted Cinnamic Acids by*
635 *Enterobacteria: the Influence on Beer Flavour*. Journal of Applied Bacteriology, 1975.
636 39(2): p. 181-187.
- 637 65. Grbic-Galic, D., *O-Demethylation, dehydroxylation, ring-reduction and cleavage of*
638 *aromatic substrates by Enterobacteriaceae under anaerobic conditions*. Journal of
639 Applied Bacteriology, 1986. 61(6): p. 491-497.
- 640 66. Huang, J., et al., *Different Flavonoids Can Shape Unique Gut Microbiota Profile In*
641 *Vitro*. J Food Sci, 2016. 81(9): p. H2273-9.
- 642 67. Gu, W., et al., *Structural Basis of Enzymatic Activity for the Ferulic Acid*
643 *Decarboxylase (FADase) from Enterobacter sp. Px6-4*. PLoS ONE, 2011. 6(1): p.
644 e16262.
- 645 68. Hunter, W.J., D.K. Manter, and D. van der Lelie, *Biotransformation of ferulic acid to*
646 *4-vinylguaiacol by Enterobacter soli and E. aerogenes*. Curr Microbiol, 2012. 65(6): p.
647 752-7.
- 648 69. Kuntz, S., et al., *Inhibition of Low-Grade Inflammation by Anthocyanins after*
649 *Microbial Fermentation in Vitro*. Nutrients, 2016. 8(7): p. 411.
- 650 70. Wust, P.K., M.A. Horn, and H.L. Drake, *Clostridiaceae and Enterobacteriaceae as*
651 *active fermenters in earthworm gut content*. Isme j, 2011. 5(1): p. 92-106.
- 652 71. Saulnier, D.M., G.R. Gibson, and S. Kolida, *In vitro effects of selected synbiotics on*
653 *the human faecal microbiota composition*. FEMS Microbiol Ecol, 2008. 66(3): p. 516-
654 27.
- 655 72. Kemperman, R.A., et al., *Impact of polyphenols from black tea and red wine/grape*
656 *juice on a gut model microbiome*. Food Research International, 2013. 53(2): p. 659-
657 669.
- 658 73. Eckburg, P.B., et al., *Diversity of the human intestinal microbial flora*. Science, 2005.
659 308.
- 660 74. Menon, R., et al., *Oats-From Farm to Fork*. Adv Food Nutr Res, 2016. 77: p. 1-55.
- 661 75. Englyst, H.N. and J.H. Cummings, *Digestion of the polysaccharides of some cereal*
662 *foods in the human small intestine*. Am J Clin Nutr, 1985. 42(5): p. 778-87.
- 663 76. Martínez, I., et al., *Resistant Starches Types 2 and 4 Have Differential Effects on the*
664 *Composition of the Fecal Microbiota in Human Subjects*. PLoS ONE, 2010. 5(11): p.
665 e15046.
- 666 77. Grootaert, C., et al., *Microbial metabolism and prebiotic potency of arabinoxylan*
667 *oligosaccharides in the human intestine*. Trends in Food Science & Technology, 2007.
668 18(2): p. 64-71.
- 669 78. Macfarlane, S. and G.T. Macfarlane, *Regulation of short-chain fatty acid production*.
670 Proc Nutr Soc, 2003. 62(1): p. 67-72.
- 671 79. Pokusaeva, K., G.F. Fitzgerald, and D. van Sinderen, *Carbohydrate metabolism in*
672 *Bifidobacteria*. Genes & Nutrition, 2011. 6(3): p. 285-306.
- 673 80. Fooks, L.J. and G.R. Gibson, *Probiotics as modulators of the gut flora*. Br J Nutr, 2002.
674 88 Suppl 1: p. S39-49.
- 675 81. Russell, W.R., et al., *Colonic bacterial metabolites and human health*. Curr Opin
676 Microbiol, 2013. 16(3): p. 246-54.

- 677 82. Wolever, T.M., P. Spadafora, and H. Eshuis, *Interaction between colonic acetate and*
678 *propionate in humans*. Am J Clin Nutr, 1991. 53(3): p. 681-7.
- 679 83. Alvaro, A., et al., *Gene expression analysis of a human enterocyte cell line reveals*
680 *downregulation of cholesterol biosynthesis in response to short-chain fatty acids*.
681 IUBMB Life, 2008. 60(11): p. 757-64.
- 682 84. Cheickna, D. and Z. Hui, *Oat Beta-Glucan: Its Role in Health Promotion and*
683 *Prevention of Diseases*. Comprehensive Reviews in Food Science and Food Safety,
684 2012. 11(4): p. 355-365.
- 685 85. Grundy, M.M.L., et al., *The impact of oat structure and β -glucan on in vitro lipid*
686 *digestion*. Journal of Functional Foods, 2017. 38: p. 378-388.
- 687 86. Ziegler, K., et al., *Butyric acid increases transepithelial transport of ferulic acid*
688 *through upregulation of the monocarboxylate transporters SLC16A1 (MCT1) and*
689 *SLC16A3 (MCT4)*. Archives of Biochemistry and Biophysics, 2016. 599: p. 3-12.

690

Table 1. Macronutrient, fibre and phenolic content of oat bran before and after *in vitro* digestion and bioaccessibility (%) through digestion

Components	Before digestion	After digestion	Bioaccessibility %
Amount (g)	60	40	33
Total Carbohydrate (g)	37.8	17.6	53
Available Carbohydrate (g)	32.6	12.5	62
Total dietary fibre(g)	5.2	5.1	2
β-glucan (g)	1.7	1.5	12
Total protein (g)	7	5	29
Fat (g)	3	1.1	63
<i>Free+conjugated polyphenols</i>	6.2	3.2	48
Hydroxybenzoic acids (mg)	1.2	1.1	8
Hydrocinnamic acids (mg)	4.6	2.1	54
Avenanthramides (mg)	0.4	0.1	75
<i>Bound polyphenols</i>	16.4	12.2	26
Hydroxybenzoic acids (mg)	1	0.4	60
Hydrocinnamic acids (mg)	15.4	11.8	23
Avenanthramides (mg)	ND	ND	-
Total polyphenols (mg)	22.6	15.4	32

Bioaccessibility - evaluated following *in vitro* upper gut digestion procedures, we have used the term *bioavailability* to indicate the percentage of compound remaining, thus available to the large gut microbiota. This term was first defined by Dall'Asta et al. in *Nutrients* 2016, 8(1), 42.

Table 2. *In vitro* fermentation treatments

Treatment	β -glucan (mg)	Polyphenol (mg)
OAT1.5	56.67	0.58
OAT4.5	170	1.73
BG (94% of extract)	180	0.1
POLY	ND	1.73
PC	ND	ND
NC	ND	ND

Digested oat bran 1.5g (OAT1.5) and 4.5g (OAT4.5), β -glucan extract (BG), polyphenol mix (POLY), Synergy 1 positive control (PC) and negative control (NC), not detected (ND)

Table 3. Significant changes (FDR $p < 0.05$) in relative abundance (%) of bacterial taxa at 5h, 10h and 24h of *in vitro* batch culture fermentation inoculated with faeces and administered with digested oat bran 1.5g (OAT1.5) and 4.5g (OAT4.5), β -glucan extract (BG), polyphenol mix (POLY), Synergy 1 (PC) and negative control (NC) as substrates. This experiment was conducted three times, with a different faecal donor used for each run.

Phylum	Family	Genus	Species	Time	Treatment ¹						<i>p</i> value
					OAT 1.5	OAT 4.5	BG	POLY	PC	NC	
Actinobacteria	Bifidobacteriaceae	<i>Bifidobacterium</i>	<i>adolescentis</i>	5h	2.73±1.9	0.82±0.7	0.31±0.1	0.38±0.1	18.48±3.5*	0.39±0.2	0.001
Firmicutes	Ruminococcaceae	<i>Ruminococcus</i>	<i>unassigned</i>	5h	0.2±0*	0.1±0.1	0.52±0.1	0.67±0.1	0.32±0.1	0.74±0.1	0.03
Firmicutes	Lachnospiraceae	<i>Coproccoccus</i>	<i>unassigned</i>	5h	0.2±0.1*	0.12±0*	0.48±0.2	0.44±0	0.76±0.3	0.54±0	0.02
Proteobacteria	Enterobacteriaceae	<i>unassigned</i>	<i>unassigned</i>	5h	55.46±8.4*	61.2±9*	23±6	31.9±8.6	11.33*±3.7	25.9±11	0.05
Actinobacteria	Bifidobacteriaceae	<i>Bifidobacterium</i>	<i>unassigned</i>	10h	1.17±10.1*	1.1±0.9	0.19±0.1	0.17±0.1	4.12±2.1*	0.27±0.1	0.03
Actinobacteria	Bifidobacteriaceae	<i>Bifidobacterium</i>	<i>adolescentis</i>	10h	1.93±0.5*	0.66±0.5	0.19±0.1	0.27±0.1	6.7±2.6*	0.26±0	0.02
Firmicutes	Ruminococcaceae	<i>unassigned</i>	<i>unassigned</i>	10h	4.26±5	0.05±0*	8.96±2.8	8.68±1.7	9.76±10	8.85±0.8	0.03
Firmicutes	Lachnospiraceae	<i>unassigned</i>	<i>unassigned</i>	10h	5.14±4	0.08±0*	10.1±2.4	8.14±5	5.11±1.2	10.1±0.2	0.03
Firmicutes	Lachnospiraceae	<i>Blautia</i>	<i>unassigned</i>	10h	0.35±0.1*	0.02±0*	1.39±0.3	1.21±0.4	1.15±0.8	1.14±0.4	0.03
Proteobacteria	Enterobacteriaceae	<i>unassigned</i>	<i>unassigned</i>	10h	50.2±12.2	67.2±14*	12.75±6*	36.1±9.2	6.47±3.7*	29.2±1.4	0.02
Actinobacteria	Bifidobacteriaceae	<i>Bifidobacterium</i>	<i>adolescentis</i>	24h	2.03±0.1*	1.91±1.8	0.31±0.1	0.31±0.2	6.87±3.7	0.41±0	0.05

¹ The difference in relative abundance of taxa between treatments within the same time points was assessed by ANOVA. The *p*-value after correction for multiple tests (69 taxa) with the FDR method.

* indicate significant differences ($p < 0.05$) to NC at the same time point. Values are mean ± SE

Figure Legend


Figure 1. Principal coordinates analysis (PCoA) plots of 16S rRNA gene profiles based on (A) unweighted and (B) weighted phylogenetic Unifrac distance matrices calculated from 10 rarefied OTU tables (9800 reads per samples) unweighted (A) showing clear clustering according to donors (ANOSIM test, $p = 0.01$). The degree of variation between 10 jackknifed replicates of PCoA is displayed with confidence ellipsoids around each sample. Weighted (B) quantitative information used to generate the bacterial relative abundance of donors showed no clear clustering (ANOSIM test, $p > 0.05$) for a whole dataset (24h *in vitro* batch culture fermentation inoculated with 3 healthy donors faeces and administered with digested oat bran 1.5g (OAT1.5) and 4.5g (OAT4.5), β -glucan extract (BG), polyphenol mix (POLY), Synergy 1 (PC) and negative control (NC) as the substrates). Each colour represents a different donor

Figure 2. Changes in bacterial phyla (relative abundances (%)) 0, 5, 10 and 24h *in vitro* batch culture fermentation. This experiment was conducted three times, with a different faecal donor used for each run. The media was supplemented with digested oat bran 1.5g (OAT1.5) and 4.5g (OAT4.5), β -glucan extract (BG), polyphenol mix (POLY), Synergy 1 (PC) and negative control (NC). Samples were analysed at 0, 5, 10, 24 h. Values are mean (%).


Figure 3. Change in concentrations of acetic acid (A), propionic acid (B), butyric acid (C) from negative control (mM) throughout 24h *in vitro* batch culture fermentation. This experiment was conducted three times, with a different faecal donor used for each run. The media supplemented with digested oat bran 1.5g (OAT1.5) and 4.5g (OAT4.5), β -glucan extract (BG), polyphenol mix (POLY), Synergy 1 (PC) as the substrates. Samples were analysed at 0, 5, 10, 24 h

Figure 1.

A



B



■ Donor 1 ■ Donor 2 ■ Donor 3

Figure 2.





Figure 3.



Values are reported after subtracted of negative control value

*Significant from initial value * $p < 0.05$, ** $p < 0.01$; Significant from 5h value # $p < 0.05$, ## $p < 0.01$,*

Significant from 10h value ^ $p < 0.05$, ^^ $p < 0.01$; Significant from negative control value at the same time point & $p < 0.05$, && $p < 0.01$