

*Integrated microscopy approaches in  
archaeobotany: proceedings of the 2016  
and 2017 workshops, University of  
Reading, UK*

Article

Accepted Version

Banerjea, R. Y., Barnett, C., Morandi, L. and Portillo, M.  
(2020) Integrated microscopy approaches in archaeobotany:  
proceedings of the 2016 and 2017 workshops, University of  
Reading, UK. *Environmental Archaeology: The Journal of  
Human Palaeoecology*, 25 (2). pp. 131-134. ISSN 1461-4103  
doi: <https://doi.org/10.1080/14614103.2019.1593654> Available  
at <http://centaur.reading.ac.uk/83200/>

It is advisable to refer to the publisher's version if you intend to cite from the  
work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1080/14614103.2019.1593654>

Publisher: Taylor & Francis

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

## **CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

## Editorial: Proceedings of the 2016 and 2017 workshops on ‘Integrated Microscopy Approaches in Archaeobotany’

Rowena Y. Banerjea<sup>1</sup>, Catherine Barnett<sup>1</sup>, Lionello Morandi<sup>1,2</sup>, Marta Portillo<sup>1</sup>

<sup>1</sup> University of Reading, Department of Archaeology, Reading, UK

<sup>2</sup> Eberhard Karl University of Tübingen, Institute of Scientific Archaeology, Tübingen, Germany

### ORCID ID

Rowena Y. Banerjea 0000-0002-1786-357X

Catherine Barnett 0000-0002-6156-1640

Lionello Morandi 0000-0001-8840-6096

Marta Portillo 0000-0002-2703-031X

The collection of papers in this special issue arises from research presented at the 2016 and 2017 workshops in ‘Integrated Microscopy Approaches in Archaeobotany’ held at the University of Reading, UK. The aim of the workshops was to bring together students and specialists in geoarchaeology, biogeochemistry, plant macroremains, anthracology, palynology, non-pollen palynomorphs and phytoliths who are working in universities, major heritage organisations, curatorial bodies, and commercial archaeology companies in order to share research and skills. Since then, the IMAA workshops have been successfully celebrated in Reading every year, testifying to the growing interest for archaeobotanical microscopy approaches in conjunction with other proxies, and particularly from the field of geosciences.

Archaeobotany is the study of archaeological plant remains, which include plant macroremains, such as seeds and charcoal, and a range of associated microfossils, such as phytoliths, pollen, spores and starch grains. These papers showcase a range of microscopy approaches that can be applied to examine and interpret botanical and organic assemblages in the archaeological record, and to address a range of topical research themes and current debates in archaeology, such as environmental management, human responses to environmental change and sustainable life-ways spanning a broad geographical and chronological time periods. Soil micromorphology is a technique that enables the formation and post-depositional processes of archaeological deposits to be understood by reconstructing sediment histories. It allows archaeobotanical remains to be examined within their depositional context to provide a micro-contextual interpretation of the evidence, and to understand different types of archaeobotanical remains in a range of preservation conditions (e.g. Matthews 2010; Vrydaghs *et al.* 2016; Ismail-Meyer 2017; Ismail-Meyer *et al.* 2018). As with many other proxies, formation processes and taphonomy are key issues in the study of any type of plant remains (e.g. Van der Veen 2007; Matthews 2010), and this is reflected in several contributions of this volume.

Banerjea *et al.* and Borderie *et al.* both address an absence of multi-proxy environmental archaeological studies on European castle sites from the medieval period, and particularly to

understand the early development of these sites. In both these papers, the integrated results have allowed the potential of these proxies to be powerfully exploited to show new archaeobotanical perspectives on the function and development of these castle complexes and their relationships with the castle's hinterland.

Banerjea *et al.* use soil micromorphology, plant macroremains, phytoliths and palynology results from two castles in the Baltic (Elbląg, Poland, and Karksi, Estonia) to show changes in animal husbandry through time, whereabouts within the castles the animals were stabled, and livestock alimentation. The study represents an important investigation of deposits from medieval castles, particularly from the period of active Crusading that are rarely revealed through excavation and in the past have rarely been subjected to an integrated environmental investigation.

Borderie *et al.* use soil micromorphology, plant macroremains and phytoliths, including their morphometric analysis, to understand floor preparation materials, maintenance practices, and domestic, metallurgical and animal husbandry activities to provide new evidence of the richness and the complexity of the town, and of the activities which could take place within early *castra*. Morphometric analysis of phytoliths in micromorphological thin-sections showed cultivated cereals which were processed *in situ*, such as *Triticum durum*, *Secale cereale* and *Hordeum vulgare*.

The paper by Vrydaghs and Devos propose a system for the simultaneous observation and description of phytoliths in thin sections of soils and archaeological sediments. It presents three aspects of phytoliths in thin section, namely VPC index: (V) visibility, expressing to what extent these are masked and/or surrounded by fine material; (P) preservation, as indicator for the physico-chemical alterations that affected them; (C) colour, describing the extent to which the phytoliths are coated by other particles and indicator for the charring of the organic material surrounding them. This paper presents an important contribution towards the standardised description and recording of phytoliths in thin section to facilitate comparison and interpretations of microfossil datasets from different sites and research areas.

A good knowledge of livestock management and herding practices is crucial when investigating the subsistence of farming societies. These aspects do not always have a good visibility within the archaeological record, so that archaeobotanical evidence should be employed to shed light on them, along with other analytical techniques including contextual micromorphology as above outlined, and direct microscopic dung indicators such as calcitic spherulites, coprophilous fungal spores, isotopic and biomolecular signatures, along with zooarchaeological remains from different origin. Of particular interest for unveiling pastoral and agro-pastoral activities in archaeology is understanding the nature and taphonomy of animal dung remains, which is in this issue explored in archaeological and modern ethnographical contexts through the use of direct and indirect proxies of various origin, including in vegetal micro-remains such as opal phytoliths and calcium oxalates, as well as faecal indicators such as fungal spores and calcitic spherulites that form in the digestive system of a variety of animals (e.g. Shahack-Gross 2011; Friesem 2016, and references therein).

Two papers deal with both ethnoarchaeological plant and dung microfossil evidence for livestock management (Dalton and Ryan; Morandi).

The study of Dalton and Ryan on a New Kingdom pharaonic settlement in Sudan combines dung spherulite and phytolith analyses of desiccated and charred ovicaprid dung pellets as well as faecal material in thin-sections, with further support provided by micromorphological evidence and chemical signatures. Their experimentally produced datasets from modern reference materials suggest differences in dietary calcium intake and feed Ca availability implicated in dung spherulite

crystallization within ovicaprid digestive tracts, thus providing new insights into spherulite formation aspects poorly-defined to date.

Analysing a sequence from a recent pastoral cave context in northwestern Italy, the paper by Morandi also follows an integrated approach combining direct faecal microfossil indicators from calcitic spherulites and coprophilous fungal spores, which are more widely utilised in wetland palaeoecology to assess past herbivore presence (Baker *et al.* 2013; Johnson *et al.* 2015). This study highlights the contribution of ethnoarchaeological approaches to understand microfossil taphonomy and their distributions better within penning deposits.

The paper by García-Suárez, Portillo and Matthews presents micro-contextual studies in open site areas through simultaneous examination in thin section micromorphology and integrated phytolith and dung spherulite analyses for the identification of early animal management strategies in the Konya Plain, a key region in the origins and spread of Neolithic innovations in Central Anatolia, Turkey (e.g. Hodder 2006; Baird *et al.* 2018). Dung assemblages are identified at high-resolution studies from extensive midden deposits at the early agricultural site of Boncuklu, as well as at the later occupations of Çatalhöyük and the Late Neolithic settlement of the Pınarbaşı rockshelter, thus proving direct evidence for animal management, diet and ecological diversity, as well as for site formation and organization of the space of early settled life. In sum, all three papers within this section show the potential of archaeobotanical microfossils and other micro-remains for reconstructing livestock management, animal diet and potential indicators of seasonality.

No proxy record has the ability to enable reconstruction of past landscapes in isolation. However, substantial on-site archaeobotanical macrofossil assemblages including charcoal do offer the opportunity to examine aspects of the local landscape and presence/absence of exploited habitats across the wider source area (Théry-Parisot *et al.* 2010; Asouti and Austen 2005, Chabal 1997). The presence of charred wood on an archaeological site relies on it having been gathered and transformed through use or discard, and so its examination can also provide insights into a range of human activities and targeted resource exploitation (Théry-Parisot *et al.* 2010; Smart and Hoffman 1988). Despite these opportunities for knowledge gain, analysis of charcoal, and to a degree other on-site macroscopic proxies, such as mollusca and coleoptera, are often overlooked, underestimated or felt to be a poor relation of plant macrofossil analysis as a tool in archaeological studies.

Reflection is made in the paper presented in this volume by Barnett on whether systematic, large-scale, analysis of wood charcoal is of significant value in understanding landscape, resource use and lifestyles at and around Iron Age Silchester and the wider implications of that reflection are considered. Barnett's paper shows that anthracology has high potential to inform not only on the presence of tree and shrub taxa in the past but also on the structure and location of former woodlands, heathlands and wetland fringes.

Lastly, Portillo and colleagues present an extensive and up-to-date review of the literature dedicated to morphometric investigations in archaeobotany. Over the last few decades, morphometrics have particularly contributed to the study of the domestication and spread of many crops around the world, such as cereals and legumes, underground storage organs (USO), and fruits, including olives, grapes, dates, and bananas (e.g. Terral *et al.* 2004, 2010; Willcox 2004; Fuller 2007, 2017; Ball *et al.* 2016, and references therein). This paper presents an overview on current methodologies, recent applications, and advances in the use of morphometrics for various types of plant remains (mainly seeds, pollen, phytoliths, and starch grains), its applications for improving taxonomic resolution, and

discusses its contributions to addressing major research questions, challenges and possible future directions in archaeobotanical research.

Collectively, the papers explore ways to address the challenges of integrating and presenting datasets that were recovered from a) multiple sites and b) through different types of excavation, often within the same project. Arising from these issues stems a necessity to understand the sampling and data interrogation requirements of fellow specialists. Overall, the articles included in this volume illustrate a range of perspectives by which microscopy approaches in archaeobotany contribute to address main research questions and current debates in archaeology, making these studies a valuable research tool especially when integrated with other proxies.

## Disclosure statement

No potential conflict of interest was reported by the authors.

## Funding

Funding, administrative and technical support has been provided by the School of Archaeology, Geography and Environmental Science for the successful development of the IMAA workshops, and we thank the Association for Environmental Archaeology, Beta Analytic and Quaternary Scientific (QUEST) for their sponsorship. The 2017 workshop also received funding from the University of Reading's Endowment Fund. Rowena Banerjea's research was carried out within the framework of The Ecology of Crusading project, which received funding from the European Union's Seventh Framework Programme (FP7/ 2007-2013) under grant agreement No. 263735. Catherine Barnett's research was funded by the Calleva Foundation as part of the wider Silchester Environs project. The work of Marta Portillo was carried out within the framework of the EU Horizon 2020 MICROARCHEODUNG project. This project has received funding from the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement No H2020-MSCA-IF-2015-702529.

## References

Asouti, E., and P. Austin 2005. Reconstructing woodland vegetation and its relation to human societies, based on the analysis and interpretation of archaeological wood charcoal macro-remains. *Environmental Archaeology* 10: 1-18.

Baird, D., A. Fairbairn, E. Jenkins, L. Martin, C. Middleton, J. Pearson, E. Asouti, Y. Edwards, C. Kabukcu, G. Mustafaoğlu, N. Russell, O. Bar-Yosef, G. Jacobsen, X. Wu, A. Baker, S. Elliott. 2018. "Agricultural origins on the Anatolian Plateau." *Proceedings of the National Academy of Sciences*. doi: <https://doi.org/10.1073/pnas.1800163115>.

Baker, A. G., Cornelissen, P., Bhagwat, S. A., Vera, F. W., Willis, K. J. 2016. Quantification of population sizes of large herbivores and their long-term functional role in ecosystems using dung fungal spores. *Methods in Ecology and Evolution* 7(11): 1273-1281.

- Ball, T.B., A. Davis, R.R. Evett, J.L. Ladwig, M. Tromp, W.A. Out, and M. Portillo. 2016. "Morphometric analysis of phytoliths: recommendations towards standardization from the International Committee for Phytolith Morphometrics". *Journal of Archaeological Science* 68: 106–111.
- Chabal, L., 1992. La représentativité paléoécologique des charbons de bois archéologiques issus du bois de feu. In: Vernet, J.L. (Ed.), *Les charbons de bois les anciens écosystèmes et le rôle de l'Homme*. Bul. de la Soc. Bot. de France, 139, 213–236.
- Friesem, D.E. 2016. "Geo-ethnoarchaeology in action". *Journal of Archaeological Science* 70: 145–157.
- Fuller, D.Q. 2007. "Contrasting patterns in crop domestication and domestication rates: recent archaeobotanical insights from the Old World". *Annals of Botany* 100 (5): 903–924.
- Fuller, D.Q., Colledge, S., Murphy, C., and C. Stevens. 2017. "Sizing up cereal variation: patterns in grain evolution revealed in chronological and geographical comparisons". In *Miscelánea en homenaje a Lydia Zapata Peña (1965-2015)*, edited by J. FernándezEraso, J.A. MujikaAlustiza, A. ArrizabalagaValbuena, and M. GarcíaDíez, 131–149. Bilbao: Universidad del País Vasco.
- Hodder, I., 2006. *The Leopard's Tale*. Thames & Hudson, London.
- Ismail-Meyer, K. 2017. Plant remains. In: Nicosia, C., Stoops, G. *Archaeological Soil and Sediment Micromorphology*. John Wiley & Sons, Chichester, pp. 121-135
- Ismail-Meyer, K., Stolt, M.H. & Lindbo, D.L. 2018. Soil organic matter. In: Stoops, G., Marcelino, V., Mees, F. (eds.) *Interpretation of micromorphological features of soils and regoliths*. Elsevier, Amsterdam, pp 471-512.
- Johnson, C. N., Rule, S., Haberle, S. G., Turney, C. S., Kershaw, A. P., & Brook, B. W. 2015. Using dung fungi to interpret decline and extinction of megaherbivores: problems and solutions. *Quaternary Science Reviews* 110: 107-113.
- Matthews, W. 2010. Geoarchaeology and taphonomy of plant remains in early urban environments in the Ancient Near East. *Quaternary International*, 214 (1-2). pp. 98-113. doi: <https://doi.org/10.1016/j.quaint.2009.10.019>
- Shahack-Gross, R. 2011. Herbivorous livestock dung: formation, taphonomy, methods for identification, and archaeological significance. *Journal of Archaeological Science* 38 (2):205-18.
- Smart, T. L. and Hoffman, E. S. 1988. Environmental interpretation of archaeological charcoal, in Hastorf, C. A. and Popper, V. S. (eds.), *Current Paleoethnobotany*. Chicago and London: University of Chicago Press, 165-205.
- Terral J.F, N. Alonso, R. Buxó i Capdevila, N. Chatti, L. Fabre, G. Fiorentino, Ph. Marival, G. Pérez Jordá, B. Pradat, N. Rovira, and P. Alibert. 2004. "Historical biogeography of olive domestication (*Olea europaea* L.) as revealed by geometrical morphometry applied to biological and archaeological material". *Journal of Biogeography* 31: 63–77.
- Terral, J.F., E. Tabard, L. Bouby, S. Ivorra, T. Pastor, I. Figueiral, S. Picq, J.B. Chevance, C. Jung, L. Fabre, C. Tardy, M. Compan, R. Bacilieri, T. Lacombe, and P. This. 2010. Evolution and history of grapevine (*Vitis vinifera*) under domestication: new morphometric perspectives to understand seed domestication syndrome and reveal origins of ancient European cultivars". *Annals of Botany* 105 (3): 443–455

Théry-Parisot, I., L. Chabal, and J. Chravzez. 2010. Anthracology and taphonomy, from wood gathering to charcoal analysis: A review of the taphonomic processes modifying charcoal assemblages, in archaeological contexts. *Palaeogeography, Palaeoclimatology, Palaeoecology* 291: 142–153.

Van der Veen, M. 2007. Formation processes of desiccated and carbonized plant remains – the identification of routine practice. *Journal of Archaeological Science* 34: 968–990

Vrydaghs, L., Ball, T.B., Devos, Y. 2016. Beyond redundancy and multiplicity. Integrating phytolith analysis and micromorphology to the study of Brussels Dark Earth. *Journal of Archaeological Science* 68, 79-88 <https://doi.org/10.1016/j.jas.2015.09.004>

Willcox, G. 2004. “Measuring grain size and identifying Near Eastern cereal domestication: evidence from the Euphrates valley.” *Journal of Archaeological Science* 31(2): 145–150.