

Symmetries of the Feinberg-Zee random hopping matrix

Article

Accepted Version

Hagger, R. (2015) Symmetries of the Feinberg-Zee random hopping matrix. *Random Matrices: Theory and Applications*, 4 (4). 1550016. ISSN 2010-3271 doi: 10.1142/S2010326315500161 Available at <https://centaur.reading.ac.uk/84017/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1142/S2010326315500161>

Publisher: World Scientific

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Symmetries of the Feinberg-Zee Random Hopping Matrix

Raffael Hagger*

October 29, 2018

Abstract

We study the symmetries of the spectrum of the Feinberg-Zee Random Hopping Matrix introduced in [6] and studied in various papers thereafter (e.g. [1], [2], [3], [7], [9]). In [3], Chandler-Wilde and Davies proved that the spectrum of the Feinberg-Zee Random Hopping Matrix is invariant under taking square roots, which implied that the unit disk is contained in the spectrum (a result already obtained slightly earlier in [1]). In a similar approach we show that there is an infinite sequence of symmetries at least in the periodic part of the spectrum (which is conjectured to be dense). Using these symmetries and the result of [3], we can exploit a considerably larger part of the spectrum than the unit disk. As a further consequence we find an infinite sequence of Julia sets contained in the spectrum. These facts may serve as a part of an explanation of the seemingly fractal-like behaviour of the boundary.

2010 Mathematics Subject Classification: Primary 47B80; Secondary 47A10, 47B36

Keywords: random hopping, random operator, spectrum, tridiagonal, periodic, symmetry

1 Introduction

In recent years some progress has been made in the study of non-self-adjoint random operators (see [1], [2], [3], [7], [9] and references therein). However, still a lot of questions remain open even in the tridiagonal case. In particular, the spectrum of most tridiagonal random operators is yet unknown. Although the techniques used here are possibly also suited for more general types of operators, we focus on the Feinberg-Zee Random Hopping Matrix [6] here. It is defined as follows:

$$A^b := \begin{pmatrix} \ddots & \ddots & & & \\ & \ddots & 0 & 1 & \\ & & b_0 & 0 & 1 \\ & & & b_1 & 0 & \ddots \\ & & & & \ddots & \ddots \end{pmatrix} \in \mathcal{L}(\ell^2(\mathbb{Z}))$$

for a random sequence $b \in \{\pm 1\}^{\mathbb{Z}}$, i.e. a sequence with randomly (i.i.d.) distributed entries. Similarly we define A^c for arbitrary sequences $c \in \{\pm 1\}^{\mathbb{Z}}$. The spectrum

$$\text{sp}(A^b) := \{\lambda \in \mathbb{C} : A^b - \lambda I \text{ is not invertible}\}$$

of A^b is independent of the sequence b in the following sense:

*raffael.hagger@tuhh.de

Proposition 1. (e.g. [2, Lemma 2.3, Theorem 2.5]) For a sequence of i.i.d. random variables $(b_j)_{j \in \mathbb{Z}}$ taking values in $\{\pm 1\}$ with non-zero probability each, the spectrum of A^b is given by

$$\text{sp}(A^b) = \Sigma := \bigcup_{c \in \{\pm 1\}^{\mathbb{Z}}} \text{sp}(A^c) \quad (1)$$

almost surely.

In [1] Chandler-Wilde, Chonchaiya and Lindner discovered a beautiful connection to the Sierpinski triangle. This connection was then used to show that the unit disk is contained in Σ , which disproved earlier conjectures that Σ may be of fractal dimension. In [2] the same authors also gave an upper bound using the numerical range. This upper bound was further improved in [7]. An image of Σ (or more precisely: what it is conjectured to be) and the upper bounds computed in [2] and [7] are provided in Figure 1.




Figure 1: The conjectured shape of Σ (black), the upper bound computed in [2] (red) and the upper bound computed in [7] (blue).

Another way of deriving lower bounds is using equation (1) directly. The spectra of many operators on the right hand side can be computed explicitly. This is in particular the case for periodic operators. We call an operator $A \in \mathcal{L}(\ell^2(\mathbb{Z}))$ (not necessarily tridiagonal) m -periodic if $A_{i,j} = A_{i+m,j+m}$ for all $i, j \in \mathbb{Z}$. In the tridiagonal case, we will also use the notation A_{per}^k for

$k \in \{\pm 1\}^m$, i.e.

$$A_{per}^k = \begin{pmatrix} \ddots & \ddots & & & & & \\ \ddots & 0 & 1 & & & & \\ & k_m & 0 & 1 & & & \\ & & k_1 & \ddots & \ddots & & \\ & & & \ddots & \ddots & 1 & \\ & & & & k_m & 0 & 1 \\ & & & & & k_1 & 0 & \ddots \\ & & & & & & \ddots & \ddots \end{pmatrix} \in \mathcal{L}(\ell^2(\mathbb{Z})).$$

Note that k is not unique because $(k_1, \dots, k_m) \in \{\pm 1\}^m$ and $(k_1, \dots, k_m, k_1, \dots, k_m) \in \{\pm 1\}^{2m}$ define the same operator. We will make use of this fact later on.

The good thing about periodic operators is that they can be diagonalized in some sense.

Proposition 2. (e.g. [4, Theorem 4.4.9])

Let $m \in \mathbb{N}$, let $A \in \mathcal{L}(\ell^2(\mathbb{Z}))$ be an m -periodic operator and define

$$a_r := \begin{pmatrix} A_{rm+1,1} & \dots & A_{rm+1,m} \\ \vdots & \ddots & \vdots \\ A_{rm+m,1} & \dots & A_{rm+m,m} \end{pmatrix}$$

for all $r \in \mathbb{Z}$. If $\sum_{r \in \mathbb{Z}} \|a_r\| < \infty$, then A is unitarily equivalent to the (generalized) multiplication operator $M_a \in \mathcal{L}(L^2([0, 2\pi), \mathbb{C}^m))$. The function $a \in L^\infty([0, 2\pi), \mathbb{C}^{m \times m})$ is called the symbol of A and given by

$$a(\varphi) = \sum_{r \in \mathbb{Z}} a_r e^{ir\varphi} \quad (\varphi \in [0, 2\pi)).$$

Moreover, $\text{sp}(A) = \{\lambda \in \mathbb{C} : \det(a(\varphi) - \lambda I_m) = 0 \text{ for some } \varphi \in [0, 2\pi)\} = \bigcup_{\varphi \in [0, 2\pi)} \text{sp}(a(\varphi))$.

In the following, we will always have a finite number of non-vanishing coefficients a_r . Thus we will not have to worry about the convergence of $\sum_{r \in \mathbb{Z}} \|a_r\|$. Proposition 2 enables us to compute the spectra of periodic operators explicitly (numerically). We call the set

$$\pi_\infty := \bigcup_{m \in \mathbb{N}} \bigcup_{k \in \{\pm 1\}^m} \text{sp}(A_{per}^k)$$

the periodic part of Σ . Another related set is the set of eigenvalues of finite matrices of this kind. For $n \in \mathbb{N}$ and $k \in \{\pm 1\}^n$ we define

$$A_{fin}^k := \begin{pmatrix} 0 & 1 & & & \\ k_1 & \ddots & \ddots & & \\ & \ddots & \ddots & 1 & \\ & & k_n & 0 & \end{pmatrix} \in \mathbb{C}^{(n+1) \times (n+1)}.$$

Then we call the set

$$\sigma_\infty := \bigcup_{n \in \mathbb{N}} \bigcup_{k \in \{\pm 1\}^n} \text{sp}(A_{fin}^k)$$

the finite part of Σ . It is clear by equation (1) that π_∞ is a subset of Σ . Furthermore, it was shown in [2] that $\sigma_\infty \subset \pi_\infty$ holds. It was then conjectured in [1] that $\text{clos}(\sigma_\infty) = \text{clos}(\pi_\infty) = \Sigma$ and that Σ is a simply connected set which is the closure of its interior and which has a fractal boundary. That the first equality holds was recently shown in [8] whereas the second equality and the other assertions remain open. Some light was shed on this question in [3], where it was proved that $\mathbb{D} \subset \text{clos}(\pi_\infty)$. Combined with the result of [8] and equation (1), this implies

$$\mathbb{D} \subset \text{clos}(\sigma_\infty) = \text{clos}(\pi_\infty) \subset \Sigma. \quad (2)$$

The result of Chandler-Wilde and Davies in [3] is based on the observation that π_∞ (and also Σ) is invariant under taking square roots, i.e.

$$\lambda^2 \in \pi_\infty \implies \lambda \in \pi_\infty.$$

In this paper we extend this result to an infinite number of symmetries. We prove that for $p \in S$, where S is an infinite set of polynomials made precise below, the following holds:

$$p(\lambda) \in \pi_\infty \implies \lambda \in \pi_\infty.$$

In other words, π_∞ (and hence also $\text{clos}(\pi_\infty)$) is invariant under taking roots of polynomials $p \in S$. This implies that (2) can be extended to

$$p^{-1}(\mathbb{D}) \subset p^{-1}(\text{clos}(\sigma_\infty)) = p^{-1}(\text{clos}(\pi_\infty)) \subset \text{clos}(\pi_\infty) \subset \Sigma$$

for all $p \in S$. Thus

$$\bigcup_{p \in S} p^{-1}(\mathbb{D}) \subset \text{clos}(\sigma_\infty) = \text{clos}(\pi_\infty) \subset \Sigma. \quad (3)$$

This improvement in comparison with (2) is significant as Figure 2 shows.

Clearly, this construction can also be iterated, i.e.

$$p^{-n}(\mathbb{D}) \subset p^{-n}(\text{clos}(\sigma_\infty)) = p^{-n}(\text{clos}(\pi_\infty)) \subset \text{clos}(\pi_\infty) \subset \Sigma$$

and hence

$$\bigcup_{n \in \mathbb{N}} p^{-n}(\mathbb{D}) \subset \bigcup_{n \in \mathbb{N}} p^{-n}(\text{clos}(\sigma_\infty)) = \bigcup_{n \in \mathbb{N}} p^{-n}(\text{clos}(\pi_\infty)) \subset \text{clos}(\pi_\infty) \subset \Sigma.$$

This implies that Σ contains an infinite sequence of (presumably filled) Julia sets (see Remark 12 below), e.g. the set indicated in Figure 3.

These two approaches can also be combined as follows. Let T be the closure of S with respect to composition, i.e.

$$T = \{q: \mathbb{C} \rightarrow \mathbb{C} : q = p_1 \circ \dots \circ p_n \text{ for } p_1, \dots, p_n \in S, n \in \mathbb{N}\}.$$

Then

$$\bigcup_{q \in T} q^{-1}(\mathbb{D}) \subset \text{clos}(\sigma_\infty) = \text{clos}(\pi_\infty) \subset \Sigma.$$




Figure 2: The left-hand side of (3) (polynomials in S up to order 15 to be precise), the upper bound to Σ computed in [7] and the unit circle as a reference.

In this way one can construct even more Julia sets that are contained in Σ . This richness of symmetries might be a part of an explanation of the seemingly fractal boundary of Σ . Surely, this observation needs further investigation.

Note that in [3] it was also shown that

$$\lambda \in \Sigma \implies \pm\sqrt{\lambda} \in \Sigma.$$

Here we only have the (possibly weaker) statement

$$\lambda \in \Sigma \implies p^{-1}(\{\lambda\}) \cap \Sigma \neq \emptyset$$

for $p \in S$.

In addition to the polynomial symmetries mentioned above, there are also the following symmetries (see [2, Lemma 3.4]):

$$\lambda \in \pi_\infty \implies i\lambda, \bar{\lambda} \in \pi_\infty \quad \text{and} \quad \lambda \in \Sigma \implies i\lambda, \bar{\lambda} \in \Sigma.$$

We start with some well-known preparatory results and end up with the two main theorems of this paper. A short list of polynomials in S and selected pictures of new subsets of Σ are provided at the end.




Figure 3: The filled Julia set corresponding to $p(\lambda) = \lambda^3 - \lambda$.

2 Symmetries

Let $m \in \mathbb{N}$ and $k \in \{\pm 1\}^m$. Then we denote the corresponding m -periodic operator

$$\begin{pmatrix} \ddots & \ddots & & & & & \\ \ddots & 0 & 1 & & & & \\ & k_m & 0 & 1 & & & \\ & & k_1 & \ddots & \ddots & & \\ & & & \ddots & \ddots & 1 & \\ & & & & k_m & 0 & 1 \\ & & & & & k_1 & 0 & \ddots \\ & & & & & & \ddots & \ddots \end{pmatrix} \in \mathcal{L}(\ell^2(\mathbb{Z}))$$

by A_{per}^k . By Proposition 2, we can use the symbol a^k to compute the spectrum of A_{per}^k . In our case, due to tridiagonality, the formula can be simplified as follows.

Lemma 3. *Let $m \in \mathbb{N}$, $k \in \{\pm 1\}^m$ and let a^k denote the symbol of A_{per}^k . Then the only φ -dependent term in the characteristic polynomial of $a^k(\varphi)$ is the term of order zero. More precisely,*

there exists a polynomial $p_k: \mathbb{C} \rightarrow \mathbb{C}$ of degree m such that

$$\det(a^k(\varphi) - \lambda I_m) = (-1)^m \left(p_k(\lambda) - e^{i\varphi} \prod_{j=1}^m k_j - e^{-i\varphi} \right) \quad (4)$$

for all $\varphi \in [0, 2\pi)$. The polynomial p_k is monic and given by

$$p_k(\lambda) = (-1)^m \left(\det \begin{pmatrix} -\lambda & 1 & & \\ k_1 & \ddots & \ddots & \\ & \ddots & \ddots & 1 \\ & & k_{m-1} & -\lambda \end{pmatrix} - k_m \det \begin{pmatrix} -\lambda & 1 & & \\ k_2 & \ddots & \ddots & \\ & \ddots & \ddots & 1 \\ & & k_{m-2} & -\lambda \end{pmatrix} \right).$$

Furthermore, p_k is an even (odd) function if m is even (odd).

Proof. The symbol a^k is given by

$$a^k(\varphi) = \begin{pmatrix} 0 & 1 & & k_m e^{i\varphi} \\ k_1 & \ddots & \ddots & \\ & \ddots & \ddots & 1 \\ e^{-i\varphi} & & k_{m-1} & 0 \end{pmatrix}.$$

Using Laplace's formula, we get

$$\begin{aligned} \det(a^k(\varphi) - \lambda I_m) &= -\lambda \det \begin{pmatrix} -\lambda & 1 & & \\ k_2 & \ddots & \ddots & \\ & \ddots & \ddots & 1 \\ & & k_{m-1} & -\lambda \end{pmatrix} \\ &\quad - k_1 \left(\det \begin{pmatrix} -\lambda & 1 & & \\ k_3 & \ddots & \ddots & \\ & \ddots & \ddots & 1 \\ & & k_{m-1} & -\lambda \end{pmatrix} + (-1)^{m-2} k_m e^{i\varphi} \prod_{j=2}^{m-1} k_j \right) \\ &\quad + (-1)^{m-1} e^{-i\varphi} \left(1 + (-1)^{m-2} k_m e^{i\varphi} \det \begin{pmatrix} -\lambda & 1 & & \\ k_2 & \ddots & \ddots & \\ & \ddots & \ddots & 1 \\ & & k_{m-2} & -\lambda \end{pmatrix} \right) \\ &= \det \begin{pmatrix} -\lambda & 1 & & \\ k_1 & \ddots & \ddots & \\ & \ddots & \ddots & 1 \\ & & k_{m-1} & -\lambda \end{pmatrix} - k_m \det \begin{pmatrix} -\lambda & 1 & & \\ k_2 & \ddots & \ddots & \\ & \ddots & \ddots & 1 \\ & & k_{m-2} & -\lambda \end{pmatrix} \\ &\quad + (-1)^{m-1} e^{i\varphi} \prod_{j=1}^m k_j + (-1)^{m-1} e^{-i\varphi}. \end{aligned}$$

That p_k is an even (odd) function if m is even (odd) follows easily by induction over m . \square

The most important part of Lemma 3 is that there are no mixed terms of λ and φ in equation (4). This leads to the fact (see Corollary 5 below) that the spectrum of every periodic operator A_{per}^k can be written as the preimage of the interval $[-2, 2]$ under some polynomial p_k . In this way the various parts of π_∞ are connected. We will make great use of this fact in Theorem 10. But first observe the following. The term in (4) involving φ can be simplified to $-2\cos(\varphi)$ or $2i\sin(\varphi)$ depending on the product $\prod_{j=1}^m k_j$. To avoid unnecessary paperwork, we give the following definition.

Definition 4. Let $m \in \mathbb{N}$ and $k \in \{\pm 1\}^m$. Then we call k even if $\prod_{j=1}^m k_j = 1$ and odd if $\prod_{j=1}^m k_j = -1$.

Note that we can always assume that a periodic operator A_{per}^k has an even period k . This is because we can always double the period, i.e. take $(k_1, \dots, k_m, k_1, \dots, k_m) \in \{\pm 1\}^{2m}$ instead of $(k_1, \dots, k_m) \in \{\pm 1\}^m$ as mentioned in the introduction.

Lemma 3 has the following important corollary.

Corollary 5. Let $m \in \mathbb{N}$ and $k \in \{\pm 1\}^m$. Then $\text{sp}(A_{per}^k) = p_k^{-1}([-2, 2])$ if k is even and $\text{sp}(A_{per}^k) = p_k^{-1}(i[-2, 2])$ if k is odd.

Proof. Let k be even first. By Proposition 2 and Lemma 3 we have

$$\begin{aligned} \text{sp}(A_{per}^k) &= \{\lambda \in \mathbb{C} : \det(a(\varphi) - \lambda I_m) = 0 \text{ for some } \varphi \in [0, 2\pi)\} \\ &= \{\lambda \in \mathbb{C} : p_k(\lambda) - 2\cos(\varphi) = 0 \text{ for some } \varphi \in [0, 2\pi)\} \\ &= \{\lambda \in \mathbb{C} : p_k(\lambda) \in [-2, 2]\} \\ &= p_k^{-1}([-2, 2]). \end{aligned}$$

If k is odd, just replace $-2\cos(\varphi)$ by $2i\sin(\varphi)$. \square

The next proposition not only shows that $\text{sp}(p_k(A_{per}^k)) = [-2, 2]$ ($i[-2, 2]$) for all even (odd) $k \in \{\pm 1\}^m$ but also that $p_k(A_{per}^k)$ has a very simple form.

Proposition 6. Let $m \in \mathbb{N}$, $k \in \{\pm 1\}^m$ and let p_k be the corresponding polynomial given by Lemma 3. If k is even, then

$$p_k(a^k(\varphi)) = 2\cos(\varphi)I_m$$

and $p_k(A_{per}^k)$ is the Laurent operator with 1 on its m -th sub- and superdiagonal (and 0 everywhere else):

$$p_k(A_{per}^k) = \begin{pmatrix} & & & & & \\ & & \ddots & & & \\ & & & 1 & & \\ \ddots & & & & 1 & \\ & 1 & & & & 1 \\ & & 1 & & & \ddots \\ & & & 1 & & \\ & & & & \ddots & \end{pmatrix}. \quad (5)$$

If k is odd, then

$$p_k(a^k(\varphi)) = -2i \sin(\varphi) I_m$$

and $p_k(A_{per}^k)$ is the Laurent operator with -1 on its m -th subdiagonal and 1 on its m -th superdiagonal (and 0 everywhere else):

$$p_k(A_{per}^k) = \begin{pmatrix} & & \ddots & & & \\ & & & 1 & & \\ \ddots & & & & 1 & \\ & -1 & & & & 1 \\ & & -1 & & & \ddots \\ & & & -1 & & \\ & & & & \ddots & \end{pmatrix}. \quad (6)$$

Proof. In both cases the first part follows immediately from Lemma 3 and the theorem of Cayley-Hamilton. For the second part observe that $2 \cos(\varphi) I_m = (e^{i\varphi} + e^{-i\varphi}) I_m$ is the symbol of (5) and $-2i \sin(\varphi) I_m = (-e^{i\varphi} + e^{-i\varphi}) I_m$ is the symbol of (6). The assertion thus follows by Proposition 2. \square

This simple observation now enables us to prove the first of our main results.

Theorem 7. Let $m \geq 2$, $k := (k_1, \dots, k_{m-2}, -1, 1) \in \{\pm 1\}^m$ and $\hat{k} := (k_1, \dots, k_{m-2}, 1, -1) \in \{\pm 1\}^m$. Furthermore, let $b \in \{\pm 1\}^{\mathbb{Z}}$ and

$$A^b := \begin{pmatrix} \ddots & \ddots & & & \\ \ddots & 0 & 1 & & \\ & b_0 & \boxed{0} & 1 & \\ & & b_1 & 0 & \ddots \\ & & & \ddots & \ddots \end{pmatrix},$$

where the box indicates $A_{0,0}^b$. If the corresponding polynomials p_k and $p_{\hat{k}}$ are equal, then there exist $c \in \{\pm 1\}^{\mathbb{Z}}$ and $B \in \mathcal{L}(\ell^2(\mathbb{Z}))$ such that

$$p_k(A^c) \cong B \oplus A^b,$$

where we consider the following decomposition of the Hilbert space $\ell^2(\mathbb{Z})$:

$$\ell^2(\mathbb{Z}) \cong \ell^2(\mathbb{Z} \setminus m\mathbb{Z}) \oplus \ell^2(m\mathbb{Z}) \cong \ell^2(\mathbb{Z}) \oplus \ell^2(\mathbb{Z}).$$

In particular, $\text{sp}(A^b) \subset \text{sp}(p_k(A^c))$.

Proof. Let $c \in \{\pm 1\}^{\mathbb{Z}}$ be the sequence defined by:

- $c_0 = 1$, $c_1 = -1$,
- $c_{rm+j} = k_{j-1}$ for $j \in \{2, \dots, m-1\}$, $r \in \mathbb{Z}$,

- $\prod_{j=0}^{m-1} c_{rm-j} = b_r$ for $r \in \mathbb{Z}$,

- $c_{rm+1} = -c_{rm}$ for $r \in \mathbb{Z}$.

Note that A^c is very similar to A_{per}^k and $A_{per}^{\hat{k}}$. The difference is that, depending on the sequence b , the entries c_{rm} and c_{rm+1} are swapped for some $r \in \mathbb{Z}$. For $m = 2$ this is exactly the same construction as in [3, Lemma 5]. First we will prove the following claim by induction:

Claim 1: $((A^c)^s)_{i,i-s+2j}$ only depends on the coefficients $c_{i-s+j+1}, \dots, c_{i+j}$ for $s \in \mathbb{N}$, $i \in \mathbb{Z}$ and $j \in \{0, \dots, s-1\}$. Furthermore, $((A^c)^s)_{i,i+s} = 1$ for all $s \in \mathbb{N}$, $i \in \mathbb{Z}$ and all other entries are 0.

For $s = 1$ we have $(A^c)_{i,i-1+2j} = c_{i+j}$ for $j = 0$, $i \in \mathbb{Z}$, and $(A^c)_{i,i+1} = 1$ for $i \in \mathbb{Z}$. All other entries are 0. So assume that the claim holds for $s-1$. Then

$$\begin{aligned} ((A^c)^s)_{i,i-s+2j} &= (A^c)_{i,i+1}((A^c)^{s-1})_{i+1,i-s+2j} + (A^c)_{i,i-1}((A^c)^{s-1})_{i-1,i-s+2j} \\ &= ((A^c)^{s-1})_{i+1,i-s+2j} + c_i((A^c)^{s-1})_{i-1,i-s+2j} \\ &= f(c_{i-s+j+2}, \dots, c_{i+j}) + c_i g(c_{i-s+j+1}, \dots, c_{i+j-1}), \end{aligned} \quad (7)$$

where f and g are some polynomials. Observe that c_i is contained in $\{c_{i-s+j+1}, \dots, c_{i+j}\}$ for all $i \in \mathbb{Z}$, $j \in \{0, \dots, s-1\}$. Thus $((A^c)^s)_{i,i-s+2j}$ only depends on the coefficients $c_{i-s+j+1}, \dots, c_{i+j}$ for all $i \in \mathbb{Z}$, $j \in \{0, \dots, s-1\}$. Plugging $j = s$ into (7) yields $((A^c)^s)_{i,i+s} = 1$ for all $i \in \mathbb{Z}$ by the same induction argument. Plugging in $j \in \mathbb{Z} + \frac{1}{2}$ and $j \in \mathbb{Z} \setminus \{0, \dots, s\}$ shows that all other entries are 0. This finishes the proof of the claim.

Using Claim 1 with $s = m$ and $i = rm$ for $r \in \mathbb{Z}$, we get that $((A^c)^m)_{rm,(r-1)m+2j}$ only depends on the coefficients $c_{(r-1)m+j+1}, \dots, c_{rm+j}$ for all $j \in \{0, \dots, m-1\}$, $((A^c)^m)_{rm,rm+m} = 1$ and all other entries in row rm are 0. Similarly, $((A^c)^m)_{(r-1)m+2j,rm}$ only depends on the coefficients $c_{(r-1)m+j+1}, \dots, c_{rm+j}$ for all $j \in \{0, \dots, m-1\}$, $((A^c)^m)_{rm+m,rm} = 1$ and all other entries in column rm are 0. Moreover, Claim 1 also implies that the same is true for $p_k(A^c)$ because p_k is an even/odd monic polynomial of degree m by Lemma 3.

Claim 2: $((A^c)^m)_{i,i-m} = \prod_{j=0}^{m-1} c_{i-j}$ for all $i \in \mathbb{Z}$.

This again follows easily by induction:

$$\begin{aligned} ((A^c)^m)_{i,i-m} &= (A^c)_{i,i-1}((A^c)^{m-1})_{i-1,i-m} + (A^c)_{i,i+1}((A^c)^{m-1})_{i+1,i-m} \\ &= c_i \cdot ((A^c)^{m-1})_{i-1,i-m} + 1 \cdot 0 \\ &= c_i \cdot \dots \cdot c_{i-(m-1)}. \end{aligned}$$

Since $((A^c)^s)_{i,i-m} = 0$ for all $s < m$, also $(p_k(A^c))_{i,i-m} = \prod_{j=0}^{m-1} c_{i-j}$ for all $i \in \mathbb{Z}$. By definition of c , it thus follows

$$(p_k(A^c))_{rm,(r-1)m} = \prod_{j=0}^{m-1} c_{rm-j} = b_r$$

for all $r \in \mathbb{Z}$. Furthermore, $(p_k(A^c))_{rm, (r-1)m+2j}$ only depends on the coefficients $c_{(r-1)m+j+1}, \dots, c_{rm+j}$ for $j \in \{1, \dots, m-1\}$. In particular, these numbers all depend on c_{rm} and c_{rm+1} but not on $c_{(r-1)m}, c_{(r-1)m+1}, c_{(r+1)m}$ or $c_{(r+1)m+1}$. This implies

$$(p_k(A^c))_{rm, (r-1)m+2j} = (p_k(A_{per}^{\hat{k}}))_{rm, (r-1)m+2j} = 0 \quad (\text{if } c_{rm} = -1, c_{rm+1} = 1)$$

or (using $p_k = p_{\hat{k}}$)

$$(p_k(A^c))_{rm, (r-1)m+2j} = (p_{\hat{k}}(A_{per}^{\hat{k}}))_{rm, (r-1)m+2j} = 0 \quad (\text{if } c_{rm} = 1, c_{rm+1} = -1)$$

for $j \in \{1, \dots, m-1\}$ by Proposition 6. In other words, the entries $(p_k(A^c))_{rm, (r-1)m+2j}$ ($j \in \{1, \dots, m-1\}$) can not “know” whether we swapped some of the entries c_{lm} and c_{lm+1} ($l \in \mathbb{Z}$) or not. Thus they have to remain zero. Similarly, the entries $(p_k(A^c))_{(r-1)m+2j, rm}$ ($j \in \{1, \dots, m-1\}$) remain 0. Therefore $p_k(A^c)$ looks like this (where $*$ means “some unimportant entries”):

$$p_k(A^c) = \begin{pmatrix} \vdots & \vdots & \vdots & \ddots & & & & & & & \\ \dots & * & \vdots & * & \dots & * & & & & & \\ \dots & \dots & 0 & \dots & \dots & 0 & 1 & & & & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & & & & \\ \dots & * & \vdots & * & \dots & * & 0 & * & & & \\ \ddots & \vdots & \vdots & \vdots & & \vdots & \vdots & \vdots & \ddots & & \\ & * & 0 & * & \dots & * & \vdots & * & \dots & * & \\ \vdots & b_r & 0 & \dots & \dots & 0 & \dots & \dots & 0 & 1 & \\ & & & * & \dots & * & \vdots & * & \dots & * & 0 & * \\ & & & & \ddots & \vdots & \vdots & \vdots & & \vdots & \vdots & \ddots \\ & & & & & * & 0 & * & \dots & * & \vdots & * & \dots \\ & & & & & & b_{r+1} & 0 & \dots & \dots & 0 & \dots & \dots \\ & & & \underbrace{\hspace{2cm}} & & & & * & \dots & * & \vdots & * & \dots \\ & & & & & & & \ddots & \vdots & \vdots & \vdots & \vdots & \vdots \end{pmatrix}.$$

Decomposing our Hilbert space $\ell^2(\mathbb{Z}) \cong \ell^2(\mathbb{Z} \setminus m\mathbb{Z}) \oplus \ell^2(m\mathbb{Z}) \cong \ell^2(\mathbb{Z}) \oplus \ell^2(\mathbb{Z})$, we get the following decomposition of $p_k(A^c)$:

$$p_k(A^c) \cong B \oplus A^b$$

for some $B \in \mathcal{L}(\ell^2(\mathbb{Z}))$. In particular, $\text{sp}(A^b) \subset \text{sp}(p_k(A^c))$. \square

By construction of the sequence c , we also have the following important corollary for periodic operators.

Corollary 8. *Under the same assumptions as in Theorem 7, we have that if $b \in \{\pm 1\}^{\mathbb{Z}}$ is an n -periodic sequence with even period (b_1, \dots, b_n) , then c , as defined in the proof of Theorem 7, is an nm -periodic sequence with even period (c_1, \dots, c_{nm}) . $B \in \mathcal{L}(\ell^2(\mathbb{Z}))$ then is a periodic operator, too. Furthermore, if we denote the symbols of A^b , A^c and B by a^b , a^c and a^B , then also $p_k(a^c)$ can be decomposed as $p_k(a^c) \cong a^B \oplus a^b$. In particular, $\text{sp}(a^b(\varphi)) \subset \text{sp}(p_k(a^c(\varphi)))$ for every $\varphi \in [0, 2\pi)$.*

Proof. The first part follows by construction of c . It remains to prove that the symbol $p_k(a^c)$ can be decomposed in a similar way. By Proposition 2, A^c is unitarily equivalent to the multiplication operator $M_{a^c} \in \mathcal{L}(L^2([0, 2\pi), \mathbb{C}^{nm}))$. Let us denote this equivalence by \mathcal{F}_{nm} , i.e. $A^c = \mathcal{F}_{nm}^* M_{a^c} \mathcal{F}_{nm}$. It follows

$$p_k(A^c) = p_k(\mathcal{F}_{nm}^* M_{a^c} \mathcal{F}_{nm}) = \mathcal{F}_{nm}^* p_k(M_{a^c}) \mathcal{F}_{nm} = \mathcal{F}_{nm}^* M_{p_k(a^c)} \mathcal{F}_{nm}.$$

Furthermore, A^b is unitarily equivalent to the multiplication operator $M_{a^b} \in \mathcal{L}(L^2([0, 2\pi), \mathbb{C}^m))$ and B is unitarily equivalent to the multiplication operator $M_{a^B} \in \mathcal{L}(L^2([0, 2\pi), \mathbb{C}^{(n-1)m}))$. Let us denote these equivalences by \mathcal{F}_m and $\mathcal{F}_{(n-1)m}$. Furthermore, let us denote the decomposition $\ell^2(\mathbb{Z}) \cong \ell^2(\mathbb{Z} \setminus m\mathbb{Z}) \oplus \ell^2(m\mathbb{Z})$ by U and the decomposition $L^2([0, 2\pi), \mathbb{C}^{nm}) \cong L^2([0, 2\pi), \mathbb{C}^{(n-1)m}) \oplus L^2([0, 2\pi), \mathbb{C}^m)$ (in the obvious way) by V . It is not hard to see that

$$(\mathcal{F}_{(n-1)m} \oplus \mathcal{F}_m) U \mathcal{F}_{nm}^* = V$$

holds. Thus

$$\begin{aligned} M_{p_k(a^c)} &= \mathcal{F}_{nm} p_k(A^c) \mathcal{F}_{nm}^* \\ &= \mathcal{F}_{nm} U^* (B \oplus A^b) U \mathcal{F}_{nm}^* \\ &= \mathcal{F}_{nm} U^* (\mathcal{F}_{(n-1)m}^* M_{a^B} \mathcal{F}_{(n-1)m} \oplus \mathcal{F}_m^* M_{a^b} \mathcal{F}_m) U \mathcal{F}_{nm}^* \\ &= V^* (M_{a^B} \oplus M_{a^b}) V. \end{aligned}$$

This implies $p_k(a^c) \cong a^B \oplus a^b$ (as functions of φ). \square

Definition 9. We define

$$S := \{p_k : p_k \text{ is a polynomial such that the assumptions of Theorem 7 are satisfied}\}$$

as our set of symmetries of π_∞ .

In the case of periodic operators we can prove the following stronger version of Theorem 7 that justifies the definition of S .

Theorem 10. Let $n \in \mathbb{N}$, let $b \in \{\pm 1\}^{\mathbb{Z}}$ be an n -periodic sequence and let $k \in \{\pm 1\}^m$ be such that $p := p_k \in S$. Moreover, let $c \in \{\pm 1\}^{\mathbb{Z}}$ be the sequence as constructed in the proof of Theorem 7. Then the following assertion holds:

$$p(\lambda) \in \text{sp}(A^b) \iff \lambda \in \text{sp}(A^c).$$

Proof. W.l.o.g. we can assume that (b_1, \dots, b_n) is even. By Corollary 8, (c_1, \dots, c_{nm}) is even, too. Let q and r be the polynomials given by Lemma 3 corresponding to b and c . Also denote by a^b and a^c the symbols of A^b and A^c . Fix some $\varphi \in [0, 2\pi)$ and let $\mu \in \mathbb{C}$ be an eigenvalue of $a^b(\varphi)$. Again by Corollary 8, there exists some $\lambda \in \text{sp}(a^c(\varphi))$ such that $p(\lambda) = \mu$. Since $q(\mu) = 2 \cos(\varphi)$ by Proposition 6, we have $q(p(\lambda)) = 2 \cos(\varphi)$. On the other hand, also $r(\lambda) = 2 \cos(\varphi)$ by Proposition 6. Thus

$$(q \circ p)(\lambda) = 2 \cos(\varphi) = r(\lambda)$$

(cf. Figure 4). Since both $q \circ p$ and r are polynomials and the above argument is valid for every $\varphi \in [0, 2\pi)$, we conclude that $q \circ p$ and r are equal.

It follows

$$\text{sp}(A^c) = r^{-1}([-2, 2]) = p^{-1}(q^{-1}([-2, 2])) = p^{-1}(\text{sp}(A^b))$$

by Corollary 5. \square




Figure 4: Schematic picture of the maps involved in the proof of Theorem 10.

Theorem 10 combined with the result $\mathbb{D} \subset \text{clos}(\pi_\infty)$ from [3] implies the following corollary.

Corollary 11. *Let $p \in S$. Then $p^{-1}(\pi_\infty) \subset \pi_\infty$ and hence $p^{-1}(\mathbb{D}) \subset p^{-1}(\text{clos}(\pi_\infty)) \subset \text{clos}(\pi_\infty)$.*

Recall that $k = (k_1, \dots, k_m) \in \{\pm 1\}^m$ generates a polynomial $p = p_k \in S$ if $k_{m-1} \neq k_m$, so that $\hat{k} = (k_1, \dots, k_{m-2}, k_m, k_{m-1}) \neq k$ but still $p_k = p_{\hat{k}}$. Thus it is immediate that all k of the form $(1, \dots, 1, -1, 1)$ and $(-1, \dots, -1, -1, 1)$ generate a polynomial $p_k \in S$. Indeed, if $k = (1, \dots, 1, -1, 1)$ and $\hat{k} = (1, \dots, 1, 1, -1)$, then A_{per}^k and $A_{per}^{\hat{k}}$ are unitarily equivalent by a simple shift. This implies that S contains a countable number of polynomials. However, there are also a lot more than these trivial examples as the following table shows. We conjecture that there are approximately $2^{\lceil \frac{m}{2} \rceil - 1}$ polynomials of degree m in S .

No.	k	$p_k(\lambda)$
2.1	$(-1, 1)$	λ^2
3.1	$(1, -1, 1)$	$\lambda^3 - \lambda$
3.2	$(-1, -1, 1)$	$\lambda^3 + \lambda$
4.1	$(1, 1, -1, 1)$	$\lambda^4 - 2\lambda^2$
4.2	$(-1, -1, -1, 1)$	$\lambda^4 + 2\lambda^2$
5.1	$(1, 1, 1, -1, 1)$	$\lambda^5 - 3\lambda^3 + \lambda$
5.2	$(1, -1, 1, -1, 1)$	$\lambda^5 - \lambda^3 + \lambda$
5.3	$(-1, 1, -1, -1, 1)$	$\lambda^5 + \lambda^3 + \lambda$
5.4	$(-1, -1, -1, -1, 1)$	$\lambda^5 + 3\lambda^3 + \lambda$
6.1	$(1, 1, 1, 1, -1, 1)$	$\lambda^6 - 4\lambda^4 + 3\lambda^2$
6.2	$(1, -1, -1, 1, -1, 1)$	$\lambda^6 - \lambda^2$
6.3	$(-1, -1, -1, -1, -1, 1)$	$\lambda^6 + 4\lambda^4 + 3\lambda^2$
7.1	$(1, 1, 1, 1, 1, -1, 1)$	$\lambda^7 - 5\lambda^5 + 6\lambda^3 - \lambda$
7.2	$(1, 1, -1, 1, 1, -1, 1)$	$\lambda^7 - 3\lambda^5 + 2\lambda^3 + \lambda$
7.3	$(1, -1, 1, -1, 1, -1, 1)$	$\lambda^7 - \lambda^5 + 2\lambda^3 - \lambda$
7.4	$(1, -1, -1, -1, 1, -1, 1)$	$\lambda^7 + \lambda^5 - 2\lambda^3 + \lambda$
7.5	$(-1, 1, 1, 1, -1, -1, 1)$	$\lambda^7 - \lambda^5 - 2\lambda^3 - \lambda$
7.6	$(-1, 1, -1, 1, -1, -1, 1)$	$\lambda^7 + \lambda^5 + 2\lambda^3 + \lambda$
7.7	$(-1, -1, 1, -1, -1, -1, 1)$	$\lambda^7 + 3\lambda^5 + 2\lambda^3 - \lambda$
7.8	$(-1, -1, -1, -1, -1, -1, 1)$	$\lambda^7 + 5\lambda^5 + 6\lambda^3 + \lambda$

Table 1: Short list of elements in S .

Remark 12. As mentioned in the introduction, we can iterate Corollary 11 to get


$$U := \bigcup_{n \in \mathbb{N}} p^{-n}(\mathbb{D}) \subset \bigcup_{n \in \mathbb{N}} p^{-n}(\text{clos}(\pi_\infty)) \subset \text{clos}(\pi_\infty)$$


for every $p \in S$. In other words, $z \in U$ if and only if $|p^n(z)| \leq 1$ for some $n \in \mathbb{N}$. Thus there is clearly a connection to the filled Julia set corresponding to p which is given by

$$J_f(p) := \{z \in \mathbb{C} : (p^n(z))_{n \in \mathbb{N}} \text{ is bounded}\}.$$


(see [10, Lemma 17.1]). Indeed, the boundary $J(p) := \partial J_f(p)$ (which is usually just called the Julia set corresponding to p) is contained in the closure of $\bigcup_{n \in \mathbb{N}} p^{-n}(z)$ for every $z \in \mathbb{C}$ except for at most one point (see [5, Corollary 14.8(a)]). Hence $J(p) \subset \text{clos}(U) \subset \text{clos}(\pi_\infty)$. Considering the pictures (viii) and (ix) below, it seems natural to conjecture that even the filled Julia set $J_f(p)$ is contained in $\text{clos}(U)$.

We conclude with some pictures of subsets of Σ . The red unit circle serves as a reference.







(v) $\bigcup_{\substack{p \in S \\ \deg(p) \leq 7}} p^{-1}(\mathbb{D})$




(vi) $\bigcup_{\substack{p \in S \\ \deg(p) = 12}} p^{-1}(\mathbb{D})$




(vii) $\bigcup_{\substack{p \in S \\ \deg(p) \leq 15}} p^{-1}(\mathbb{D})$



(viii) $p_k^{-9}(\mathbb{D})$ for $k = (1, -1, 1)$



(ix) $p_k^{-6}(\mathbb{D})$ for $k = (-1, 1, -1, -1, 1)$



(x) $\bigcup_{\substack{p \in S \\ \deg(p) \leq 8}} p^{-5}(\mathbb{D})$

References

- [1] S.N. Chandler-Wilde, R. Chonchaiya and M. Lindner: *Eigenvalue Problem meets Sierpinski Triangle: Computing the Spectrum of a Non-Self-Adjoint Random Operator*, Operators and Matrices, 5 (2011), 633-648.
- [2] S.N. Chandler-Wilde, R. Chonchaiya and M. Lindner: *On the Spectra and Pseudospectra of a Class of non-self-adjoint Random Matrices and Operators*, Operators and Matrices, 7 (2013), 739-775.
- [3] S.N. Chandler-Wilde and E.B. Davies: *Spectrum of a Feinberg-Zee Random Hopping Matrix*, Journal of Spectral Theory, 2 (2012), 147-179.
- [4] E.B. Davies: *Linear Operators and their Spectra*, Cambridge Studies in Advanced Mathematics 106, Cambridge University Press, Cambridge, 2007.
- [5] K. Falconer: *Fractal Geometry: Mathematical Foundations and Applications*, 2nd Ed., John Wiley, Chichester, 2003.
- [6] J. Feinberg and A. Zee: *Spectral Curves of Non-Hermitian Hamiltonians*, Nucl. Phys. B 552 (1999), 599-623.
- [7] R. Hagger: *On the Spectrum and Numerical Range of Tridiagonal Random Operators*, Preprint at arXiv: 1407.5486.
- [8] R. Hagger: *The Eigenvalues of Tridiagonal Sign Matrices are Dense in the Spectra of Periodic Tridiagonal Sign Operators*, Preprint at arXiv: 1412.1724
- [9] D.E. Holz, H. Orland and A. Zee: *On the Remarkable Spectrum of a Non-Hermitian Random Matrix Model*, J. Phys. A Math. Gen 36 (2003), 3385-3400.
- [10] J. Milnor: *Dynamics in one Complex Variable. Introductory Lectures*, Vieweg & Sohn, Braunschweig, 1999.

Author's address:

Raffael Hagger
Institute of Mathematics
Hamburg University of Technology
Schwarzenbergstr. 95 E
D-21073 Hamburg
GERMANY

`raffael.hagger@tuhh.de`