

Fredholmness of Toeplitz operators on the Fock space

Article

Accepted Version

Fulsche, R. and Hagger, R. (2019) Fredholmness of Toeplitz operators on the Fock space. *Complex Analysis and Operator Theory*, 13 (2). pp. 375-403. ISSN 1661-8262 doi: <https://doi.org/10.1007/s11785-018-0803-8> Available at <https://centaur.reading.ac.uk/84026/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1007/s11785-018-0803-8>

Publisher: Springer

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Fredholmness of Toeplitz operators on the Fock space

Robert Fulsche, Raffael Hagger

November 9, 2018

Abstract

The Fredholm property of Toeplitz operators on the p -Fock spaces F_α^p on \mathbb{C}^n is studied. A general Fredholm criterion for arbitrary operators from the Toeplitz algebra $\mathcal{T}_{p,\alpha}$ on F_α^p in terms of the invertibility of limit operators is derived. This paper is based on previous work, which establishes corresponding results on the unit balls \mathbb{B}^n [10].

AMS subject classification: Primary: 47B35; Secondary: 47L80, 47A53, 47A10

Keywords: Toeplitz operators, Fock spaces, essential spectrum, limit operators

1 Introduction

Consider the weighted Gaussian measure $d\mu_\nu(z) = (\nu/\pi)^n e^{-\nu|z|^2} dz$ on \mathbb{C}^n , where $\nu > 0$ and dz denotes the usual Lebesgue measure on $\mathbb{C}^n \cong \mathbb{R}^{2n}$. The Fock space F_α^p for $\alpha > 0$, $1 < p < \infty$ is the closed subspace of $L^p(\mathbb{C}^n, \mu_{p\alpha/2})$ consisting of entire functions. Toeplitz operators on these spaces are defined to be the composition of a multiplication operator M_f , where $f \in L^\infty(\mathbb{C}^n)$, and a certain projection P_α (see below) back onto the closed subspace F_α^p , i.e.

$$T_f = P_\alpha M_f.$$

The function f is then called the symbol of T_f or M_f , respectively.

When studying such Toeplitz operators a natural property to consider is the Fredholmness of such operators, that is: are the kernel and the cokernel of T_f finite dimensional? Inspired by Toeplitz operators on other spaces, e.g. on the Hardy space or on the Bergman space over the unit ball, one expects that the information about the Fredholmness of T_f can be extracted from the symbol, more precisely from the behaviour of f near infinity [5, 24]. For symbols which extend continuously to the boundary sphere of \mathbb{C}^n theorems of the following form are known: If f is nowhere zero on the boundary sphere, then T_f is Fredholm [6][7, Theorem 2.1]. Results for a more general class of symbols are known for the case $p = 2$: If f is of vanishing oscillation, then T_f is Fredholm if f is bounded away from zero close to the boundary (cf. [17, 25] for the Bergman space or [4, 22] for the Fock space). Corresponding results can be proven without the restriction $p = 2$, cf. Section 6 in this article. For Toeplitz operators with more

general symbols, or even other operators from the Toeplitz algebra $\mathcal{T}_{p,\alpha}$, which is just the norm closure of the algebra generated by Toeplitz operators, such results were missing.

Only recently the methods of limit operators, known from the theory of band-dominated operators on sequence spaces, were adapted to the case of Toeplitz operators on Bergman- and Fock spaces. It was realized that the notion of “boundary of \mathbb{C}^n (resp. \mathbb{B}^n in the Bergman space case)” in the usual sense was too restrictive. Instead, one densely embeds \mathbb{C}^n (resp. \mathbb{B}^n) into the maximal ideal space \mathcal{M} of $BUC(\mathbb{C}^n)$ (resp. $BUC(\mathbb{B}^n)$), the space of bounded uniformly continuous functions. We thus consider $\mathcal{M} \setminus \mathbb{C}^n$ as the boundary of \mathbb{C}^n and the boundary values of an operator $A \in \mathcal{T}_{p,\alpha}$ at $\mathcal{M} \setminus \mathbb{C}^n$ are obtained by “shifting” A to the boundary (we will make this precise below). For each $x \in \mathcal{M} \setminus \mathbb{C}^n$ we will get a boundary operator A_x , called a limit operator. In [16] and [23] a limit operator theory for the Bergman space over the unit ball was developed, whereas the corresponding results for the Fock space were derived in [3]. In both cases it was shown that operators in the Toeplitz algebra are compact if and only if all of their limit operators vanish. In [10] some ideas from the limit operator theory on sequence spaces were adapted to show that an operator in the Toeplitz algebra over the unit ball is Fredholm if and only if all of its limit operators are invertible. Let us briefly review a few of these ideas and where they originate. The main objective of study in limit operator theory is the class of band-dominated operators, which was first studied in its entirety by Simonenko [20, 21]. Specific classes of band-dominated operators were also considered earlier, e.g. by Gohberg and Krein [8]. In 1985 it was shown by Lange and Rabinovich [13] (see also [18, 19]) that a band-dominated operator is Fredholm if and only if all of its limit operators are invertible and their inverses are uniformly bounded. For the subsequent 25 years it was unclear whether the uniform boundedness condition is necessary or not. Various subclasses of operators have been studied by many different authors and in every case it was shown that the uniform boundedness condition is actually redundant, i.e. the inverses are automatically uniformly bounded if all limit operators are invertible. For the general case this was then shown by Lindner and Seidel [15] in 2014. The corresponding result for (essential) norms can be found in [11]. For a more detailed history of limit operators on sequence spaces we refer to [14] and [19].

In this paper, we further adapt these results to the Fock space to obtain the following main theorem:

Theorem 1. *An operator $A \in \mathcal{T}_{p,\alpha}$ is Fredholm if and only if all of its limit operators are invertible.*

Theorem 1 improves an earlier result by Bauer and Isralowitz ([3, Theorem 7.2]) for $p = 2$, where the inverses of the limit operators are additionally assumed to be uniformly bounded. Hence, besides generalizing to arbitrary p , we show that the uniform boundedness condition is redundant. We closely follow the lines of [10] in this paper. Since the Fock space is in some aspects simpler than the Bergman space, we can avoid some technical difficulties and can focus more on the actual ideas.

The paper is organized as follows: In Section 2 we introduce our notation and recall some basic results. In Section 3 we introduce band-dominated operators and provide some properties of them. Section 4 will be devoted to the theory of limit operators and the main theorem of this paper. In Section 5 methods

similar to those from Section 4 will be sketched to derive results on the essential norm of operators from the Toeplitz algebra. In the end, Section 6 will be used to show how the expected results on the Fredholmness of Toeplitz operators with symbols of vanishing oscillation can be derived from our main theorem.

2 Notation and basic definitions

In this section we present the main definitions and some basic results which are well-known and/or easy to prove.

For $\nu > 0$ let $d\mu_\nu$ denote the Gaussian measure

$$d\mu_\nu(z) = \left(\frac{\nu}{\pi}\right)^n e^{-\nu|z|^2} dz$$

on \mathbb{C}^n , where dz denotes the Lebesgue measure on $\mathbb{C}^n \simeq \mathbb{R}^{2n}$ and $|\cdot|$ denotes the norm coming from the standard hermitian inner product $\langle \cdot, \cdot \rangle$ on \mathbb{C}^n , which is linear in the first and antilinear in the second component. $d\mu_\nu$ is easily seen to be a probability measure. The space L_α^p is given by

$$L_\alpha^p = \{f : \mathbb{C}^n \rightarrow \mathbb{C}; f \text{ measurable and } \|f\|_{p,\alpha} < \infty\} = L^p(\mathbb{C}^n, d\mu_{p\alpha/2})$$

for $\alpha > 0$ and $1 < p < \infty$, where

$$\|f\|_{p,\alpha}^p := \int_{\mathbb{C}^n} |f(z)|^p d\mu_{p\alpha/2}(z).$$

Further, F_α^p denotes the closed subspace of entire functions in L_α^p . Throughout this paper we will assume, unless stated otherwise, $p \in (1, \infty)$ and $\alpha \in (0, \infty)$ without further mentioning it.

For a Banach space X we denote by $\mathcal{L}(X)$ the space of bounded linear operators on X and by $\mathcal{K}(X)$ the ideal of compact operators. By M_f we will denote the operator of multiplication by the function $f \in L^\infty(\mathbb{C}^n)$. We will use this symbol for both multiplication operators acting as $L_\alpha^p \rightarrow L_\alpha^p$ or $F_\alpha^p \rightarrow F_\alpha^p$ without mentioning p or α in the notation. A Toeplitz operator is an operator of the form $P_\alpha M_f : F_\alpha^p \rightarrow F_\alpha^p$, where f is called the symbol of the operator. Here P_α is the projection $L_\alpha^p \rightarrow F_\alpha^p$ onto the closed subspace given by the formula in Proposition 2 below. By $\mathcal{T}_{p,\alpha}$ we denote the norm-closed subalgebra of $\mathcal{L}(F_\alpha^p)$ generated by all Toeplitz operators with bounded symbols. A net of bounded linear operators $(A_\gamma)_\gamma$ on some Banach space X is said to converge $*$ -strongly to $A \in \mathcal{L}(X)$ if $A_\gamma \rightarrow A$ strongly and $A_\gamma^* \rightarrow A^*$ strongly, where B^* denotes the Banach space adjoint of $B \in \mathcal{L}(X)$. For a set $M \subseteq \mathbb{C}^n$ we denote its characteristic function by χ_M . By $B(z, r)$ we will denote the Euclidean ball around $z \in \mathbb{C}^n$ with radius $r > 0$.

We will need the following result regarding projections from L_α^p to F_α^p :

Proposition 2 ([12, Theorem 7.1]). *The linear operator $P_\alpha : L_\alpha^p \rightarrow F_\alpha^p$ given by*

$$(P_\alpha f)(z) = \int_{\mathbb{C}^n} e^{\alpha\langle z, w \rangle} f(w) d\mu_\alpha(w)$$

is a bounded projection onto F_α^p . In particular, $P_\alpha|_{F_\alpha^p} = \text{Id}$.

The following duality results will also be of importance:

Proposition 3. *Let $\frac{1}{p} + \frac{1}{q} = 1$. Then the following assertions hold under the usual dual pairing induced by the scalar product on L_α^2 :*

(i) $(L_\alpha^p)' \cong L_\alpha^q$,

(ii) for $g \in L_\alpha^q$ it holds

$$\|\langle \cdot, g \rangle_\alpha\|_{(L_\alpha^p)'} = \frac{2^n}{p^{n/p} q^{n/q}} \|g\|_{q,\alpha},$$

(iii) $(P_\alpha : L_\alpha^p \rightarrow L_\alpha^p)^* \cong (P_\alpha : L_\alpha^q \rightarrow L_\alpha^q)$.

Proof. The standard proof of $L^p \cong L^q$ yields (i) and (ii). Using the symmetry of P_α (see Proposition 2), one immediately gets (iii). We refer to [12, 27] for details. \square

Similarly, we have the following duality of Fock spaces:

Proposition 4. *Let $\frac{1}{p} + \frac{1}{q} = 1$. Then the following assertions hold under the usual dual pairing induced by the scalar product on F_α^2 :*

(i) $(F_\alpha^p)' \cong F_\alpha^q$,

(ii) for $g \in F_\alpha^q$ it holds

$$\|g\|_{q,\alpha} \leq \|\langle \cdot, g \rangle_\alpha\|_{(F_\alpha^p)'} \leq \frac{2^n}{p^{n/p} q^{n/q}} \|g\|_{q,\alpha}.$$

Proof. (i) is again standard (cf. [12, 27]), (ii) is [9, Theorem 1.2]. \square

Note that in both cases the isomorphism is not isometric for $p \neq 2$. However, these quasi-isometries still allow for the usual adjoint arguments, which we will use occasionally.

Proposition 5. *Let $(U_j)_{j \in \mathbb{N}}$ be a sequence of measurable subsets of \mathbb{C}^n such that every $z \in \mathbb{C}^n$ belongs to at most N of the sets U_j for some $N \in \mathbb{N}$. Further, let $(f_j)_{j \in \mathbb{N}}$ be a sequence of measurable functions $f_j : \mathbb{C}^n \rightarrow \mathbb{C}$ such that $\text{supp } f_j \subseteq U_j$ and $|f_j(z)| \leq 1$ for all $z \in \mathbb{C}^n$. Then, for every $g \in L_\alpha^p$*

$$\sum_{j=1}^{\infty} \int_{\mathbb{C}^n} |f_j(z)g(z)|^p d\mu_{p\alpha/2}(z) \leq N \|g\|_{p,\alpha}^p.$$

In particular,

$$\sum_{j=1}^{\infty} \|M_{f_j} g\|_{p,\alpha}^p \leq N \|g\|_{p,\alpha}^p.$$

Proof. As in [10, Proposition 5]. \square

The following two results are well-known and are provided here for completeness:

Lemma 6. *Let $\alpha, \beta, \gamma > 0$. The function*

$$(z, w) \mapsto e^{\alpha\langle z, w \rangle - \beta|z|^2 - \gamma|w|^2}$$

is bounded on $\mathbb{C}^n \times \mathbb{C}^n$ if and only if $4\beta\gamma - \alpha^2 \geq 0$.

Proof. For

$$|e^{\alpha\langle z, w \rangle - \beta|z|^2 - \gamma|w|^2}| = e^{\alpha \operatorname{Re}\langle z, w \rangle - \beta|z|^2 - \gamma|w|^2}$$

to be bounded it suffices to show that $\alpha \operatorname{Re}\langle z, w \rangle - \beta|z|^2 - \gamma|w|^2$ is bounded from above. Since

$$\alpha \operatorname{Re}\langle z, w \rangle - \beta|z|^2 - \gamma|w|^2 \leq \alpha|z||w| - \beta|z|^2 - \gamma|w|^2$$

and the right-hand side of this inequality is just the polynomial $p(x, y) = \alpha xy - \beta x^2 - \gamma y^2$ evaluated at $x = |z|, y = |w|$, the boundedness follows from the well-known fact that $p(x, y)$ is bounded from above for $4\beta\gamma - \alpha^2 \geq 0$.

Conversely, if $4\beta\gamma - \alpha^2 < 0$, set $w = \sqrt{\frac{\beta}{\gamma}}z$ to obtain

$$e^{\alpha\langle z, w \rangle - \beta|z|^2 - \gamma|w|^2} = e^{|z|^2(\alpha\sqrt{\frac{\beta}{\gamma}} - 2\beta)}.$$

Since $\alpha > 2\sqrt{\beta\gamma}$, this function is unbounded. \square

Proposition 7. *Let $D \subset \mathbb{C}^n$ be compact. Then the operators $P_\alpha M_{\chi_D}$ and $M_{\chi_D} P_\alpha : L_\alpha^p \rightarrow L_\alpha^p$ are compact operators.*

Proof. It is

$$\begin{aligned} P_\alpha M_{\chi_D}(f)(z) &= \left(\frac{\alpha}{\pi}\right)^n \int_{\mathbb{C}^n} e^{\alpha\langle z, w \rangle} \chi_D(w) e^{-\alpha|w|^2} f(w) dw \\ &= \left(\frac{2}{p}\right)^n \int_{\mathbb{C}^n} e^{\alpha\langle z, w \rangle - \frac{2\alpha - p\alpha}{2}|w|^2} \chi_D(w) f(w) d\mu_{p\alpha/2}(w). \end{aligned}$$

By the Hille-Tamarkin theorem [26, Theorem 41.6] it suffices to check (recall: $L_\alpha^p = L^p(\mathbb{C}^n, \mu_{p\alpha/2})$) that

$$\int_{\mathbb{C}^n} \left(\int_{\mathbb{C}^n} |e^{\alpha\langle z, w \rangle - \frac{2\alpha - p\alpha}{2}|w|^2}|^q \chi_D(w) d\mu_{p\alpha/2}(z) \right)^{\frac{p}{q}} d\mu_{p\alpha/2}(w) < \infty,$$

where $\frac{1}{p} + \frac{1}{q} = 1$. A direct computation shows

$$\begin{aligned}
& \int_{\mathbb{C}^n} \left(\int_{\mathbb{C}^n} |e^{\alpha\langle z, w \rangle - \frac{2\alpha - p\alpha}{2}|w|^2}|^q \chi_D(w) d\mu_{p\alpha/2}(z) \right)^{\frac{p}{q}} d\mu_{p\alpha/2}(w) \\
&= \left(\frac{p\alpha}{2\pi} \right)^{np} \int_D \left(\int_{\mathbb{C}^n} e^{q\alpha \operatorname{Re}\langle z, w \rangle - \frac{2q\alpha - pq\alpha}{2}|w|^2 - \frac{p\alpha}{2}|z|^2} dz \right)^{\frac{p}{q}} e^{-\frac{p\alpha}{2}|w|^2} dw \\
&= \left(\frac{p\alpha}{2\pi} \right)^{np} \int_D \left(\int_{\mathbb{C}^n} e^{q\alpha \operatorname{Re}\langle z, w \rangle - \frac{p\alpha}{4}|z|^2 - \frac{q^2\alpha}{p}|w|^2} e^{-\frac{p\alpha}{4}|z|^2} dz \right. \\
&\quad \left. \cdot e^{\frac{-2pq\alpha + p^2q\alpha + 2q^2\alpha}{2p}|w|^2} \right)^{\frac{p}{q}} e^{-\frac{p\alpha}{2}|w|^2} dw \\
&= \left(\frac{p\alpha}{2\pi} \right)^{np} \int_D \left(\int_{\mathbb{C}^n} e^{q\alpha \operatorname{Re}\langle z, w \rangle - \frac{p\alpha}{4}|z|^2 - \frac{q^2\alpha}{p}|w|^2} e^{-\frac{p\alpha}{4}|z|^2} dz \right)^{\frac{p}{q}} \\
&\quad \cdot e^{\frac{-3p\alpha + p^2\alpha + 2q\alpha}{2}|w|^2} dw \\
&\leq \left(\frac{p\alpha}{2\pi} \right)^{np} \int_D \left(\int_{\mathbb{C}^n} C e^{-\frac{p\alpha}{4}|z|^2} dz \right)^{\frac{p}{q}} e^{\frac{-3p\alpha + p^2\alpha + 2q\alpha}{2}|w|^2} dw \\
&< \infty,
\end{aligned}$$

where C is the bound from Lemma 6 with $\gamma = \frac{q^2\alpha}{p}$. The proof for $M_{\chi_D} P_\alpha$ is similar. \square

The next lemma is essentially a sloppy version of Jensen's inequality. Since we won't need the stronger form and the notation will be more convenient with this variant, we will just mention this weak estimate. It can easily be shown and will frequently be used.

Lemma 8. For $k \in \mathbb{N}$, $p \in (1, \infty)$ and $x_1, \dots, x_k \geq 0$ it is

$$\left(\sum_{j=1}^k x_j \right)^p \leq k^p \sum_{j=1}^k x_j^p.$$

3 Band-dominated operators

The aim of this section is to introduce band-dominated operators in $\mathcal{L}(L_\alpha^p)$ and to provide some basic properties of them. Most of the proofs are similar to those in [10] and use techniques adapted from the sequence space case (see [14, 19] and the references therein).

Definition 9. (i) An operator $A \in \mathcal{L}(L_\alpha^p)$ is called a band operator if there is a positive real number ω such that $M_f A M_g = 0$ for all $f, g \in L^\infty(\mathbb{C}^n)$ with $\operatorname{dist}(\operatorname{supp} f, \operatorname{supp} g) > \omega$. The infimum over all such ω will be denoted by $\omega(A)$ and is called the band width of A .

(ii) An operator $A \in \mathcal{L}(L_\alpha^p)$ is called a band-dominated operator if it is the norm limit of a sequence of band operators. The set of band-dominated operators on L_α^p will be denoted by $\operatorname{BDO}_\alpha^p$.




Figure 1: The function ϕ

Denote by $|z|_\infty$ the induced sup-norm from $\mathbb{R}^{2n} \cong \mathbb{C}^n$ and $\text{dist}_\infty(z, B) = \inf\{|z - w|_\infty; w \in B\}$ for $z \in \mathbb{C}^n$ and $B \subseteq \mathbb{C}^n$. Set

$$\zeta = \{[-3, 3]^{2n} + \sigma \subset \mathbb{R}^{2n}; \sigma \in 6\mathbb{Z}^{2n}\}$$

and enumerate ζ as $\zeta = \{B_j\}_{j=1}^\infty$ such that it is $0 \in B_1$. Furthermore, we denote

$$\Omega_k(B_j) := \{z \in \mathbb{C}^n; \text{dist}_\infty(z, B_j) \leq k\}$$

for $k = 1, 2, 3$ and $j \in \mathbb{N}$.

Proposition 10 ([3, Lemma 3.1]). *The B_j satisfy*

- (i) $B_j \cap B_k = \emptyset$ for $j \neq k$;
- (ii) every $z \in \mathbb{C}^n$ belongs to at most 2^{2n} of the sets $\Omega_1(B_j)$ and at most 4^{2n} of the sets $\Omega_3(B_j)$;
- (iii) $\text{diam}(B_j) = 6\sqrt{2n}$, where $\text{diam}(B_j)$ denotes the Euclidean diameter of B_j .

We will now construct a sequence of auxiliary functions, which will give a partition of unity of \mathbb{C}^n with particularly nice properties. Define the function $\phi : \mathbb{R} \rightarrow [0, 1]$ as in Figure 1. Further, define $\tilde{\varphi}_0 : \mathbb{C}^n \simeq \mathbb{R}^{2n} \rightarrow [0, 1]$ by

$$\tilde{\varphi}_0(x_1, \dots, x_{2n}) = \phi(x_1) \cdot \phi(x_2) \cdot \dots \cdot \phi(x_{2n}).$$

Now, for every $\sigma = (\sigma_1, \dots, \sigma_{2n}) \in 6\mathbb{Z}^{2n}$ set

$$\tilde{\varphi}_\sigma(x_1, \dots, x_{2n}) = \tilde{\varphi}_0(x_1 - \sigma_1, \dots, x_{2n} - \sigma_{2n}).$$

Let $(\sigma_j)_{j \in \mathbb{N}}$ be the enumeration of $6\mathbb{Z}^{2n}$ which coincides with the enumeration of ζ , i.e. $\sigma_j \in B_j$ for all $j \in \mathbb{N}$. We set $\varphi_j = \tilde{\varphi}_{\sigma_j}$. It is easily seen that the φ_j fulfill the following properties:

- (i) $\text{supp } \varphi_j = \Omega_1(B_j)$ for all j ,
- (ii) $\sum_{j=1}^\infty \varphi_j(z) = 1$ for all $z \in \mathbb{C}^n$,
- (iii) the sequence $(\varphi_j)_{j \in \mathbb{N}}$ is uniformly equicontinuous (every function is even Lipschitz continuous with Lipschitz constant at most $\frac{1}{2} \cdot 2n = n$).

In a similar way we can construct another sequence $(\psi_j)_{j \in \mathbb{N}}$ of functions, now such that the functions are non-negative and uniformly equicontinuous mappings from \mathbb{C}^n to $[0, 1]$ with

- (i) $\psi_j(z) = 1$ for all $z \in \Omega_2(B_j)$ for all $j \in \mathbb{N}$,
- (ii) $\text{supp } \psi_j = \Omega_3(B_j)$ for all $j \in \mathbb{N}$.

For each $t \in (0, 1)$ and each $j \in \mathbb{N}$ we define the functions $\varphi_{j,t}(z) := \varphi_j(tz)$ and $\psi_{j,t}(z) = \psi_j(tz)$. The following proposition gives a few characterisations of band-dominated operators:

Proposition 11. *Let $A \in \mathcal{L}(L_\alpha^p)$. The following are equivalent:*

- (i) *A is band-dominated;*
- (ii) $\lim_{t \rightarrow 0} \sup_{\|f\|=1} \sum_{j=1}^{\infty} \|M_{\varphi_{j,t}} A M_{1-\psi_{j,t}} f\|^p = 0$;
- (iii) $\lim_{t \rightarrow 0} \left\| \sum_{j=1}^{\infty} M_{\varphi_{j,t}} A M_{1-\psi_{j,t}} \right\| = 0$, where the convergence of the operator sum should be understood as strong convergence;
- (iv) $\lim_{t \rightarrow 0} \sup_{\|f\|=1} \sum_{j=1}^{\infty} \|[A, M_{\varphi_{j,t}}]f\|^p = 0$, where $[A, M_{\varphi_{j,t}}] = A M_{\varphi_{j,t}} - M_{\varphi_{j,t}} A$ is the commutator.

Proof. The strong convergence in (iii) follows from the fact that $\sum_{j=1}^{\infty} M_{\varphi_{j,t}} A$ converges strongly, which can easily be seen, and the following Lemma 12.
(i) \implies (ii): Let $\varepsilon > 0$ and B be a band operator such that $\|A - B\| < \varepsilon$. Further, let $t > 0$ be small enough such that

$$\text{dist}(\text{supp}(\varphi_{j,t}), \text{supp}(1 - \psi_{j,t})) > \omega(B),$$

where the distance on the left-hand side is by construction independent of j . Then, for all $j \in \mathbb{N}$

$$M_{\varphi_{j,t}} B M_{1-\psi_{j,t}} = 0.$$

We thus get for $f \in L_\alpha^p$

$$\begin{aligned} \sum_{j=1}^{\infty} \|M_{\varphi_{j,t}} A M_{1-\psi_{j,t}} f\|^p &= \sum_{j=1}^{\infty} \|M_{\varphi_{j,t}} (A - B) M_{1-\psi_{j,t}} f\|^p \\ &\leq 2^p \sum_{j=1}^{\infty} (\|M_{\varphi_{j,t}} (A - B) f\|^p \\ &\quad + \|M_{\varphi_{j,t}} (A - B) M_{\psi_{j,t}} f\|^p) \\ &\leq 2^p 2^{2n} (\|(A - B) f\|^p + \|(A - B) M_{\psi_{j,t}} f\|^p) \\ &\leq 2^{p+1} 2^{2n} \varepsilon^p \|f\|^p, \end{aligned}$$

where we used Proposition 5 and Lemma 8. Since $\varepsilon > 0$ was arbitrary, the result follows.

(ii) \implies (iii): Using Lemma 8 combined with Proposition 10 it is

$$\begin{aligned}
\left\| \sum_{j=1}^{\infty} M_{\varphi_{j,t}} AM_{1-\psi_{j,t}} f \right\|^p &= \int_{\mathbb{C}^n} \left| \sum_{j=1}^{\infty} (M_{\varphi_{j,t}} AM_{1-\psi_{j,t}} f)(z) \right|^p d\mu_{p\alpha/2} \\
&\leq \int_{\mathbb{C}^n} (2^{2n})^p \sum_{j=1}^{\infty} |(M_{\varphi_{j,t}} AM_{1-\psi_{j,t}} f)(z)|^p d\mu_{p\alpha/2} \\
&= (2^{2n})^p \sum_{j=1}^{\infty} \int_{\mathbb{C}^n} |(M_{\varphi_{j,t}} AM_{1-\psi_{j,t}} f)(z)|^p d\mu_{p\alpha/2} \\
&= (2^{2n})^p \sum_{j=1}^{\infty} \|M_{\varphi_{j,t}} AM_{1-\psi_{j,t}} f\|^p.
\end{aligned}$$

Taking the supremum over all f with $\|f\| = 1$ and then the limit $t \rightarrow 0$ gives the result.

(iii) \implies (i): The operator

$$A_m := \sum_{j=1}^{\infty} M_{\varphi_{j,\frac{1}{m}}} AM_{\psi_{j,\frac{1}{m}}}$$

can easily be seen to be a band operator. Since $\sum_j \varphi_{j,t} = 1$ for all $t > 0$ it is

$$\begin{aligned}
\|A - A_m\| &= \left\| \sum_{j=1}^{\infty} (M_{\varphi_{j,\frac{1}{m}}} A - M_{\varphi_{j,\frac{1}{m}}} AM_{\psi_{j,\frac{1}{m}}}) \right\| \\
&= \left\| \sum_{j=1}^{\infty} M_{\varphi_{j,\frac{1}{m}}} AM_{1-\psi_{j,\frac{1}{m}}} \right\| \\
&\rightarrow 0
\end{aligned}$$

for $m \rightarrow \infty$.

The equivalence (i) \iff (iv) is more technical and we refer to the identical proof in the unit ball case [10, Proposition 11]. \square

Lemma 12. *For every $j \in \mathbb{N}$ let $a_j, b_j : \mathbb{C}^n \rightarrow [0, 1]$ be measurable functions and assume that there is some $N \in \mathbb{N}$ such that each $z \in \mathbb{C}$ belongs to at most N of the sets $\text{supp}(a_j)$ and at most M of the sets $\text{supp}(b_j)$. If $A \in \mathcal{L}(L^p_\alpha)$, then the series*

$$\sum_{j=1}^{\infty} M_{a_j} AM_{b_j}$$

converges strongly and $\|\sum_{j=1}^{\infty} M_{a_j} AM_{b_j}\| \leq NM\|A\|$.

Proof. Observe that for each $f \in L^p_\alpha$ we have, as a consequence of Proposition 5, $\sum_{j=m}^{\infty} \|M_{b_j} f\|^p \rightarrow 0$ for $m \rightarrow \infty$. To prove the lemma it suffices to show that

$$\left\| \sum_{j=m}^{\infty} M_{a_j} AM_{b_j} f \right\|^p \rightarrow 0$$

for each $f \in L_\alpha^p$ as $m \rightarrow \infty$. So let $f \in L_\alpha^p$. Then

$$\begin{aligned} \left\| \sum_{j=m}^{\infty} M_{a_j} A M_{b_j} f \right\|^p &= \int_{\mathbb{C}^n} \left| \sum_{j=m}^{\infty} (M_{a_j} A M_{b_j} f)(z) \right|^p d\mu_{p\alpha/2}(z) \\ &\leq \int_{\mathbb{C}^n} \left(\sum_{j=m}^{\infty} |(M_{a_j} A M_{b_j} f)(z)| \right)^p d\mu_{p\alpha/2}(z). \end{aligned}$$

By assumption, the sum under the integral is pointwise a finite sum with at most N terms. Using Lemma 8 we can continue the estimate as follows:

$$\begin{aligned} &\leq \int_{\mathbb{C}^n} N^p \sum_{j=m}^{\infty} |(M_{a_j} A M_{b_j} f)(z)|^p d\mu_{p\alpha/2}(z) \\ &= N^p \sum_{j=m}^{\infty} \int_{\mathbb{C}^n} |(M_{a_j} A M_{b_j} f)(z)|^p d\mu_{p\alpha/2}(z) \\ &= N^p \sum_{j=m}^{\infty} \|M_{a_j} A M_{b_j} f\|^p. \end{aligned}$$

Using $|a_j(z)| \leq 1$, it follows:

$$\begin{aligned} &\leq N^p \|A\|^p \sum_{j=m}^{\infty} \|M_{b_j} f\|^p \\ &\rightarrow 0 \end{aligned}$$

as $m \rightarrow \infty$. The norm estimate follows easily as well. \square

Here are some of the properties of BDO_α^p :

Proposition 13. (i) $M_f \in \text{BDO}_\alpha^p$ for all $f \in L^\infty(\mathbb{C}^n)$.

(ii) BDO_α^p is a closed subalgebra of $\mathcal{L}(L_\alpha^p)$.

(iii) If $A \in \text{BDO}_\alpha^p$ is Fredholm and B is a Fredholm regularizer of A , then $B \in \text{BDO}_\alpha^p$. In particular, BDO_α^p is inverse closed.

(iv) $\mathcal{K}(L_\alpha^p)$ is a closed and two-sided ideal in BDO_α^p .

(v) $A \in \text{BDO}_\alpha^p$ if and only if $A^* \in \text{BDO}_\alpha^q$, where $\frac{1}{p} + \frac{1}{q} = 1$. In particular, BDO_α^2 is a C^* -algebra.

Proof. (i), (ii) and (v) are easy consequences of the definition of BDO_α^p . The proofs of (iii) and (iv) are quite technical. Since we will not need those statements for our purposes, we only refer to the identical proofs in the unit ball case in [10, Proposition 13]. \square

In the following we will show that Toeplitz operators are, in a sense made precise below, in BDO_α^p . The next lemma will be crucial for this.

Lemma 14. *There is a function $\beta_{p,\alpha} : [1, \infty) \rightarrow [0, \infty)$ with $\lim_{\sigma \rightarrow \infty} \beta_{p,\alpha}(\sigma) = 0$ which satisfies the following property: If $(a_j)_{j \in \mathbb{N}}, (b_j)_{j \in \mathbb{N}}$ are sequences of measurable functions from \mathbb{C}^n to $[0, 1]$ such that*

- *there exists $N \in \mathbb{N}$ such that each $z \in \mathbb{C}^n$ is contained in at most N of the sets $\text{supp}(a_j)$ and in at most N of the sets $\text{supp}(b_j)$,*
- *there exists $\sigma \geq 1$ such that $\text{dist}(\text{supp } a_j, \text{supp}(1 - b_j)) \geq \sigma$ for all $j \in \mathbb{N}$,*

then

$$\left\| \sum_{j=1}^{\infty} M_{a_j} P_{\alpha} M_{1-b_j} \right\| \leq N^2 \beta_{p,\alpha}(\sigma).$$

In particular,

$$\left\| \sum_{j=1}^{\infty} M_{a_j} P_{\alpha} M_{1-b_j} \right\| \rightarrow 0$$

for $\inf_{j \in \mathbb{N}} \text{dist}(\text{supp } a_j, \text{supp}(1 - b_j)) \rightarrow \infty$.

Proof. Observe that all operator series mentioned in the statement above and in the following proof converge $*$ -strongly as an easy consequence of Lemma 12. We borrow ideas from the proof of [3, Lemma 2.6] and sketch them here. It will be appropriate to start with the case $N = 1$. We first consider the limit case $p = \infty$. Define

$$L_{\alpha}^{\infty} := \{f : \mathbb{C}^n \rightarrow \mathbb{C}; f \text{ measurable and } \|f\|_{\infty, \alpha} < \infty\}$$

with

$$\|f\|_{\infty, \alpha} := \text{esssup}_{z \in \mathbb{C}^n} |f(z)| e^{-\frac{\alpha}{2}|z|^2}.$$

P_{α} , that is the integral operator with the same integral kernel as for the case $p < \infty$, is a projection from L_{α}^{∞} to F_{α}^{∞} , the closed subspace of holomorphic functions, and can hence be considered as an operator on L_{α}^{∞} (see [27, Corollary 2.22]). For $f \in L_{\alpha}^{\infty}$ and $z \in \mathbb{C}^n$, we get

$$\begin{aligned} & \left| \sum_{j=1}^{\infty} (M_{a_j} P_{\alpha} M_{1-b_j} f)(z) \right| e^{-\frac{\alpha}{2}|z|^2} \\ & \leq \left(\frac{\alpha}{\pi}\right)^n \sum_{j=1}^{\infty} |a_j(z)| \int_{\mathbb{C}^n} |1 - b_j(w)| |f(w)| e^{-\frac{\alpha}{2}|w|^2} e^{-\frac{\alpha}{2}|w-z|^2} dw \\ & \leq 2^n \|f\|_{\infty, \alpha} \end{aligned}$$

and hence it is $\|\sum_{j=1}^{\infty} M_{a_j} P_{\alpha} M_{1-b_j}\| \leq 2^n$ for $p = \infty$. Observe that, by the same argument with $a_1 \equiv 1 \equiv b_j$ and $a_j \equiv 0 \equiv b_1$ for all $j > 1$ we get that P_{α} is bounded on L_{α}^{∞} with norm $\leq 2^n$. If we can prove an estimate of the form $\|\sum_{j=1}^{\infty} M_{a_j} P_{\alpha} M_{1-b_j}\| \leq \beta_{p,\alpha}(\sigma)$ for $p = 2$, the result follows by interpolation for all $2 \leq p < \infty$ (see e.g. [12, Section 9] or [27, Chapter 2.4] for results on

interpolation of the spaces L_α^p). As in the proof of [3, Lemma 2.6], one can prove the estimate for $p = 2$ using the Schur test.

For $1 < p < 2$, instead of proving $\|\sum_{j=1}^{\infty} M_{a_j} P_\alpha M_{1-b_j}\| \leq \beta_{p,\alpha}(\sigma)$ directly, we will prove an estimate $\|\sum_{j=1}^{\infty} M_{1-b_j} P_\alpha M_{a_j}\| \leq \beta'_{q,\alpha}(\sigma)$ for the operator norm on $L_\alpha^q \cong (L_\alpha^p)'$ and then consider adjoints (see Proposition 3). As before, the estimate on L_α^q can be proven with the two limit steps $q = \infty$, $q = 2$ and then using interpolation. For the case $q = \infty$, observe that in the same way as above one can show

$$\left| \sum_{j=1}^{\infty} (M_{a_j} P_\alpha M_{b_j} f)(z) \right| e^{-\frac{\alpha}{2}|z|^2} \leq 2^n \|f\|_{\infty,\alpha}$$

(i.e. replace $1 - b_j$ by b_j) for $f \in L_\alpha^\infty$. Using

$$\begin{aligned} \left\| \sum_{j=1}^{\infty} P_\alpha M_{a_j} f \right\|_{\infty,\alpha} &\leq \|P_\alpha\| \left\| \sum_{j=1}^{\infty} M_{a_j} f \right\|_{\infty,\alpha} \\ &\leq \|P_\alpha\| \|f\|_{\infty,\alpha} \end{aligned}$$

one gets

$$\left\| \sum_{j=1}^{\infty} M_{1-b_j} P_\alpha M_{a_j} \right\| \leq 2^n + \|P_\alpha\|$$

for the case $q = \infty$. For $q = 2$, we already have the estimate since

$$\left(\sum_{j=1}^{\infty} M_{1-b_j} P_\alpha M_{a_j} \right)^* = \sum_{j=1}^{\infty} M_{a_j} P_\alpha M_{1-b_j}$$

and the two series converge strongly.

For the case $N > 1$ we set $\Lambda_1(z) = \{j \in \mathbb{N}; z \in \text{supp}(a_j)\}$ and $\Lambda_2(z) = \{j \in \mathbb{N}; z \in \text{supp}(b_j)\}$, both sets are considered to be ordered in the natural way. With

$$A_j^k = \{z \in \text{supp}(a_j); j \text{ is the } k\text{-th element of } \Lambda_1(z)\}$$

and

$$B_j^l = \{z \in \text{supp}(b_j); j \text{ is the } l\text{-th element of } \Lambda_2(z)\}$$

we have the disjoint unions $\text{supp}(a_j) = A_j^1 \cup \dots \cup A_j^N$ and $\text{supp}(b_j) = B_j^1 \cup \dots \cup B_j^N$ and both $A_{j_1}^k \cap A_{j_2}^k = \emptyset$ and $B_{j_1}^l \cap B_{j_2}^l = \emptyset$ hold for $j_1 \neq j_2$ and all $k, l = 1, \dots, N$. It follows

$$\sum_{j=1}^{\infty} M_{a_j} P_\alpha M_{1-b_j} = \sum_{k=1}^N \sum_{l=1}^N \sum_{j=1}^{\infty} M_{a_j \chi_{A_j^k}} P_\alpha M_{1-b_j \chi_{B_j^l}}$$

and we can write the operator as a finite sum of operators which fulfill the requirements of the lemma for $N = 1$. \square

For an operator $A \in \mathcal{L}(F_\alpha^p)$ we define its extension to L_α^p by $\hat{A} = AP_\alpha + Q_\alpha$, where $Q_\alpha = \text{Id} - P_\alpha$. Now we can prove the announced result about Toeplitz operators being band-dominated:

Theorem 15. *For any $A \in \mathcal{T}_{p,\alpha}$ it holds $\hat{A} \in \text{BDO}_\alpha^p$.*

Proof. We obtain that P_α is in BDO_α^p by combining Lemma 14 and Proposition 11 with the fact that

$$\liminf_{t \rightarrow 0} \inf_{j \in \mathbb{N}} \text{dist}(\text{supp}(\varphi_{j,t}), \text{supp}(1 - \psi_{j,t})) \geq \lim_{t \rightarrow 0} \frac{2}{t} = \infty.$$

By Proposition 13, the extension of every Toeplitz operator is in BDO_α^p and hence, since BDO_α^p is a norm-closed algebra, the result follows. \square

The last result in this section will be a criterion about Fredholmness for band-dominated operators.

Proposition 16. *Let $A \in \text{BDO}_\alpha^p$ be such that $[A, P_\alpha] = 0$. Assume that there is a positive constant M such that for every $t > 0$ there is an integer $j_0(t) > 0$ such that for all $j \geq j_0(t)$ there are operators $B_{j,t}, C_{j,t} \in \mathcal{L}(L_\alpha^p)$ with*

$$\|B_{j,t}\|, \|C_{j,t}\| \leq M$$

and

$$B_{j,t}AM_{\psi_{j,t}} = M_{\psi_{j,t}} = M_{\psi_{j,t}}AC_{j,t}.$$

Then $A|_{F_\alpha^p}$ is Fredholm and $\|(A|_{F_\alpha^p} + \mathcal{K}(F_\alpha^p))^{-1}\| \leq 2^{6n+1}\|P_\alpha\|M$.

Proof. The proof goes similarly to the unit ball case [10, Proposition 17]. We give a sketch of the proof here: For $t > 0$ define an operator

$$B_t := \sum_{j=j_0(t)}^{\infty} M_{\psi_{j,t}}B_{j,t}M_{\varphi_{j,t}},$$

where the series converges strongly $\|B_t\| \leq 2^{6n}M$ by Lemma 12. Using the identity

$$\begin{aligned} B_tA &= \sum_{j=j_0(t)}^{\infty} M_{\psi_{j,t}}B_{j,t}AM_{\varphi_{j,t}}M_{\psi_{j,t}} + \sum_{j=j_0(t)}^{\infty} M_{\psi_{j,t}}B_{j,t}[M_{\varphi_{j,t}}, A]M_{\psi_{j,t}} \\ &\quad + \sum_{j=j_0(t)}^{\infty} M_{\psi_{j,t}}B_{j,t}M_{\varphi_{j,t}}AM_{1-\psi_{j,t}} \end{aligned}$$

and some properties of band-dominated operators from Proposition 11, one can show

$$\lim_{t \rightarrow 0} \left\| B_tA - \sum_{j=j_0(t)}^{\infty} M_{\varphi_{j,t}} \right\| = 0$$

(see [10, Proposition 17]). With this fact we directly obtain

$$\lim_{t \rightarrow 0} \left\| P_\alpha B_t A|_{F_\alpha^p} - \sum_{j=j_0(t)}^{\infty} P_\alpha M_{\varphi_{j,t}}|_{F_\alpha^p} \right\| = 0.$$

Now since $\sum_{j=1}^{j_0(t)-1} P_\alpha M_{\varphi_{j,t}}|_{F_\alpha^p}$ is compact (Proposition 7) and

$$\sum_{j=1}^{\infty} P_\alpha M_{\varphi_{j,t}}|_{F_\alpha^p} = P_\alpha \sum_{j=1}^{\infty} M_{\varphi_{j,t}}|_{F_\alpha^p} = \text{Id},$$

$P_\alpha B_t A|_{F_\alpha^p} + \mathcal{K}(F_\alpha^p)$ converges to $\text{Id} + \mathcal{K}(F_\alpha^p)$ in the norm of the Calkin algebra $\mathcal{L}(F_\alpha^p)/\mathcal{K}(F_\alpha^p)$ as $t \rightarrow 0$. Using a standard Neumann series argument, we obtain the existence of $B \in \mathcal{L}(F_\alpha^p)$ such that $BA|_{F_\alpha^p} \in \text{Id} + \mathcal{K}(F_\alpha^p)$ and

$$\|B + \mathcal{K}(F_\alpha^p)\| \leq 2\|P_\alpha\| \|B_t\| \leq 2^{6n+1} \|P_\alpha\| M.$$

The other Fredholm regularizer (i.e. $A|_{F_\alpha^p} C \in \text{Id} + \mathcal{K}(F_\alpha^p)$ for some $C \in \mathcal{L}(F_\alpha^p)$) can be obtained similarly, defining operators C_t :

$$C_t := \sum_{j=j_0(t)}^{\infty} M_{\varphi_{j,t}} C_{j,t} M_{\psi_{j,t}}.$$

Using that $A^* \in \text{BDO}_\alpha^q$ one can analogously show

$$\lim_{t \rightarrow 0} \left\| AC_t - \sum_{j=j_0(t)}^{\infty} M_{\varphi_{j,t}} \right\| = \lim_{t \rightarrow 0} \left\| C_t^* A^* - \sum_{j=j_0(t)}^{\infty} M_{\varphi_{j,t}} \right\| = 0$$

and conclude again, using $[A, P_\alpha] = 0$, that

$$\lim_{t \rightarrow 0} \left\| AP_\alpha C_t|_{F_\alpha^p} - \sum_{j=j_0(t)}^{\infty} P_\alpha M_{\varphi_{j,t}}|_{F_\alpha^p} \right\| = 0.$$

Now proceed as in the first case. □

4 Limit operators

In this section we show our main result. As in Section 3, we use some techniques from limit operator theory on sequence spaces (see [14, 19]). The construction in Lemma 27 is due to Lindner and Seidel [15].

For each $z \in \mathbb{C}^n$ consider the weighted shift operators $C_z : L_\alpha^p \rightarrow L_\alpha^p$ given by

$$(C_z f)(w) = f(w - z) e^{\alpha \langle w, z \rangle - \frac{\alpha}{2} |z|^2}.$$

C_z is an isometry from L_α^p onto itself and from F_α^p onto itself for each $z \in \mathbb{C}^n$. Also, $C_z^{-1} = C_{-z}$. By \tilde{C}_z we will denote the restriction to F_α^p . It is easy to verify that the adjoint of $\tilde{C}_z : F_\alpha^p \rightarrow F_\alpha^p$ (in the sense of Proposition 4) is given by

$\tilde{C}_{-z} : F_\alpha^q \rightarrow F_\alpha^q$, where q is the dual exponent of p . For an operator $A \in \mathcal{L}(F_\alpha^p)$ and $z \in \mathbb{C}^n$ we define the shifted operator A_z by

$$A_z = \tilde{C}_z A \tilde{C}_{-z}.$$

Let \mathcal{M} denote the maximal ideal space of $\text{BUC}(\mathbb{C}^n)$, the unital C^* -algebra of bounded and uniformly continuous functions on \mathbb{C}^n , where \mathcal{M} is equipped with the weak- $*$ topology. We consider \mathbb{C}^n as a subset of \mathcal{M} by identifying each $z \in \mathbb{C}^n$ with the functional of point evaluation at z , $\delta_z : f \mapsto f(z)$. In this sense, \mathbb{C}^n is known to be a dense subspace of \mathcal{M} . If $A \in \mathcal{T}_{p,\alpha}$ and (z_γ) is a net in \mathbb{C}^n converging to $x \in \mathcal{M} \setminus \mathbb{C}^n$, then A_{z_γ} is known to converge $*$ -strongly to some limit operator, denoted by A_x , which does not depend on the particular choice of the net (z_γ) [3, Corollary 5.4].

In the following we will denote by $\tau_z : \mathbb{C}^n \rightarrow \mathbb{C}^n$ the function $w \mapsto w - z$ for each $z \in \mathbb{C}^n$. For later reference we collect a few results in the following lemma.

Lemma 17. (i) For $f \in L^\infty(\mathbb{C}^n)$ and $z \in \mathbb{C}^n$ it is

$$C_z M_f C_{-z} = M_{f \circ \tau_z}.$$

(ii) For $z \in \mathbb{C}^n$ it is

$$P_\alpha C_z = \tilde{C}_z P_\alpha.$$

(iii) For $f \in L^\infty(\mathbb{C}^n)$ and $z \in \mathbb{C}^n$ it is

$$(T_f)_z = T_{f \circ \tau_z}.$$

Proof. (iii) is a direct consequence of (i) and (ii). For (i), observe that for $g \in L_\alpha^p$ and $w \in \mathbb{C}^n$ it is

$$\begin{aligned} (C_z M_f C_{-z} g)(w) &= e^{\alpha\langle w, z \rangle - \frac{\alpha}{2}|z|^2} (M_f C_{-z} g)(w - z) \\ &= e^{\alpha\langle w, z \rangle - \frac{\alpha}{2}|z|^2} f(w - z) (C_{-z} g)(w - z) \\ &= f(w - z) g(w) \end{aligned}$$

and hence

$$C_z M_f C_{-z} = M_{f \circ \tau_z}$$

for every $z \in \mathbb{C}^n$. (ii) holds since

$$\begin{aligned} (P_\alpha C_z g)(w) &= \int_{\mathbb{C}^n} e^{\alpha\langle w, u \rangle} (C_z g)(u) d\mu_\alpha(u) \\ &= \left(\frac{\alpha}{\pi}\right)^n \int_{\mathbb{C}^n} e^{\alpha\langle w, u \rangle + \alpha\langle u, z \rangle - \frac{\alpha}{2}|z|^2} g(u - z) e^{-\alpha|u|^2} du \\ &= \left(\frac{\alpha}{\pi}\right)^n \int_{\mathbb{C}^n} e^{\alpha\langle w, v+z \rangle + \alpha\langle v+z, z \rangle - \frac{\alpha}{2}|z|^2} g(v) e^{-\alpha|v+z|^2} dv \\ &= e^{\alpha\langle w, z \rangle - \frac{\alpha}{2}|z|^2} \int_{\mathbb{C}^n} e^{\alpha\langle w-z, v \rangle} g(v) d\mu_\alpha(v) \\ &= (\tilde{C}_z P_\alpha g)(w) \end{aligned}$$

for all $w \in \mathbb{C}^n$ and $g \in L_\alpha^p$. \square

Proposition 18. *Let $A \in \mathcal{T}_{p,\alpha}$ and let (z_γ) be a net in \mathbb{C}^n converging to $x \in \mathcal{M} \setminus \mathbb{C}^n$ such that A_x is invertible. Let $f \in L^\infty(\mathbb{C}^n)$ be with compact support. Then there is a γ_0 such that for all $\gamma \geq \gamma_0$ there are operators $B_\gamma, D_\gamma \in \mathcal{L}(L_\alpha^p)$ satisfying*

$$\|B_\gamma\|, \|D_\gamma\| \leq 2(\|A_x^{-1}\| \|P_\alpha\| + \|Q_\alpha\|)$$

and

$$B_\gamma \hat{A} M_{f \circ \tau_{-z_\gamma}} = M_{f \circ \tau_{-z_\gamma}} = M_{f \circ \tau_{-z_\gamma}} \hat{A} D_\gamma.$$

Proof. The proof is similar to the proof of [10, Proposition 19]. Let $R > 0$ such that $\text{supp } f \subset B(0, R)$. $P_\alpha M_{\chi_{B(0,R)}}$ is compact by Proposition 7 and therefore it follows

$$\begin{aligned} & \left\| \left(C_{z_\gamma} (A P_\alpha + Q_\alpha) C_{-z_\gamma} - (A_x P_\alpha + Q_\alpha) \right) M_{\chi_{B(0,R)}} \right\| \\ &= \left\| \left(C_{z_\gamma} A P_\alpha C_{-z_\gamma} - A_x P_\alpha \right) M_{\chi_{B(0,R)}} \right\| \\ &= \left\| \left(\tilde{C}_{z_\gamma} A \tilde{C}_{-z_\gamma} - A_x \right) P_\alpha M_{\chi_{B(0,R)}} \right\| \\ &\rightarrow 0 \end{aligned}$$

for $z_\gamma \rightarrow x$, where we also used $C_{z_\gamma} Q_\alpha C_{-z_\gamma} = Q_\alpha$ (which is a consequence of Lemma 17 (ii)). Therefore there exists a γ_0 such that

$$\begin{aligned} R_\gamma &:= (A_x^{-1} P_\alpha + Q_\alpha) \left(C_{z_\gamma} (A P_\alpha + Q_\alpha) C_{-z_\gamma} - (A_x P_\alpha + Q_\alpha) \right) M_{\chi_{B(0,R)}} \\ &= (A_x^{-1} P_\alpha + Q_\alpha) C_{z_\gamma} (A P_\alpha + Q_\alpha) C_{-z_\gamma} M_{\chi_{B(0,R)}} - M_{\chi_{B(0,R)}} \end{aligned}$$

fulfills $\|R_\gamma\| < \frac{1}{2}$ for $\gamma \geq \gamma_0$. Here we used that $A_x^{-1} P_\alpha + Q_\alpha$ is the inverse of $A_x P_\alpha + Q_\alpha$. In particular, $\text{Id} + R_\gamma \in \mathcal{L}(L_\alpha^p)$ is invertible for all $\gamma \geq \gamma_0$. Multiplying R_γ by M_f yields

$$(A_x^{-1} P_\alpha + Q_\alpha) C_{z_\gamma} (A P_\alpha + Q_\alpha) C_{-z_\gamma} M_f = (\text{Id} + R_\gamma) M_f$$

and thus

$$M_f = (\text{Id} + R_\gamma)^{-1} (A_x^{-1} P_\alpha + Q_\alpha) C_{z_\gamma} (A P_\alpha + Q_\alpha) C_{-z_\gamma} M_f.$$

Multiplying by C_{-z_γ} from the left and C_{z_γ} from the right and using Lemma 17 (i) gives

$$C_{-z_\gamma} (\text{Id} + R_\gamma)^{-1} (A_x^{-1} P_\alpha + Q_\alpha) C_{z_\gamma} (A P_\alpha + Q_\alpha) M_{f \circ \tau_{-z_\gamma}} = M_{f \circ \tau_{-z_\gamma}},$$

and the claimed norm estimate follows easily with

$$B_\gamma := C_{-z_\gamma} (\text{Id} + R_\gamma)^{-1} (A_x^{-1} P_\alpha + Q_\alpha) C_{z_\gamma}.$$

The result for D_γ can be derived similarly: Since $M_{\chi_{B(0,R)}} P_\alpha$ is also compact,

$$\begin{aligned} & \left\| M_{\chi_{B(0,R)}} \left(C_{z_\gamma} (A P_\alpha + Q_\alpha) C_{-z_\gamma} - (A_x P_\alpha + Q_\alpha) \right) \right\| \\ &= \left\| M_{\chi_{B(0,R)}} P_\alpha (C_{z_\gamma} A C_{-z_\gamma} - A_x) P_\alpha \right\| \\ &\rightarrow 0 \end{aligned}$$

for $z_\gamma \rightarrow x$. Therefore

$$S_\gamma := M_{\chi_{B(0,R)}} \left(C_{z_\gamma} (AP_\alpha + Q_\alpha) C_{-z_\gamma} - (A_x P_\alpha + Q_\alpha) \right) (A_x^{-1} P_\alpha + Q_\alpha)$$

has norm $< \frac{1}{2}$ for large γ and we get

$$M_{f \circ \tau_{-z_\gamma}} (AP_\alpha + Q_\alpha) D_\gamma = M_{f \circ \tau_{-z_\gamma}}$$

with

$$D_\gamma := C_{-z_\gamma} (A_x^{-1} P_\alpha + Q_\alpha) (\text{Id} + S_\gamma)^{-1} C_{z_\gamma}. \quad \square$$

We get the following theorem:

Theorem 19. *If $A \in \mathcal{T}_{p,\alpha}$ is such that A_x is invertible for every $x \in \mathcal{M} \setminus \mathbb{C}^n$ and $\sup_{x \in \mathcal{M} \setminus \mathbb{C}^n} \|A_x^{-1}\| < \infty$, then A is Fredholm.*

Proof. The proof works entirely as in [10, Theorem 20]. For the readers convenience we reproduce it here.

Assume A is not Fredholm. One easily sees that $[\hat{A}, P_\alpha] = 0$. By Proposition 16, there exists a strictly increasing sequence $(j_m)_{m \in \mathbb{N}}$ and some $t > 0$ with

$$B \hat{A} M_{\psi_{j_m, t}} \neq M_{\psi_{j_m, t}}$$

or

$$M_{\psi_{j_m, t}} \hat{A} B \neq M_{\psi_{j_m, t}}$$

for all $m \in \mathbb{N}$ and all $B \in \mathcal{L}(L_\alpha^p)$ with $\|B\| \leq 2 \left(\sup_{x \in \mathcal{M} \setminus \mathbb{C}^n} \|A_x^{-1}\| \|P_\alpha\| + \|Q_\alpha\| \right)$.

Since both cases can be dealt with in the same way, we may assume

$$B \hat{A} M_{\psi_{j_m, t}} \neq M_{\psi_{j_m, t}}.$$

As $\text{diam}(\text{supp } \psi_{j, t}) \leq \frac{12\sqrt{2n}}{t} =: R$ for all $j \in \mathbb{N}$ by definition of $\psi_{j, t}$ and Proposition 10 (iii), there is a sequence $(w_{j_m})_{m \in \mathbb{N}}$ with $|w_{j_m}| \rightarrow \infty$ such that

$$\text{supp } \psi_{j_m, t} \subseteq B(w_{j_m}, R).$$

By the compactness of \mathcal{M} we may choose a convergent subnet (w_γ) of (w_{j_m}) such that $(-w_\gamma)$ converges to some $y \in \mathcal{M} \setminus \mathbb{C}^n$. By Proposition 18 there is a γ_0 such that for each $\gamma \geq \gamma_0$ there is an operator $B_\gamma \in \mathcal{L}(L_\alpha^p)$ with $\|B_\gamma\| \leq 2(\|A_y^{-1}\| \|P_\alpha\| + \|Q_\alpha\|)$ and

$$B_\gamma \hat{A} M_{\chi_{B(w_\gamma, R)}} = B_\gamma \hat{A} M_{\chi_{B(0, R)} \circ \tau_{w_\gamma}} = M_{\chi_{B(0, R)} \circ \tau_{w_\gamma}} = M_{\chi_{B(w_\gamma, R)}},$$

which is a contradiction. \square

We will need the following proposition:

Proposition 20. *Let $A \in \mathcal{T}_{p,\alpha}$ be compact and (z_γ) be a net in \mathbb{C}^n converging to $x \in \mathcal{M} \setminus \mathbb{C}^n$. Then A_{z_γ} converges $*$ -strongly to 0.*

Proof. This is the statement of [3, Theorem 1.1 and Lemma 6.1]. \square

The following theorem provides the converse of the previous theorem:

Theorem 21. *Let $A \in \mathcal{L}(F_\alpha^p)$ be Fredholm. Let (z_γ) be a net in \mathbb{C}^n converging to $x \in \mathcal{M} \setminus \mathbb{C}^n$ such that A_{z_γ} converges $*$ -strongly to $A_x \in \mathcal{L}(F_\alpha^p)$. Then A_x is invertible with $\|A_x^{-1}\| \leq \|(A + \mathcal{K}(F_\alpha^p))^{-1}\|$. If further B is a Fredholm regularizer of A , then B_{z_γ} converges $*$ -strongly to A_x^{-1} as $z_\gamma \rightarrow x$.*

Proof. As in [10, Theorem 21]. Since $AB - \text{Id}$ and $BA - \text{Id}$ are both compact, $(AB - \text{Id})_{z_\gamma}$ and $(BA - \text{Id})_{z_\gamma}$ converge $*$ -strongly to 0 for $z_\gamma \rightarrow x$. Further,

$$\begin{aligned} \|f\| &= \|\tilde{C}_{z_\gamma} \tilde{C}_{-z_\gamma} f\| \\ &\leq \|\tilde{C}_{z_\gamma} B A \tilde{C}_{-z_\gamma} f\| + \|\tilde{C}_{z_\gamma} (I - BA) \tilde{C}_{-z_\gamma} f\| \\ &\leq \|\tilde{C}_{z_\gamma} B \tilde{C}_{-z_\gamma}\| \|\tilde{C}_{z_\gamma} A \tilde{C}_{-z_\gamma} f\| + \|\tilde{C}_{z_\gamma} (I - BA) \tilde{C}_{-z_\gamma} f\| \\ &= \|B\| \|\tilde{C}_{z_\gamma} A \tilde{C}_{-z_\gamma} f\| + \|\tilde{C}_{z_\gamma} (I - BA) \tilde{C}_{-z_\gamma} f\| \end{aligned}$$

for all $f \in F_\alpha^p$. Letting $z_\gamma \rightarrow x$ we get $\|f\| \leq \|B\| \|A_x f\|$. A_x is hence injective with closed range. Using the same argument for the adjoint operators (see Proposition 4), we get $\|g\| \leq \|B\| \|A_x^* g\|$ for all $g \in F_\alpha^q$ and hence the surjectivity of A_x . A_x is therefore invertible. We also get $\|A_x^{-1}\| \leq \|B\|$ from these estimates. Since B was an arbitrary Fredholm regularizer of A , we have $\|A_x^{-1}\| \leq \|(A + \mathcal{K}(F_\alpha^p))^{-1}\|$. Using the identity

$$B_{z_\gamma} (A_x - A_{z_\gamma}) A_x^{-1} + (BA - \text{Id})_{z_\gamma} A_x^{-1} = B_{z_\gamma} - A_x^{-1},$$

which can easily be established, we also get the $*$ -strong convergence of B_{z_γ} to A_x^{-1} , since $A_{z_\gamma} \rightarrow A_x$, $(BA - I)_{z_\gamma} \rightarrow 0$ and $\|B_{z_\gamma}\| \leq \|B\|$. \square

Combining Theorem 21 and Theorem 19 with [3, Corollary 5.4], the fact that all limit operators exist for operators in $\mathcal{T}_{p,\alpha}$, we obtain:

Proposition 22. *$A \in \mathcal{T}_{p,\alpha}$ is Fredholm if and only if A_x is invertible for all $x \in \mathcal{M} \setminus \mathbb{C}^n$ and $\sup_{x \in \mathcal{M} \setminus \mathbb{C}^n} \|A_x^{-1}\| < \infty$.*

The condition $\sup \|A_x^{-1}\| < \infty$ is actually redundant. This will be shown in the remaining part of this section. Denote

$$r_t := \text{diam}(\text{supp } \varphi_{j,t}) = \frac{8\sqrt{2n}}{t},$$

which, of course, is independent of j , and define for $t > 0$, $F \subseteq \mathbb{C}^n$ and $A \in \mathcal{L}(L_\alpha^p)$

$$\nu(A|_F) := \inf\{\|Af\|; f \in L_\alpha^p, \|f\| = 1, \text{supp } f \subseteq F\}$$

and

$$\nu_t(A|_F) := \inf_{w \in \mathbb{C}^n} \nu(A|_{F \cap B(w, r_t)}).$$

We also use the notation $\nu(A) := \nu(A|_{\mathbb{C}^n})$.

Proposition 23. *For $A, B \in \mathcal{L}(L_\alpha^p)$ and $F \subseteq \mathbb{C}^n$ it is*

$$(i) \quad |\nu(A|_F) - \nu(B|_F)| \leq \|(A - B)M_{\chi_F}\| \leq \|A - B\|,$$

$$(ii) \quad |\nu_t(A|_F) - \nu_t(B|_F)| \leq \|A - B\|.$$

Proof. (i): As in [10, Proposition 27]: For the first statement let $\varepsilon > 0$ and pick $f \in L_\alpha^p$ with $\|f\| = 1$, $\text{supp } f \subseteq F$ and $\|Bf\| \leq \nu(B|_F) + \varepsilon$. Then

$$\begin{aligned} \nu(A|_F) - \nu(B|_F) - \varepsilon &\leq \nu(A|_F) - \|Bf\| \\ &\leq \|Af\| - \|Bf\| \\ &\leq \|(A - B)f\| \\ &\leq \|(A - B)M_{\chi_F}\|. \end{aligned}$$

Since the inequalities are symmetric in A and B , the result follows.

(ii): Let again $\varepsilon > 0$, pick $w \in \mathbb{C}^n$ such that

$$\nu(B|_{F \cap B(w, r_t)}) \leq \nu_t(B|_F) + \varepsilon.$$

Then

$$\begin{aligned} \nu_t(A|_F) - \nu_t(B|_F) - 2\varepsilon &\leq \nu_t(A|_F) - \nu(B|_{F \cap B(w, r_t)}) - \varepsilon \\ &\leq \nu(A|_{F \cap B(w, r_t)}) - \nu(B|_{F \cap B(w, r_t)}) - \varepsilon \\ &\leq \|(A - B)M_{\chi_{F \cap B(w, r_t)}}\| \\ &\leq \|A - B\|, \end{aligned}$$

where the second-to-last estimate can be concluded as in the first statement. Now use again the symmetry in A and B . \square

Proposition 24. *Let $A \in \mathcal{T}_{p, \alpha}$. For every $\varepsilon > 0$ there exists some $t > 0$ such that for all $F \subseteq \mathbb{C}^n$ and all $B \in \{\hat{A}\} \cup \{\hat{A}_x; x \in \mathcal{M} \setminus \mathbb{C}^n\}$:*

$$\nu(B|_F) \leq \nu_t(B|_F) \leq \nu(B|_F) + \varepsilon.$$

Proof. The first inequality follows by definition. For the second inequality: Let $(A_m)_{m \in \mathbb{N}}$ be a sequence of band operators that converges to \hat{A} in norm. Further, let $\varepsilon > 0$ and choose $m \in \mathbb{N}$ such that $\|\hat{A} - A_m\| < \frac{\varepsilon}{4}$. For $x \in \mathcal{M} \setminus \mathbb{C}^n$ let (z_γ) be a net in \mathbb{C}^n converging to x . $(A_m)_{z_\gamma}$ is a bounded net in $\mathcal{L}(L_\alpha^p)$, hence we may pass to a weakly convergent subnet, which we also denote by $(A_m)_{z_\gamma}$. Let the limit of this net be denoted by $(A_m)_x$. The strong convergence of \hat{A}_{z_γ} to \hat{A}_x implies that $C_{z_\gamma}(\hat{A} - A_m)C_{-z_\gamma}$ converges weakly to $\hat{A}_x - (A_m)_x$. Thus,

$$\|\hat{A}_x - (A_m)_x\| \leq \sup_\gamma \|C_{z_\gamma}(\hat{A} - A_m)C_{-z_\gamma}\| = \|\hat{A} - A_m\| < \frac{\varepsilon}{4}.$$

Now let $f, g \in L^\infty(\mathbb{C}^n)$ be such that $\text{dist}(\text{supp } f, \text{supp } g) > \omega(A_m)$. Then Lemma 17 gives

$$M_f(C_{z_\gamma}A_mC_{-z_\gamma})M_g = C_{z_\gamma}M_{f \circ \tau_{-z_\gamma}}A_mM_{g \circ \tau_{-z_\gamma}}C_{-z_\gamma} = 0,$$

since $\text{dist}(\text{supp } f \circ \tau_{-z_\gamma}, \text{supp } g \circ \tau_{-z_\gamma}) = \text{dist}(\text{supp } f, \text{supp } g)$. This implies $\omega((A_m)_{z_\gamma}) \leq \omega(A_m)$ and hence $\omega((A_m)_x) \leq \omega(A_m)$ by passing to the limit.

Observe now that, if we know that there exists a $t \in (0, 1)$ such that for all $F \subseteq \mathbb{C}^n$ and all $B \in \{A_m\} \cup \{(A_m)_x; x \in \mathcal{M} \setminus \mathbb{C}^n\}$ it is

$$\nu_t(B|_F) \leq \nu(B|_F) + \frac{\varepsilon}{2},$$

we are done, since by Proposition 23 it is

$$|\nu(\hat{A}|_F) - \nu(A_m|_F)| \leq \|\hat{A} - A_m\| < \frac{\varepsilon}{4}$$

and

$$|\nu(\hat{A}_x|_F) - \nu((A_m)_x|_F)| \leq \|\hat{A}_x - (A_m)_x\| < \frac{\varepsilon}{4}.$$

For the existence of such a t , we refer to the corresponding part of the proof of the unit ball case in [10, Proposition 23], which is identical to the situation in the Fock space. \square

Proposition 25. $\{A_x; x \in \mathcal{M}\}$ and $\{A_x; x \in \mathcal{M} \setminus \mathbb{C}^n\}$ are both compact in the strong operator topology for each $A \in \mathcal{T}_{p,\alpha}$.

Proof. \mathcal{M} and $\mathcal{M} \setminus \mathbb{C}^n$ are compact and $x \mapsto A_x$ is continuous w.r.t. the strong operator topology [3, Proposition 5.3]. \square

Lemma 26. Let $A \in \mathcal{T}_{p,\alpha}$, $w \in \mathbb{C}^n$ and $r > 0$. Then, for each $f \in L_\alpha^p$ with $\text{supp } f \subseteq B(w, r)$ and every $x \in \mathcal{M} \setminus \mathbb{C}^n$ there exists $g \in L_\alpha^p$ and $y \in \mathcal{M} \setminus \mathbb{C}^n$ with $\|g\| = \|f\|$, $\text{supp } g \subseteq B(0, r)$ and $\|\hat{A}_x f\| = \|\hat{A}_y g\|$. Further, $\nu(\hat{A}_y|_{B(0, r+|w|)}) \leq \nu(\hat{A}_x|_{B(0, r)})$.

Proof. Using the definition, one can quickly check that

$$\tilde{C}_{w_1} \tilde{C}_{w_2} = \tilde{C}_{w_1+w_2} e^{\frac{\alpha}{2}(\langle w_2, w_1 \rangle - \langle w_1, w_2 \rangle)}$$

for every $w_1, w_2 \in \mathbb{C}^n$. Let (z_γ) be a net in \mathbb{C}^n that converges to x . Taking a suitable subsequence if necessary, we may assume that

$$\tilde{C}_{-w} \tilde{C}_{z_\gamma} \tilde{A} \tilde{C}_{-z_\gamma} \tilde{C}_w = \tilde{C}_{z_\gamma-w} \tilde{A} \tilde{C}_{-(z_\gamma-w)} \rightarrow \tilde{A}_y$$

for some $y \in \mathcal{M} \setminus \mathbb{C}^n$ by Proposition 25, and hence $\tilde{C}_{-w} A_x \tilde{C}_w = \tilde{A}_y$. Since $P_\alpha C_w = \tilde{C}_w P_\alpha$, we also have $C_{-w} \hat{A}_x C_w = \hat{A}_y$. Now let $f \in L_\alpha^p$ be such that $\text{supp } f \subseteq B(w, r)$. Then $g := C_{-w} f$ satisfies $\|g\| = \|f\|$, $\text{supp } g \subseteq B(0, r)$ and $\|\hat{A}_y g\| = \|\hat{A}_x f\|$.

For the second statement, pick a function $h \in L_\alpha^p$ such that $\text{supp } h \subseteq B(0, r)$. Then $C_{-w} h$ satisfies $\text{supp}(C_{-w} h) \subseteq B(-w, r) \subseteq B(0, r + |w|)$ and $\|\hat{A}_x h\| = \|\hat{A}_y C_{-w} h\|$. \square

Lemma 27. Let $A \in \mathcal{T}_{p,\alpha}$. Then there exists a $y \in \mathcal{M} \setminus \mathbb{C}^n$ such that

$$\nu(\hat{A}_y) = \inf\{\nu(\hat{A}_x); x \in \mathcal{M} \setminus \mathbb{C}^n\}.$$

Proof. We only give a sketch of the proof here, since it is identical (up to the obvious changes) to the proof in [10, Lemma 25].

Using Proposition 24 we get a sequence $(t_k)_{k \in \mathbb{N}}$ with $r_{t_{k+1}} > 2r_{t_k}$ and

$$\nu_{t_k}(B|_F) \leq \nu(B|_F) + \frac{1}{2^{k+1}}$$

for all $k \in \mathbb{N}$, $F \subseteq \mathbb{C}^n$ and $B \in \{\hat{A}\} \cup \{\hat{A}_x; x \in \mathcal{M} \setminus \mathbb{C}^n\}$. Further, let $(x_j)_{j \in \mathbb{N}}$ be a sequence in $\mathcal{M} \setminus \mathbb{C}^n$ such that

$$\lim_{j \rightarrow \infty} \nu(\hat{A}_{x_j}) = \inf\{\nu(\hat{A}_x); x \in \mathcal{M} \setminus \mathbb{C}^n\}.$$

Using Lemma 26 repeatedly we can construct a sequence $(y_j)_{j \in \mathbb{N}} \subseteq \mathcal{M} \setminus \mathbb{C}^n$ such that

$$\nu(\hat{A}_{y_j}|_{B(0,4r_{t_k})}) \leq \nu(\hat{A}_{x_j}) + \frac{1}{2^{k-1}}.$$

Passing to a strongly convergent subnet $(A_{y_{j_\gamma}})_\gamma$ of $(A_{y_j})_{j \in \mathbb{N}}$ (Proposition 25), which converges to A_y for some $y \in \mathcal{M} \setminus \mathbb{C}^n$, we get

$$\|(\hat{A}_{y_{j_\gamma}} - \hat{A}_y)M_{\chi_{B(0,4r_{t_k})}}\| \rightarrow 0$$

by Proposition 7 and hence

$$\nu(\hat{A}_{y_{j_\gamma}}|_{B(0,4r_{t_k})}) \rightarrow \nu(\hat{A}_y|_{B(0,4r_{t_k})})$$

by Proposition 23. Then

$$\begin{aligned} \nu(\hat{A}_y) &\leq \nu(\hat{A}_y|_{B(0,4r_{t_k})}) = \lim_{\gamma} \nu(\hat{A}_{y_{j_\gamma}}|_{B(0,4r_{t_k})}) \\ &\leq \lim_{\gamma} \nu(\hat{A}_{x_{j_\gamma}}) + \frac{1}{2^{k-1}} = \lim_{j \rightarrow \infty} \nu(\hat{A}_{x_j}) + \frac{1}{2^{k-1}}. \end{aligned}$$

Taking the limit $k \rightarrow \infty$, we get the desired result. \square

We can now finally state and prove our main result. This extends a result of Bauer and Isralowitz ([3, Theorem 7.2]) to arbitrary p and shows that the uniform boundedness condition is actually redundant. Suárez et al. showed a similar result for the unit ball ([16, Theorem 5.8], [23, Theorem 10.3]), which was then improved in [10].

Theorem 28. *For $A \in \mathcal{T}_{p,\alpha}$ the following are equivalent:*

- (i) A is Fredholm,
- (ii) A_x is invertible and $\|A_x^{-1}\| \leq \|(A + \mathcal{K}(F_\alpha^p))^{-1}\|$ for all $x \in \mathcal{M} \setminus \mathbb{C}^n$,
- (iii) A_x is invertible for all $x \in \mathcal{M} \setminus \mathbb{C}^n$ and $\sup_{x \in \mathcal{M} \setminus \mathbb{C}^n} \|A_x^{-1}\| < \infty$,
- (iv) A_x is invertible for all $x \in \mathcal{M} \setminus \mathbb{C}^n$,
- (v) \hat{A}_x is invertible for all $x \in \mathcal{M} \setminus \mathbb{C}^n$.

Proof. The equivalence of (i), (ii) and (iii) was already proven in Proposition 22. That (iii) implies (iv) is clear. If A_x is invertible, $A_x^{-1}P_\alpha + Q_\alpha$ is an inverse of \hat{A}_x . Therefore (iv) implies (v). We finish the proof by showing that (v) implies (iii): Using that $\nu(B) = \|B^{-1}\|^{-1} > 0$ if B is invertible, we get

$$\sup_{x \in \mathcal{M} \setminus \mathbb{C}^n} \|\hat{A}_x^{-1}\| = \sup_{x \in \mathcal{M} \setminus \mathbb{C}^n} \frac{1}{\nu(\hat{A}_x)} = \frac{1}{\nu(\hat{A}_y)} < \infty,$$

where y is from Lemma 27. An inverse for each A_x is given by $\hat{A}_x^{-1}|_{F_\alpha^p}$. Since $\|B\| \leq \|\hat{B}\|$ for every $B \in \mathcal{L}(F_\alpha^p)$ we also get $\sup_{x \in \mathcal{M} \setminus \mathbb{C}^n} \|A_x^{-1}\| < \infty$. The fact that $\nu(B) = \|B^{-1}\|^{-1}$ for invertible B can be found in [14, Lemma 2.35]. \square

We get the following corollary directly from the definition of the essential spectrum, which is defined as

$$\sigma_{ess}(A) := \{\lambda \in \mathbb{C}; A - \lambda \text{ is not Fredholm}\}$$

for a bounded linear operator A .

Corollary 29. *For each $A \in \mathcal{T}_{p,\alpha}$ it is*

$$\sigma_{ess}(A) = \bigcup_{x \in \mathcal{M} \setminus \mathbb{C}^n} \sigma(A_x).$$

We emphasize that the redundancy of the uniform boundedness condition in Theorem 28 is essential for this corollary (cf. [3, Theorem 7.3], [16, Corollary 5.9], [23, Corollary 10.4]).

5 Norm estimates

The aim of this section is to provide estimates of the essential norm for operators $A \in \mathcal{T}_{p,\alpha}$. Here we adapt some ideas from [10] and [11] and (slightly) improve [3, Theorem 6.2, Theorem 7.1].

Define for $t > 0$, $F \subseteq \mathbb{C}^n$ and $A \in \mathcal{L}(F_\alpha^p)$

$$\|AP_\alpha|_F\| := \sup\{\|AP_\alpha f\|; f \in L_\alpha^p, \|f\| = 1, \text{supp } f \subseteq F\}$$

and

$$\| \|AP_\alpha|_F \|_t := \sup_{w \in \mathbb{C}^n} \|AP_\alpha|_{F \cap B(w, r_t)}\|.$$

Proposition 30. *For every $A \in \mathcal{T}_{p,\alpha}$ and every $\varepsilon > 0$ there exists a $t > 0$ such that for all $F \subseteq \mathbb{C}^n$ and every $B \in \{A\} \cup \{A_x; x \in \mathcal{M} \setminus \mathbb{C}^n\}$ it is*

$$\|BP_\alpha|_F\| \geq \| \|BP_\alpha|_F \|_t \geq \|BP_\alpha|_F\| - \varepsilon.$$

Proof. The proof is very similar to the proof of Proposition 24. Only the second inequality needs to be proven, as the first follows directly from the definition. If A_m is a band operator such that $\|AP_\alpha - A_m\| < \frac{\varepsilon}{4}$, one can prove that

$$\| \|C|_F \|_t \geq \|C|_F\| - \varepsilon$$

holds for all $F \subseteq \mathbb{C}^n$ and all $C \in \{A_m\} \cup \{(A_m)_x; x \in \mathcal{M} \setminus \mathbb{C}^n\}$, where $(A_m)_x$ is defined as in the proof of Proposition 24 (cf. [10, Proposition 27]). Then, using estimates similar to those in Proposition 23, one gets the desired result. \square

This now allows us to give an alternative proof of the second part of [3, Theorem 6.2]. Our approach additionally shows that the constants there may in fact be chosen as 1 and $\|P_\alpha\|^{-1}$. We do not know whether these constants are optimal, though.

Theorem 31. *Let $A \in \mathcal{T}_{p,\alpha}$. Then*

$$\frac{1}{\|P_\alpha\|} \|A + \mathcal{K}(F_\alpha^p)\| \leq \sup_{x \in \mathcal{M} \setminus \mathbb{C}^n} \|A_x\| \leq \|A + \mathcal{K}(F_\alpha^p)\|.$$

Proof. Let (z_γ) be a net converging to $x \in \mathcal{M} \setminus \mathbb{C}^n$. $(A + K)_{z_\gamma}$ converges $*$ -strongly to A_x for every $K \in \mathcal{K}(F_\alpha^p)$ since K_{z_γ} converges to 0 by Proposition 20. Using Banach-Steinhaus and the fact that \tilde{C}_w is an isometry for each $w \in \mathbb{C}^n$, one gets

$$\|A_x\| \leq \sup_\gamma \|\tilde{C}_{z_\gamma}(A + K)\tilde{C}_{-z_\gamma}\| = \|A + K\|.$$

Since this holds for all $x \in \mathcal{M} \setminus \mathbb{C}^n$ and all compact operators K , the second inequality follows. We give a sketch for the proof of the first inequality, and refer to [10, Theorem 28] for the missing details.

It can be seen that it suffices to prove

$$\inf_{K \in \mathcal{K}(L_\alpha^p, F_\alpha^p)} \|AP_\alpha + K\| \leq \sup_{x \in \mathcal{M} \setminus \mathbb{C}^n} \|A_x P_\alpha\|,$$

where $\mathcal{K}(L_\alpha^p, F_\alpha^p)$ is the set of compact operators from L_α^p to F_α^p . This will be proven by contradiction: Assume that

$$\inf_{K \in \mathcal{K}(L_\alpha^p, F_\alpha^p)} \|AP_\alpha + K\| > \sup_{x \in \mathcal{M} \setminus \mathbb{C}^n} \|A_x P_\alpha\| + \varepsilon$$

for some $\varepsilon > 0$. By Proposition 7,

$$\|AP_\alpha|_{\mathbb{C}^n \setminus B(0,s)}\| = \|AP_\alpha - AP_\alpha M_{\chi_{B(0,s)}}\| > \sup_{x \in \mathcal{M} \setminus \mathbb{C}^n} \|A_x P_\alpha\| + \varepsilon$$

for all $s > 0$. By Proposition 30 there is a $t \in (0, 1)$ with

$$\| \|AP_\alpha|_{\mathbb{C}^n \setminus B(0,s)} \| \|_t \geq \|AP_\alpha|_{\mathbb{C}^n \setminus B(0,s)}\| - \frac{\varepsilon}{2} > \sup_{x \in \mathcal{M} \setminus \mathbb{C}^n} \|A_x P_\alpha\| + \frac{\varepsilon}{2}.$$

Using the definition, for each $s > 0$ there must be some $w_s \in \mathbb{C}^n$ such that

$$\|AP_\alpha M_{\chi_{B(w_s, r_t)}}\| \geq \|AP_\alpha M_{\chi_{B(w_s, r_t)} \setminus B(0,s)}\| > \sup_{x \in \mathcal{M} \setminus \mathbb{C}^n} \|A_x P_\alpha\| + \frac{\varepsilon}{2}.$$

Using that $M_{\chi_{B(w_s, r_t)}} = C_{w_s} M_{\chi_{B(0, r_t)}} C_{-w_s}$, $P_\alpha C_{w_s} = \tilde{C}_{w_s} P_\alpha$ (Lemma 17) and the fact that \tilde{C}_{-w_s} and C_{w_s} are surjective isometries, we get

$$\|A_{-w_s} P_\alpha M_{\chi_{B(0, r_t)}}\| > \sup_{x \in \mathcal{M} \setminus \mathbb{C}^n} \|A_x P_\alpha\| + \frac{\varepsilon}{2}.$$

Since (w_s) clearly cannot converge in \mathbb{C}^n and \mathcal{M} is compact, there is a subnet of (w_s) , also denoted by (w_s) , such that $-w_s$ converges to $y \in \mathcal{M} \setminus \mathbb{C}^n$ and A_{-w_s} converges to A_y strongly, which implies by the compactness of $P_\alpha M_{\chi_{B(0, r_t)}}$ (Proposition 7)

$$\|A_{-w_s} P_\alpha M_{\chi_{B(0, r_t)}}\| \rightarrow \|A_y P_\alpha M_{\chi_{B(0, r_t)}}\|.$$

But this implies

$$\|A_y P_\alpha M_{\chi_{B(0, r_t)}}\| \geq \sup_{x \in \mathcal{M} \setminus \mathbb{C}^n} \|A_x P_\alpha\| + \frac{\varepsilon}{2},$$

which is a contradiction. \square

An improvement of this can be obtained if $p = 2$. Comparing to [3, Theorem 7.1] (or [16, Theorem 5.6], [23, Theorem 10.1] in case of the unit ball), this shows that the supremum is actually a maximum.

Theorem 32. For $A \in \mathcal{T}_{2,\alpha}$ it is

$$\|A + \mathcal{K}(F_\alpha^2)\| = \max_{x \in \mathcal{M} \setminus \mathbb{C}^n} \|A_x\|.$$

Proof. Replacing ν and ν_t by $\|\cdot\|$ and $\|\cdot\|_t$ in the proof of Lemma 27 and using Proposition 30, one can show that there is a $y \in \mathcal{M} \setminus \mathbb{C}^n$ such that $\|A_y P_\alpha\| = \sup\{\|A_x P_\alpha\|; x \in \mathcal{M} \setminus \mathbb{C}^n\}$. Since $\|A_x P_\alpha\| = \|A_x\|$ for $p = 2$, we get that the supremum in the theorem is actually a maximum. The equality follows by Theorem 31 with $\|P_\alpha\| = 1$. \square

6 Symbols of vanishing oscillation and vanishing mean oscillation

For a bounded and continuous function $f : \mathbb{C}^n \rightarrow \mathbb{C}$ we define

$$\text{Osc}_z^r(f) := \sup\{|f(z) - f(w)|; w \in \mathbb{C}^n, |z - w| \leq r\}$$

for $z \in \mathbb{C}^n$ and $r > 0$. A basic result about oscillations is that

$$\lim_{|z| \rightarrow \infty} \text{Osc}_z^r(f) = 0 \text{ for all } r > 0 \Leftrightarrow \lim_{|z| \rightarrow \infty} \text{Osc}_z^r(f) = 0 \text{ for one } r > 0.$$

Set

$$\text{VO}(\mathbb{C}^n) := \{f : \mathbb{C}^n \rightarrow \mathbb{C}; f \text{ bounded and continuous, } \lim_{|z| \rightarrow \infty} \text{Osc}_z^1(f) = 0\}.$$

It is easy to see that $\text{VO}(\mathbb{C}^n) \subset \text{BUC}(\mathbb{C}^n)$. In the case $p = 2$ it is well-known [4, 22] that for VO-symbols the Fredholm information is located at the boundary, i.e. for $f \in \text{VO}(\mathbb{C}^n)$ it holds

$$\sigma_{\text{ess}}(T_f) = f(\partial\mathbb{C}^n),$$

where $f(\partial\mathbb{C}^n)$ denotes the set of limit points of $f(z)$ as $|z| \rightarrow \infty$. We get those results for every $1 < p < \infty$ as a special case of Corollary 29:

Theorem 33. For $f \in \text{VO}(\mathbb{C}^n)$ it holds

$$\sigma_{\text{ess}}(T_f) = f(\partial\mathbb{C}^n).$$

Proof. By Corollary 29 we need to show that

$$\bigcup_{x \in \mathcal{M} \setminus \mathbb{C}^n} \sigma((T_f)_x) = f(\partial\mathbb{C}^n).$$

Let (z_γ) be a net in \mathbb{C}^n converging to $x \in \mathcal{M} \setminus \mathbb{C}^n$. Since $f \in \text{BUC}(\mathbb{C}^n)$ it is $f \circ \tau_{z_\gamma}(0) = f(z_\gamma) \rightarrow x(f)$. Further, observe that

$$|(f \circ \tau_{z_\gamma})(0) - (f \circ \tau_{z_\gamma})(w)| = |f(z_\gamma) - f(z_\gamma - w)| \leq \text{Osc}_{z_\gamma}^r(f) \rightarrow 0$$

for $|w| \leq r$. Therefore $f \circ \tau_{z_\gamma}$ converges uniformly on compact subsets to the constant function $x(f)$. Hence, using Lemma 17,

$$(T_f)_{z_\gamma} = T_{f \circ \tau_{z_\gamma}} \xrightarrow{s} T_{x(f)} = (T_f)_x$$

where $T_{x(f)}$ is just $x(f) \cdot \text{Id}$, thus $\sigma((T_f)_x) = \{x(f)\}$. But since $f(z_\gamma)$ converges to $x(f)$, this needs to be in $f(\partial\mathbb{C}^n)$ and hence

$$\sigma_{ess}(T_f) \subseteq f(\partial\mathbb{C}^n).$$

On the other hand, if $w \in f(\partial\mathbb{C}^n)$, let (z_m) be a sequence such that $f(z_m) \rightarrow w$. Using Proposition 25 we may choose a convergent subnet $((T_f)_{z_\gamma})_\gamma$ of $((T_f)_{z_m})_m$ and, as above, it converges to $w \text{Id}$ and we get the other implication. \square

We also get the following corollary for symbols with vanishing mean oscillation (see e.g. [2] for a definition):

Corollary 34. *If $f \in \text{VMO}(\mathbb{C}^n) \cap L^\infty(\mathbb{C}^n)$, then*

$$\sigma_{ess}(T_f) = \widetilde{f}(\partial\mathbb{C}^n),$$

where \widetilde{f} is the Berezin transform of f .

Proof. It is $\widetilde{f} \in \text{VO}(\mathbb{C}^n)$ [2, Corollary 2.8] and also $|\widetilde{f} - \widetilde{f}|^2 \in C_0(\mathbb{C}^n)$, i.e. $|\widetilde{f} - \widetilde{f}|^2$ vanishes at infinity [2, Theorem 5.3]. This of course implies $\widetilde{f} - \widetilde{f} \in C_0(\mathbb{C}^n)$. Therefore, $T_{\widetilde{f} - \widetilde{f}}$ is compact [3, Theorem 1.1] and hence

$$\sigma_{ess}(T_f) = \sigma_{ess}(T_{\widetilde{f}}) = \widetilde{f}(\partial\mathbb{C}^n). \quad \square$$

Remark. After submitting this paper, we noticed that Theorem 33 was independently found recently by Al-Qabani and Virtanen [1] using different methods.

References

- [1] Aamena Al-Qabani and Jani Virtanen, *Fredholm theory of Toeplitz operators on standard weighted Fock spaces*, Ann. Acad. Sci. Fenn.-M. **43** (2018).
- [2] Wolfram Bauer, *Mean Oscillation and Hankel Operators on the Segal-Bargmann Space*, Integr. Equ. Oper. Theory **52** (2005), 1–15.
- [3] Wolfram Bauer and Joshua Isralowitz, *Compactness characterization of operators in the Toeplitz algebra of the Fock space F_α^p* , J. Funct. Anal. **263** (2012), no. 5, 1323–1355.
- [4] Charles A. Berger and Lewis A. Coburn, *Toeplitz operators on the Segal-Bargmann space*, Trans. Amer. Math. Soc. **301** (1987), no. 2, 813–829.
- [5] Albrecht Böttcher and Bernd Silbermann, *Analysis of Toeplitz Operators*, 2nd ed., Springer Berlin Heidelberg, 2006.
- [6] Albrecht Böttcher and Hartmut Wolf, *Asymptotic invertibility of Bergman and Bargmann space Toeplitz operators*, Asymptot. Anal. **8** (1994), 15–33.

- [7] ———, *Spectral Approximation for Segal-Bargmann Space Toeplitz Operators*, Linear Operators, Banach Center Publ. **38** (1997), 25–48.
- [8] Israel Gohberg and Mark G. Krein, *Systems of integral equations on the semi-axis with kernels depending on the difference of arguments*, Usp. Mat. Nauk **13** (1958), no. 5, 3–72.
- [9] Willam E. Gryn and Todd Kemp, *Duality in Segal–Bargmann spaces*, J. Funct. Anal. **261** (2011), 1591–1623.
- [10] Raffael Hagger, *The Essential Spectrum of Toeplitz Operators on the Unit Ball*, Integr. Equ. Oper. Theory **89** (2017), no. 4, 519–556.
- [11] Raffael Hagger, Marko Lindner, and Markus Seidel, *Essential pseudospectra and essential norms of band-dominated operators*, J. Math. Anal. Appl. **437** (2016), no. 1, 255 – 291.
- [12] Svante Janson, Jaak Peetre, and Richard Rochberg, *Hankel Forms and the Fock Space*, Rev. Mat. Iberoamericana **3** (1987), no. 1, 61–138.
- [13] Bruno V. Lange and Vladimir S. Rabinovich, *Systems of integral equations on the semi-axis with kernels depending on the difference of arguments*, Usp. Mat. Nauk **13** (1958), no. 5, 3–72.
- [14] Marko Lindner, *Infinite Matrices and their Finite Sections*, Birkhäuser Verlag, Basel, Boston, Berlin, 2006.
- [15] Marko Lindner and Markus Seidel, *An affirmative answer to a core issue on limit operators*, J. Funct. Anal **267** (2014), no. 3, 901–917.
- [16] Mishko Mitkovski, Daniel Suárez, and Brett D. Wick, *The Essential Norm of Operators on $A_\alpha^p(\mathbb{B}_n)$* , Integr. Equ. Oper. Theory **75** (2013), 197–233.
- [17] Antti I. Perälä and Jani A. Virtanen, *A note on the Fredholm properties of Toeplitz operators on weighted Bergman spaces with matrix-valued symbols*, Oper. Matrices **5** (2011), no. 1, 97–106.
- [18] Vladimir Rabinovich, Steffen Roch, and Bernd Silbermann, *Fredholm theory and finite section method for band-dominated operators*, Integr. Equ. Oper. Theory **30** (1998), no. 4, 452 – 495.
- [19] ———, *Limit Operators and Their Applications in Operator Theory*, Birkhäuser Verlag, Basel, Boston, Berlin, 2004.
- [20] Igor B. Simonenko, *Operators of convolution type in cones*, Mat. Sb. **74** (1967), no. 116, 298–313.
- [21] ———, *On multidimensional discrete convolutions*, Mat. Issled. **3** (1968), no. 1, 108–127.
- [22] Karel Stroethoff, *Hankel and Toeplitz operators on the Fock space*, Michigan Math. J. **39** (1992), 3–16.
- [23] Daniel Suárez, *The Essential Norm of Operators in the Toeplitz Algebra on $A^p(\mathbb{B}_n)$* , Indiana Univ. Math. J. **56** (2007), no. 5, 2185–2232.

- [24] Harald Upmeyer, *Toeplitz Operators and Index Theory in Several Complex Variables*, Birkhäuser Basel, 1996.
- [25] Jingbo Xia and Dechao Zheng, *Toeplitz operators and Toeplitz algebra with symbols of vanishing oscillation*, J. Operator Theory **76** (2016), no. 1, 107 – 131.
- [26] Adriaan Zaanen, *Introduction to Operator Theory in Riesz Spaces*, Springer-Verlag, Heidelberg, 1997.
- [27] Kehe Zhu, *Analysis on Fock Spaces*, Graduate Texts in Mathematics, vol. 263, Springer US, New York, 2012.

Robert Fulsche

[✉fulsche@math.uni-hannover.de](mailto:fulsche@math.uni-hannover.de)

Raffael Hagger

[✉raffael.hagger@math.uni-hannover.de](mailto:raffael.hagger@math.uni-hannover.de)

Both authors:

Institut für Analysis

Leibniz Universität Hannover

Welfengarten 1

30167 Hannover

GERMANY