

Does increased atmospheric resolution improve seasonal climate predictions?

Article

Published Version

Creative Commons: Attribution 4.0 (CC-BY)

Open Access

Scaife, A. A., Camp, J., Comer, R., Davis, P., Dunstone, N., Gordon, M., MacLachlan, C., Martin, N., Nie, Y., Ren, H.-L., Roberts, M., Robinson, W., Smith, D. and Vidale, P. L. ORCID: <https://orcid.org/0000-0002-1800-8460> (2019) Does increased atmospheric resolution improve seasonal climate predictions? *Atmospheric Science Letters*, 20 (8). e922. ISSN 1530-261X doi: <https://doi.org/10.1002/asl.922> Available at <https://centaur.reading.ac.uk/84866/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1002/asl.922>

Publisher: Wiley

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur




CentAUR

Central Archive at the University of Reading

Reading's research outputs online

RESEARCH ARTICLE

Does increased atmospheric resolution improve seasonal climate predictions?

Adam A. Scaife^{1,2}  | Joanne Camp¹  | Ruth Comer¹ | Philip Davis¹ |
Nick Dunstone¹  | Margaret Gordon¹ | Craig MacLachlan¹ | Nicola Martin¹ |
Yu Nie³ | Hong-Li Ren^{3,4} | Malcolm Roberts¹ | Walter Robinson⁵ | Doug Smith¹ |
Pier Luigi Vidale⁶

¹Met Office Hadley Centre, Exeter, UK

²College of Engineering, Mathematics and Physical Sciences, University of Exeter, Exeter, UK

³Laboratory for Climate Studies & CMA-NJU Joint Laboratory for Climate Prediction Studies, National Climate Center, China Meteorological Administration, Beijing, China

⁴Department of Atmospheric Science, School of Environmental Studies, China University of Geoscience, Wuhan, China

⁵Department of Marine, Earth, and Atmospheric Sciences, North Carolina State University, Raleigh, North Carolina

⁶NCAS-Climate & Department of Meteorology, University of Reading, Reading, UK

Correspondence

Adam A. Scaife, Met Office Hadley Centre, Fitz Roy Road, Exeter EX1 3PB, UK.
Email: adam.scaife@metoffice.gov.uk

Funding information

Natural Environment Research Council, Grant/Award Number: NE/S004645/1; H2020 PRIMAVERA project, Grant/Award Number: GA641727; H2020 EUCP project, Grant/Award Number: GA 776613; UK Public Weather Service Program; Newton Fund; Met Office Climate Science for Service Partnership (CSSP) China; UK-China Research & Innovation Partnership Fund

Abstract

We assess the impact of atmospheric horizontal resolution on the prediction skill and fidelity of seasonal forecasts. We show the response to an increase of atmospheric resolution from 0.8 to 0.3° horizontal grid spacing in parallel ensembles of forecasts. Changes in the prediction skill of major modes of tropical El Niño Southern Oscillation (ENSO) and extratropical North Atlantic Oscillation (NAO) variability are small and not detected and there is no discernible impact on the weak signal-to-noise ratio in seasonal predictions of the winter NAO at this range of resolutions. Although studies have shown improvements in the simulation of tropical cyclones as model resolution is increased, we find little impact on seasonal prediction skill of either their numbers or intensity. Over this range of resolutions it appears that the benefit of increasing atmospheric resolution to seasonal climate predictions is minimal. However, at yet finer scales there appears to be increased eddy feedback which could strengthen weak signals in predictions of the NAO. Until prediction systems can be run operationally at these scales, it may be better to use additional computing resources for other enhancements such as increased ensemble size, for which there is a clear benefit in extratropical seasonal prediction skill.

KEY WORDS

atmospheric resolution, eddy feedback, seasonal prediction, signal to noise ratio

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2019 The Authors. *Atmospheric Science Letters* published by John Wiley & Sons Ltd on behalf of the Royal Meteorological Society.

1 | INTRODUCTION

Studies with global climate models show benefits from increasing atmospheric resolution (e.g., Roberts *et al.*, 2018), with modest increases in blocking frequency (Berckmans *et al.*, 2013; Schiemann *et al.*, 2017), improvements in regional climatology (e.g., Yao *et al.*, 2017) and improvements in rainfall variability (e.g., Kopparla *et al.*, 2013). However, the benefits of increased atmospheric resolution for seasonal prediction *skill* are less clear. Operational seasonal prediction systems are now being at resolutions well beyond one degree (e.g., MacLachlan *et al.*, 2014; Johnson *et al.*, 2018). The effects on prediction skill from further increases in resolution in the atmosphere (e.g., Jung *et al.*, 2012; Jia *et al.*, 2015; Zhu *et al.*, 2015; Prodhomme *et al.*, 2016) or ocean (e.g., Kirtman *et al.*, 2017) is an active research topic and it is not known whether this outweighs the benefits of larger ensemble size (e.g., Scaife *et al.*, 2014; Doi *et al.*, 2019), increased vertical resolution (e.g., Marshall and Scaife, 2010; Butler *et al.*, 2016) or improved initial conditions (e.g., Kumar *et al.*, 2015; Nie *et al.*, 2019).

Here we compare results from two parallel ensembles of seasonal hindcasts which are identical in all respects except atmospheric resolution, which is more than doubled in our tests. We compare the prediction skill of the main modes of tropical and extratropical variability and also tropical cyclones, all of which are known to exhibit seasonal predictability. We also examine eddy feedback on the Atlantic jet stream at different resolutions as this is one of the likely mechanisms by which increased resolution could improve simulated climate.

2 | STANDARD AND HIGHER-RESOLUTION PREDICTION SYSTEMS

Our hindcasts are taken from the GloSea5 prediction system which is used to make real-time seasonal forecasts at the UK Met Office (MacLachlan *et al.*, 2014). We examine hindcasts for the boreal summers from 1993 to 2015 and the neighbouring boreal winters from 1994 to 2016. Lagged ensembles are formed from three start dates centred on May 1 for JJA predictions and November 1 for DJF predictions. For DJF we use hindcast start dates of October 25, November 1 and November 9, while for summer we use April 25, May 1 and May 9. Further ensemble member perturbations are generated with a stochastic physics scheme as described in MacLachlan *et al.* (2014). We use a total of 21 ensemble members for each season. The system and model components are as described in MacLachlan *et al.* (2014) with initialisation of the ocean, atmosphere and sea ice. Ocean

resolution is 0.25° latitude and longitude throughout the globe with 75 quasi-horizontal ocean levels. The atmospheric component contains 85 quasi-horizontal levels and is run at two horizontal resolutions for this study: standard horizontal resolution is 0.83° while higher horizontal resolution is 0.35° longitude.

3 | PREDICTION SKILL OF ENSO

Seasonal predictability of El Nino Southern Oscillation (ENSO) has been well established for many years (Cane *et al.*, 1986) and the high predictability of ENSO is now a cornerstone of current seasonal prediction capability (Smith *et al.*, 2012). Nevertheless, outstanding questions remain (Tang *et al.*, 2018), for example regarding the predictability of different ENSO types (Imada *et al.*, 2015; Ren *et al.*, 2019), or increases in skill as models improve (e.g., Luo *et al.*, 2008). Although comprehensive models now appear to have the edge over simpler prediction models, there is also variation in prediction skill over time (Barnston *et al.*, 2012). This variation may simply be due to low-frequency fluctuations in the amount of ENSO activity itself (Chen *et al.*, 2004) so here we examine seasonal prediction skill for ENSO in our two resolutions for the same set of boreal winter and summer seasons.

Figure 1 shows ensemble ENSO predictions for the standard and higher-resolution models. Both models produce slightly overactive predictions, with ENSO anomalies in both warm El Nino and cold La Nina cases exceeding the observed amplitude. The predictions are also overconfident in the sense that the ensemble members cluster around the ensemble mean but do not span the observed values in some cases, as is well known for tropical seasonal predictions (Weisheimer and Palmer, 2014; Scaife *et al.*, 2018). Comparison of the standard and high-resolution cases gives a very clear message: the skill scores are almost identical in the two systems, albeit with slightly lower skill in summer than winter, consistent with the lower amplitude anomalies in summer. Similarly, the overprediction of the strength of anomalies and the overconfidence of the ensembles is very similar across the two resolutions. Seasonal prediction skill for ENSO therefore appears to be insensitive to the doubling of atmospheric model resolution tested here.

4 | PREDICTION SKILL OF TROPICAL CYCLONES

Seasonal forecasts of tropical cyclone numbers and activity has been carried out using empirical–statistical methods based on ENSO and other factors for many years (Klotzbach *et al.*, 2017). Over the last two decades, coupled ocean–





FIGURE 1 Seasonal prediction skill of ENSO at different atmospheric resolutions. Ensemble predictions are shown for standard resolution (left) and higher resolution (right) for summer (JJA, upper) and winter (DJF, lower). Anomalies in Niño3.4 are plotted for the observations (black), the ensemble mean (red line) and ensemble members (red dots). Correlation skill scores are shown for each case and observed indices are from HadISST1.1 (Rayner *et al.*, 2003)

atmosphere general circulation models have been shown to skilfully predict tropical cyclones on seasonal timescales (Vitart *et al.*, Vitart and Stockdale, 2001) and these are now improving to show skill on specific regional scales (Vecchi

et al., 2014), with some skill in landfall predictions (Camp *et al.*, 2015). As the record of events grows, there is also a growing number of successes in real-time prediction of extreme tropical cyclone seasons (e.g., Camp *et al.*, 2018).

FIGURE 2 Bias in tropical cyclone track frequency. The climatological bias in tropical cyclone track frequency is plotted for standard resolution (upper) and higher resolution (lower) hindcasts. Differences between the forecast and observations are plotted for the June–November period 1993–2015. Observed tropical cyclone data for the North Atlantic and East Pacific are taken from HURDAT2 (Landsea and Franklin, 2013), for all remaining basins, data are from the Joint Typhoon Warning Centre best-track database (Chu *et al.*, 2002)



Given their horizontal scale and structure, there is sensitivity of tropical cyclone simulations and short range forecasts to horizontal model resolution (e.g., Davis *et al.*, 2010; Gopalakrishnan *et al.*, 2012) and so we examine here whether there is also a sensitivity of seasonal prediction skill to resolution for the commonly used metrics of storm numbers and accumulated cyclone energy.

We first examine the climatology of storm numbers in hindcasts. Figure 2 shows the difference between simulated and observed storm track frequency. The standard resolution model produces a reasonable climatology of storm numbers in the Atlantic basin but overestimates the number in the Pacific. Although this bias may depend to some extent on the choice of tracking method, we used the same method on the higher-resolution storms to give a fair comparison (see Camp *et al.*, 2015). In this case increased resolution gives a small improvement in storm numbers over the Atlantic, but the excess of storms in the Pacific actually increases and there is no improvement to the overestimate of storm numbers in the North Indian Ocean, so there is little overall benefit to climatological storm track frequency over the range of resolutions considered here.

Table 1 shows the skill of standard resolution hindcasts compared to the skill from higher-resolution hindcasts for each tropical ocean basin and the Northern Hemisphere as a whole. Positive skill in storm numbers is found in all basins (except for the North Indian Ocean) and at both resolutions, with highly significant and potentially useful levels of skill in the Atlantic and East Pacific. However, we find no statistically significant change in skill from the doubling of resolution between standard and higher resolution. Similar results follow for accumulated cyclone energy (Table 2) where again, good skill levels are found in the Atlantic, the East Pacific and in this case, also the West Pacific. However, as with cyclone numbers, there is no significant change in skill with doubled atmospheric resolution, suggesting that any benefits to seasonal prediction of tropical storms from

TABLE 1 Prediction skill for numbers of hurricanes/tropical cyclones

	Standard resolution	Higher resolution
North Atlantic	0.62	0.53
Eastern Pacific	0.69	0.66
Western Pacific	0.26	0.28
North Indian	−0.18	−0.02
Northern Hemisphere	0.42	0.67

Note. Correlation scores are shown for ensemble predictions of hurricane strength storm numbers for the period June–November 1993–2015. Statistically significant values at the 95% confidence level are highlighted in bold.

TABLE 2 Prediction skill for accumulated cyclone energy

	Standard resolution	Higher resolution
North Atlantic	0.57	0.48
Eastern Pacific	0.80	0.84
Western Pacific	0.56	0.29
North Indian	−0.40	−0.42
Northern Hemisphere	0.76	0.76

Note. Correlation scores are shown for ensemble predictions of accumulated cyclone energy over the period June–November 1993–2015. Statistically significant values at the 95% confidence level are highlighted in bold.

increased resolution, at least in the range considered here, is likely to be small.

5 | PREDICTION SKILL OF THE NAO

The GloSea5 system produces skilful forecasts of the winter North Atlantic Oscillation (NAO) from initial conditions in early November (Scaife *et al.*, 2014) and in recent years, multiple seasonal forecast systems have been shown to produce skilful predictions of the winter NAO at this lead time (Athanasiadis *et al.*, 2014; Butler *et al.*, 2016; Baker *et al.*, 2018).

Figure 3 shows ensemble hindcasts of the winter NAO at the standard and higher resolutions. The correlation scores exceed 0.5 for the standard resolution and are similar to the skill reported by Scaife *et al.* (2014). There is no improvement in these scores at the higher resolution and correlation is nominally lower, although the value is within the uncertainty and therefore not significantly different (cf., Siebert *et al.*, 2016).

While seasonal predictions of the NAO have been demonstrated to contain significant skill, there is a prominent outstanding problem with the *amplitude* of the forecast NAO signals (Eade *et al.*, 2014; Scaife *et al.*, 2014). This results in ensemble predictions that contain inherently low levels of predictability in the sense that they are unable to skilfully predict single ensemble members and yet, they are still able to predict the observed NAO. This so-called signal-to-noise paradox (Dunstone *et al.*, 2016; Scaife and Smith, 2018) is present in different forecast systems (Baker *et al.*, 2018) and could in principle be due to the limited resolution of these systems. To test whether the signal to noise ratio of the standard and higher-resolution systems is the same, we compare the predictable (ensemble mean) and total (ensemble member) standard deviation of the NAO in our two sets of hindcasts. For the standard resolution case, the ensemble mean standard deviation is 2.5 hPa and the ensemble




FIGURE 3 Seasonal prediction skill of the NAO at different atmospheric resolutions: Ensemble predictions are shown for standard resolution (left) and higher resolution (right) for winter (DJF) NAO predictions. Anomalies in the NAO sea level pressure index between Iceland and Azores are plotted for the observations (black), the ensemble mean (red line) and ensemble members (red dots). Observed indices are taken from HadSLP2 (Allan and Ansell, 2006)

member standard deviation is 8.3 hPa, giving a ratio of around 0.3. For the higher-resolution case, the ensemble mean has a standard deviation of 2.3 hPa and the ensemble members have a standard deviation of 7.8 hPa. In this case the ratio remains 0.3 and so we detect no increase in the signal to noise ratio as the horizontal resolution of the forecast system is doubled. At least over the range considered here, the under-prediction of the strength of ensemble mean NAO signals noted in other studies of ensemble seasonal hindcasts is therefore insensitive to the horizontal atmospheric resolution of our prediction system.

6 | EFFECT OF HORIZONTAL RESOLUTION ON SMALL SCALE EDDY FEEDBACK

As resolution increases, we can expect to resolve more of the mesoscale eddy spectrum of the atmosphere which falls off relatively slowly with wavenumber ($\sim k^{-5/3}$) below scales of 100 km or so (Nastrom *et al.*, 1984). Eddies on this scale might be better resolved in our higher-resolution hindcasts. These high-frequency mesoscale eddies have also been shown to feedback positively onto larger scale anomalous flows in the atmosphere (Lau, 1988; Robinson, 1991;




FIGURE 4 Bias in the North Atlantic winter jet and eddy momentum forcing. Climatological bias of the North Atlantic zonal wind at 850 hPa (m/s, upper panels) and eddy momentum flux convergence at 200 hPa (m/s/day, lower panels) standard resolution is shown on the left and higher resolution on the right. Black contours are mean climatology from observational reanalysis (Dee *et al.*, 2011) and shading shows the difference between predictions and observational analysis. The right column displays zonal averages across the Atlantic basin

Feldstein and Lee, 1998; Kug and Jin, 2009; Kang *et al.*, 2010) and so it might be possible to increase the strength of the NAO signals discussed above if resolution were increased to the point where mesoscale eddy feedback was better resolved.

We first examine eddy feedback on the Atlantic jet stream in our standard and higher-resolution hindcasts. Figure 4 shows the mean zonal winds and their biases relative to observational reanalysis winds in the standard and higher-resolution hindcasts for DJF. Both resolutions show a poleward bias in the jet location compared to observational reanalyses (Figure 4c). We also compare the high-frequency eddy momentum flux convergence in both sets of hindcasts:

$$F_y = -\frac{\partial(u'\bar{v}')}{\partial y},$$

where u' and v' are the transient components of the wind using 6 hr data and the overbar indicates the time mean. This quantity feeds positively into the momentum budget of the mid latitude jets as shown in Figure 4d,e but consistent with the bias in mean winds there is also a corresponding lack of eddy momentum flux convergence, which is too weak in both our standard and higher-resolution model ensembles (Figure 4e). Figure 4e also shows that there is only a small improvement as resolution is increased over the range considered here.

This result is consistent with weak eddy forcing in some other models (e.g., Willison *et al.*, 2013; Lu *et al.*, 2015). So if the eddy forcing is too weak in our models and if it is also insensitive to the resolution range tested in our hindcasts, then perhaps we need to increase resolution further to better represent the mesoscale eddy activity? Indeed, recent evidence suggests that there may be some sensitivity of the eddy driven jet to resolution (Lu *et al.*, 2015) and sensitivity of simulated extratropical cyclones to resolution at the ~ 10 km scale (Sheldon *et al.*, 2017). Unfortunately, although global models are now being developed at this latter scale and beyond, we do not have seasonal hindcasts at this higher resolution due to their computational cost. However, we do have a sample of global atmosphere-only simulations of 15 winters at our highest (0.14°) atmospheric resolution, from which we can calculate high-frequency eddy feedback. For consistency we compare these with parallel sets of atmosphere only simulations at our standard and higher resolutions and a set of lower-resolution simulations, each of which have 192 winters.

In order to relate results back to the strength of NAO anomalies, which are known to be at least partly driven by synoptic eddy feedback (Limpasuvan and Hartmann, 2000; Ren *et al.*, 2009), we calculate the synoptic eddy vorticity forcing following Yu and Lin (2016):

$$F_v = -\frac{f}{g_o} \nabla^{-2} (\nabla \cdot \bar{\mathbf{V}} \bar{\zeta}'),$$

where f is the Coriolis parameter, g_o is the standard acceleration due to gravity, \mathbf{V} is horizontal wind vector and ζ is relative vorticity. Primes indicate that a 2–8-day band-pass filter was applied to the 6-hourly data and the overbar indicates the time average. We then regress this feedback onto the centres of action of the NAO to give the anomalous eddy forcing per standard deviation of the NAO in our winter simulations.

Figure 5 shows the amount of eddy vorticity forcing and also the regressed amount of upper tropospheric geopotential




FIGURE 5 North Atlantic eddy feedback as a function of resolution. Modelled eddy vorticity forcing (left) and geopotential height (right) per unit standard deviation of the NAO. Values are calculated by regression onto the NAO index. Observational reanalysis shows higher values than found in models, especially for low-resolution models and the values only approach convergence at N1280 (0.14°) resolution or finer. The horizontal scale is proportional to grid box size

height per unit change of the NAO for simulations at the lower (1.88°), standard (0.83°), higher (0.35°) and highest (0.14°) resolutions. The same quantities are also plotted for observational reanalyses, which we can consider to have very high resolution.

The comparison of standard (0.83°) and lower (1.88°) resolution simulations yields a similar result as in Figure 4, with little change in eddy feedback between resolutions. This feedback is substantially weaker than in observational reanalyses in both cases and is weakest at the lower resolution. Interestingly, increase of resolution to 0.18° shows an increase in eddy forcing (Figure 5a) and a corresponding significant increase in geopotential height signatures in the NAO (Figure 5b). Although it is still lower than the observed value, the strength of the eddy feedback at this resolution and the relationship between upper tropospheric geopotential height and the surface NAO approaches the value seen in observational analyses. It seems that this underrepresentation of eddy feedbacks may therefore be important for large scale features in the atmosphere and the strength of predicted NAO signals. Significantly higher resolution than is currently used for seasonal predictions may therefore be needed to correct this error.

7 | CONCLUSIONS

We have examined the effects of more than doubling atmospheric resolution, over a range of 0.83° to 0.35° , on seasonal prediction skill in parallel ensemble hindcasts. Although no study can carry out a fully comprehensive analysis of seasonal hindcast skill, we examined the skill of the main tropical (ENSO) and extratropical (NAO) modes of variability and what are arguably the most devastating hydro-meteorological extreme events (tropical cyclones); all of which have previously been shown to exhibit predictability on seasonal timescales.

There are still questions about whether current general circulation models converge to realistic solutions as resolution is increased (e.g., Gustafson Jr. *et al.*, 2014) but benefits of increasing atmospheric resolution beyond the 1° scale have previously been shown for seasonal prediction of surface climate (Jia *et al.*, 2015), tropical cyclones (Chen and Lin, 2013) and extratropical storm tracks (Yang *et al.*, 2015). Here we further increased atmospheric resolution and find that doubling from around 0.8° to around 0.3° resolution is not enough to make a large difference to seasonal prediction skill, as indeed has been shown for simulation of atmospheric features such as blocking and ocean–atmosphere interaction (Schiemann *et al.*, 2017; Sheldon *et al.*, 2017; Wan *et al.*, 2018). However, we did find evidence for increased eddy feedback at yet higher resolution, which appears to feedback

positively onto the NAO. This suggests that significantly higher resolution than is currently available for operational predictions may be required to increase the strength of predicted signals in seasonal forecasts of the NAO and resolve the signal to noise paradox in current climate simulations (Scaife and Smith, 2018).

It is important to note that other aspects of resolution such as ocean resolution or vertical atmospheric resolution were not tested in this study. Nevertheless, there is evidence that improved ocean resolution can improve the fidelity of simulations relevant to seasonal prediction (Scaife *et al.*, 2011; Hewitt *et al.*, 2017). In particular, higher ocean resolution may increase the strength of ocean–atmosphere coupling (Minobe *et al.*, 2008; Kirtman *et al.*, 2017), although there is as yet limited evidence for increased prediction skill from increased ocean resolution. Similarly, existing literature shows that improved vertical resolution and domain can improve long range forecasts (Marshall and Scaife, 2010; Fereday *et al.*, 2012; Sigmond *et al.*, 2013), not least due to the effects of the stratosphere on extratropical prediction skill of the NAO (Scaife *et al.*, 2016), but this was also not tested here.

In summary we find little impact of a doubling of atmospheric horizontal resolution from 0.83° to 0.35° on seasonal predictions of ENSO, the NAO or tropical cyclones. However, we did find evidence for increased feedback from small scale eddies at much higher resolution that could strengthen predicted signals in seasonal forecasts of the NAO. Until this resolution can be implemented in ensemble prediction systems, it may be better to increase the ensemble size, vertical resolution or perhaps ocean resolution to improve the skill of global operational seasonal prediction systems.

ACKNOWLEDGEMENTS

This work was supported by the UK-China Research & Innovation Partnership Fund through the Met Office Climate Science for Service Partnership (CSSP) China as part of the Newton Fund, This work was supported by the Joint DECC/Defra Met Office Hadley Centre Climate Programme (GA01101), the UK Public Weather Service Program, H2020 EUCP project grant number GA 776613, H2020 PRIMAVERA project grant number GA641727 and the Natural Environment Research Council NE/S004645/1. The authors also acknowledge the UK NERC ACSIS project.

ORCID

Adam A. Scaife  <https://orcid.org/0000-0002-5189-7538>

Joanne Camp  <https://orcid.org/0000-0002-4567-9622>

Nick Dunstone  <https://orcid.org/0000-0001-6859-6814>

REFERENCES

- Allan, R. and Ansell, T. (2006) A new globally complete monthly historical gridded mean sea level pressure dataset (HadSLP2): 1850–2004. *Journal of Climate*, 19(22), 5816–5842. <https://doi.org/10.1175/JCLI3937.1>.
- Athanasiadis, P.J., Bellucci, A., Hermanson, L., Scaife, A.A., MacLachlan, C., Arribas, A., Materia, S., Borrelli, A. and Gualdi, S. (2014) The representation of atmospheric blocking and the associated low-frequency variability in two seasonal prediction systems (CMCC, Met-Office). *Journal of Climate*, 27, 9082–9100. <https://doi.org/10.1175/JCLI-D-14-00291.1>.
- Baker, L.H., Shaffrey, L.C., Sutton, R.T., Weisheimer, A. and Scaife, A.A. (2018) An intercomparison of skill and overconfidence/underconfidence of the wintertime North Atlantic Oscillation in multimodel seasonal forecasts. *Geophysical Research Letters*, 45, 7808–7817. <https://doi.org/10.1029/2018GL078838>.
- Barnston, A.G., Tippett, M.K., L'Heureux, M.L., Li, S. and DeWitt, D. G. (2012) Skill of real-time seasonal ENSO model predictions during 2002–11: Is our capability increasing? *Bulletin of the American Meteorological Society*, 93, 631–651. <https://doi.org/10.1175/BAMS-D-11-00111.1>.
- Berckmans, J., Woollings, T., Demory, M., Vidale, P. and Roberts, M. (2013) Atmospheric blocking in a high resolution climate model: influences of mean state, orography and eddy forcing. *Atmospheric Science Letters*, 14, 34–40. <https://doi.org/10.1002/asl2.412>.
- Butler, A.H., Arribas, A., Athanassiadou, M., Baehr, J., Calvo, N., Charlton-Perez, A., Deque, M., Domeisen, D.I.V., Frohlich, K., Hendon, H., Imada, Y., Ishii, M., Iza, M., Karpechko, A.Y., Kumar, A., MacLachlan, C., Merryfield, W.J., Muller, W.A., O'Neill, A., Scaife, A.A., Scinocca, J., Sigmund, M., Stockdale, T. N. and Yasuda, T. (2016) The Climate Historical Forecast Project: Do stratosphere-resolving models make better seasonal climate predictions in boreal winter? *Quarterly Journal of the Royal Meteorological Society*, 142, 1413–1427.
- Camp, J., Roberts, M., MacLachlan, C., Wallace, E., Hermanson, L., Brookshaw, A., Arribas, A. and Scaife, A.A. (2015) Seasonal forecasting of tropical storms using the Met Office GloSea5 seasonal forecast system. *Quarterly Journal of the Royal Meteorological Society*, 141, 2206–2219. <https://doi.org/10.1002/qj.2516>.
- Camp, J., Scaife, A.A. and Heming, J. (2018) Predictability of the 2017 North Atlantic hurricane season. *Atmospheric Science Letters*, 19, e813. <https://doi.org/10.1002/asl.813>.
- Cane, M.A., Zebiak, S.E. and Dolan, S.C. (1986) Experimental forecasts of El Niño. *Nature*, 321, 827–832.
- Chen, D., Cane, M.A., Kaplan, A., Zebiak, S.E. and Huang, D. (2004) Predictability of El Niño over the past 148 years. *Nature*, 428, 733–736. <https://doi.org/10.1038/nature02439>.
- Chen, J. and Lin, S.-J. (2013) Seasonal predictions of tropical cyclones using a 25-km-resolution general circulation model. *Journal of Climate*, 26, 380–398. <https://doi.org/10.1175/JCLI-D-12-00061.1>.
- Chu, J.H., Sampson, C.R., Levine, A.S. and Fukada, E. (2002) *The Joint Typhoon Warning Center tropical cyclone best-tracks, 1945–2000*. Monterey, CA: Naval Research Laboratory. Technical report: NRL/MR/7540-02-16.
- Davis, C., Wang, W., Dudhia, J. and Torn, R. (2010) Does increased horizontal resolution improve hurricane wind forecasts? *Weather and Forecasting*, 25, 1826–1841.
- Dee, D.P., Uppala, S.M., Simmons, A.J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M.A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A.C.M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A.J., Haimberger, L., Healy, S.B., Hersbach, H., Hólm, E.V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B.M., Morcrette, J.J., Park, B.K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.N. and Vitart, F. (2011) The ERA-Interim reanalysis: configuration and performance of the data assimilation system. *Quarterly Journal of the Royal Meteorological Society*, 137, 553–597. <https://doi.org/10.1002/qj.828>.
- Doi, T., Behera, S.K. and Yamagata, T. (2019) Merits of a 108-member ensemble system in ENSO and IOD predictions. *Journal of Climate*, 32, 957–972. <https://doi.org/10.1175/JCLI-D-18-0193>.
- Dunstone, N., Smith, D., Scaife, A.A., Hermanson, L., Eade, R., Robinson, N., Andrews, M. and Knight, J. (2016) Skilful predictions of the winter North Atlantic Oscillation one year ahead. *Nature Geoscience*, 9, 809–814.
- Eade, R., Smith, D., Scaife, A.A. and Wallace, E. (2014) Do seasonal to decadal climate predictions underestimate the predictability of the real world? *Geophysical Research Letters*, 41, 5620–5628. <https://doi.org/10.1002/2014GL061146>.
- Feldstein, S. and Lee, S. (1998) Is the Atmospheric Zonal Index driven by an eddy feedback? *Journal of the Atmospheric Sciences*, 55, 3077–3086.
- Fereday, D., Maidens, A., Arribas, A., Scaife, A.A. and Knight, J.R. (2012) Seasonal forecasts of Northern Hemisphere winter 2009/10. *Environmental Research Letters*, 7, 034031. <https://doi.org/10.1088/1748-9326/7/3/034031>.
- Gopalakrishnan, S.G., Goldenberg, S., Quirino, T., Zhang, X., Marks, F., Yeh, K., Atlas, R. and Tallapragada, V. (2012) Toward improving high-resolution numerical hurricane forecasting: influence of model horizontal grid resolution, initialization, and physics. *Weather and Forecasting*, 27, 647–666.
- Gustafson, W.I., Jr., Ma, P.-L. and Singh, B. (2014) Precipitation characteristics of CAM5 physics at mesoscale resolution during MC3E and the impact of convective timescale choice. *Journal of Advances in Modeling Earth Systems*, 6, 1271–1287. <https://doi.org/10.1002/2014MS000334>.
- Hewitt, H.T., Bell, M.J., Chassignet, E.P., Czaja, A., Ferreira, D., Griffies, S.M., Hyder, P., McClean, J.L., New, A.L. and Roberts, M.J. (2017) Will high-resolution global ocean models benefit coupled predictions on short-range to climate timescales? *Ocean Modelling*, 120(2017), 120–136. <https://doi.org/10.1016/j.ocemod.2017.11.002>.
- Imada, Y., Tatebe, H., Ishii, M., Chikamoto, Y., Mori, M., Arai, M., Watanabe, M. and Kimoto, M. (2015) Predictability of two types of El Niño assessed using an extended seasonal prediction system by MIROC. *Monthly Weather Review*, 143, 4597–4617. <https://doi.org/10.1175/MWR-D-15-0007.1>.
- Jia, L., Yang, X., Vecchi, G.A., Gudgel, R.G., Delworth, T.L., Rosati, A., Stern, W.F., Wittenberg, A.T., Krishnamurthy, L., Zhang, S., Msadek, R., Kapnick, S., Underwood, S., Zeng, F., Anderson, W.G., Balaji, V. and Dixon, K. (2015) Improved seasonal prediction of temperature and precipitation over land in a high-resolution GFDL climate model. *Journal of Climate*, 28, 2044–2062. <https://doi.org/10.1175/JCLI-D-14-00112.1>.

- Johnson, S.J., Stockdale, T.N., Ferranti, L., Balmaseda, M.A., Molteni, F., Magnusson, L., Tietsche, S., Decremet, D., Weisheimer, A., Balsamo, G., Keeley, S., Mogensen, K., Zuo, H. and Monge-Sanz, B. (2018) SEAS5: the new ECMWF seasonal forecast system. *Geoscientific Model Development Discussion*, 12, 1089–1117. <https://doi.org/10.5194/gmd-2018-228>.
- Jung, T., Miller, M.J., Palmer, T.N., Towers, P., Wedi, N., Achuthavarier, D., Adams, J.M., Altshuler, E.L., Cash, B.A., Kinter, J.L., Marx, L., Stan, C. and Hodges, K.I. (2012) High-resolution global climate simulations with the ECMWF model in project Athena: experimental design, model climate, and seasonal forecast skill. *Journal of Climate*, 25, 3155–3172. <https://doi.org/10.1175/JCLI-D-11-00265.1>.
- Kang, I.-S., Kug, J.S., Lim, M.J., and Choi, D.H. (2010) Impact of transient eddies on extratropical seasonal-mean predictability in DEMETER models. *Climate Dynamics*, 37, 509–519.
- Kirtman, B., Perlin, N. and Siqueira, L. (2017) Ocean eddies and climate predictability. *Chaos*, 27(12), 126902.
- Klotzbach, P.J., Saunders, M.A., Bell, G.D. and Blake, E.S. (2017) North Atlantic seasonal hurricane prediction. In: Wang, S.S., Yoon, J., Funk, C.C. and Gillies, R.R. (Eds.) *Climate Extremes*. Washington, DC: American Geophysical Union. <https://doi.org/10.1002/9781119068020.ch19>.
- Kopparla, P., Fischer, E.M., Hannay, C. and Knutti, R. (2013) Improved simulation of extreme precipitation in a high-resolution atmosphere model. *Geophysical Research Letters*, 40, 5803–5808. <https://doi.org/10.1002/2013GL057866>.
- Kug, J.-S. and Jin, F.-F. (2009) Left-hand rule for synoptic eddy feedback on low-frequency flow. *Geophysical Research Letters*, 36, L05709. <https://doi.org/10.1029/2008GL036435>.
- Kumar, A., Chen, M., Xue, Y. and Behringer, D. (2015) An analysis of the temporal evolution of ENSO prediction skill in the context of the equatorial Pacific Ocean observing system. *Monthly Weather Review*, 143, 3204–3213. <https://doi.org/10.1175/MWR-D-15-0035.1>.
- Landsea, C.W. and Franklin, J.L. (2013) Atlantic hurricane database uncertainty and presentation of a new database format. *Monthly Weather Review*, 141, 3576–3592. <https://doi.org/10.1175/mwr-d-12-00254.1>.
- Lau, N.-C. (1988) Variability of the observed mid-latitude storm tracks in relation to low-frequency changes in the circulation pattern. *Journal of Atmospheric Sciences*, 45(19), 2718–2743.
- Limpasuvan, V. and Hartmann, D.L. (2000) Wave-maintained annular modes of climate variability. *Journal of Climate*, 13, 4414–4429. [https://doi.org/10.1175/1520-0442\(2000\)013<4414:WMAMOC>2.0.CO;2](https://doi.org/10.1175/1520-0442(2000)013<4414:WMAMOC>2.0.CO;2).
- Lu, J., Chen, G., Leung, L.R., Burrows, D.A., Yang, Q., Sakaguchi, K. and Hagos, S. (2015) Toward the dynamical convergence on the jet stream in Aquaplanet AGCMs. *Journal of Climate*, 28, 6763–6782. <https://doi.org/10.1175/JCLI-D-14-00761.1>.
- Luo, J.-J., Masson, S., Behera, S.K. and Yamagata, T. (2008) Extended ENSO predictions using a fully coupled ocean–atmosphere model. *Journal of Climate*, 21, 84–93.
- MacLachlan, C., Arribas, A., Peterson, K.A., Maidens, A., Fereday, D., Scaife, A.A., Gordon, M., Vellinga, M., Williams, A., Comer, R.E., Camp, J., Xavier, P. and Madec, G. (2014) Global seasonal forecast system version 5 (GloSea5): a high-resolution seasonal forecast system. *Quarterly Journal of the Royal Meteorological Society*, 141, 1072–1084. <https://doi.org/10.1002/qj.2396>.
- Marshall, A. and Scaife, A.A. (2010) Improved predictability of stratospheric sudden warming events in an AGCM with enhanced stratospheric resolution. *Journal of Geophysical Research*, 115, D16114. <https://doi.org/10.1029/2009JD012643>.
- Minobe, S., Kuwano-Yoshida, A., Komori, N., Xie, S.-P. and Small, R. J. (2008) Influence of the Gulf Stream on the troposphere. *Nature*, 452, 206–209.
- Nastrom, G.D., Gage, K.S. and Jasperson, W.H. (1984) Kinetic energy spectrum of large- and mesoscale atmospheric processes. *Nature*, 310, 36–38.
- Nie, Y., Scaife, A.A., Ren, H.-L., Comer, R.E., Andrews, M.B., Davis, P. and Martin, N. (2019) Stratospheric initial conditions provide seasonal predictability of the North Atlantic and Arctic Oscillations. *Environmental Research Letters*, 14, 034006.
- Prodhomme, C., Batté, L., Massonnet, F., Davini, P., Bellprat, O., Guemas, V. and Doblas-Reyes, F.J. (2016) Benefits of increasing the model resolution for the seasonal forecast quality in EC-earth. *Journal of Climate*, 29, 9141–9162. <https://doi.org/10.1175/JCLI-D-16-0117.1>.
- Rayner, N.A., Parker, D.E., Horton, E.B., Folland, C.K., Alexander, L. V., Rowell, D.P., Kent, E.C. and Kaplan, A. (2003) Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century. *Journal of Geophysical Research*, 108(D14), 4407. <https://doi.org/10.1029/2002JD002670>.
- Ren, H.-L., Jin, F.-F., Kug, J.-S., Zhao, J.-X. and Park, J. (2009) A kinematic mechanism for positive feedback between synoptic eddies and NAO. *Geophysical Research Letters*, 36, L11709. <https://doi.org/10.1029/2009GL037294>.
- Ren, H.-L., Scaife, A.A., Dunstone, N., Tian, B., Liu, Y., Ineson, S., Lee, J.Y., Smith, D., Liu, C., Thompson, V., Vellinga, M. and MacLachlan, C. (2019) Seasonal predictability of winter ENSO types in operational dynamical model predictions. *Climate Dynamics*, 52, 3869–3890. <https://doi.org/10.1007/s00382-018-4366-1>.
- Roberts, M.J., Vidale, P.L., Senior, C., Hewitt, H., Bates, C., Berthou, S., Chang, P., Christensen, H.M., Danilov, S., Demory, M., Griffies, S.M., Haarsma, R., Jung, T., Martin, G., Minobe, S., Ringler, T., Satoh, M., Schiemann, R., Scoccimarro, E., Stephens, G. and Wehner, M.F. (2018) The benefits of global high-resolution for climate simulation: process-understanding and the enabling of stakeholder decisions at the regional scale. *Bulletin of the American Meteorological Society*, 99, 2341–2359. <https://doi.org/10.1175/BAMS-D-15-00320.1>.
- Robinson, W.A. (1991) The dynamics of the zonal index in a simple model of the atmosphere. *Tellus A*, 43, 295–305.
- Scaife, A.A., Arribas, A., Blockley, E., Brookshaw, A., Clark, R.T., Dunstone, N., Eade, R., Fereday, D., Folland, C.K., Gordon, M., Hermanson, L., Knight, J.R., Lea, D.J., MacLachlan, C., Maidens, A., Martin, M., Peterson, A.K., Smith, D., Vellinga, M., Wallace, E., Waters, J. and Williams, A. (2014) Skilful long range prediction of European and North American winters. *Geophysical Research Letters*, 41, 2514–2519. <https://doi.org/10.1002/2014GL059637>.
- Scaife, A.A., Copsey, D., Gordon, C., Harris, C., Hinton, T., Keeley, S. J., O'Neill, A., Roberts, M. and Williams, K. (2011) Improved Atlantic blocking in a climate model. *Geophysical Research Letters*, 38, L23703. <https://doi.org/10.1029/2011GL049573>.
- Scaife, A.A., Ferranti, L., Alves, O., Athanasiadis, P., Baehr, J., Dequé, M., Dippe, T., Dunstone, N., Fereday, D., Gudgel, R.G., Greatbatch, R.J., Hermanson, L., Imada, Y., Jain, S., Kumar, A.,

- MacLachlan, C., Merryfield, W., Müller, W.A., Ren, H.L., Smith, D., Takaya, Y., Vecchi, G. and Yang, X. (2018) Tropical rainfall predictions from multiple seasonal forecast systems. *International Journal of Climatology*, 39, 1–15. <https://doi.org/10.1002/joc.5855>.
- Scaife, A.A., Karpechko, A.-Y., Baldwin, M.P., Brookshaw, A., Butler, A.H., Eade, R., Gordon, M., MacLachlan, C., Martin, N., Dunstone, N. and Smith, D. (2016) Seasonal forecasts and the stratosphere. *Atmospheric Science Letters*, 17, 51–56. <https://doi.org/10.1002/asl.598>.
- Scaife, A.A. and Smith, D. (2018) A signal-to-noise paradox in climate science. *Climate and Atmospheric Science*, 1, 28. <https://doi.org/10.1038/s41612-018-0038-4>.
- Schiemann, R., Demory, M., Shaffrey, L.C., Strachan, J., Vidale, P.L., Mizielinski, M.S., Roberts, M.J., Matsueda, M., Wehner, M.F. and Jung, T. (2017) The resolution sensitivity of Northern Hemisphere blocking in four 25-km atmospheric global circulation models. *Journal of Climate*, 30, 337–358. <https://doi.org/10.1175/JCLI-D-16-0100.1>.
- Sheldon, L., Czaja, A., Vannière, B., Morcrette, C., Sohet, B., Casado, M. and Smith, D. (2017) A “warm path” for Gulf Stream–troposphere interactions. *Tellus A*, 69(1), 1299397. <https://doi.org/10.1080/16000870.2017.1299397>.
- Siebert, S., Stephenson, D., Scaife, A.A., Eade, R. and Arribas, A. (2016) A Bayesian framework for verification and recalibration of ensemble forecasts: how uncertain is NAO predictability? *Journal of Climate*, 29, 995–1012.
- Sigmond, M., Scinocca, J.F., Kharin, V.V. and Shepherd, T.G. (2013) Enhanced seasonal forecast skill following stratospheric sudden warmings. *Nature Geoscience*, 6, 98–102. <https://doi.org/10.1038/ngeo1698>.
- Smith, D.M., Scaife, A.A. and Kirtman, B.P. (2012) What is the current state of scientific knowledge with regard to seasonal and decadal forecasting? *Environmental Research Letters*, 7, 015602.
- Tang, Y., Zhang, R.H., Liu, T., Duan, W., Yang, D., Zheng, F., Ren, H., Lian, T., Gao, C., Chen, D. and Mu, M. (2018) Progress in ENSO prediction and predictability study. *National Science Review*, 5(6), 826–839.
- Vecchi, G.A., Delworth, T., Gudgel, R., Kapnick, S., Rosati, A., Wittenberg, A.T., Zeng, F., Anderson, W., Balaji, V., Dixon, K., Jia, L., Kim, H., Krishnamurthy, L., Msadek, R., Stern, W.F., Underwood, S.D., Villarini, G., Yang, X. and Zhang, S. (2014) On the seasonal forecasting of regional tropical cyclone activity. *Journal of Climate*, 27, 7994–8016. <https://doi.org/10.1175/JCLI-D-14-00158.1>.
- Vitart, F. and Stockdale, T.N. (2001) Seasonal forecasting of tropical storms using coupled GCM integrations. *Monthly Weather Review*, 129, 2521–2537.
- Wan, J., Ren, H.-L. and Wu, P. (2018) Representation of the ENSO combination mode and its asymmetric SST response in different resolutions of HadGEM3. *Advances in Atmospheric Sciences*, 35 (8), 1063–1076.
- Weisheimer, A. and Palmer, T.N. (2014) On the reliability of seasonal climate forecasts. *Journal of the Royal Society Interface*, 11(96), 20131162. <https://doi.org/10.1098/rsif.2013.1162>.
- Willison, J., Robinson, W.A. and Lackmann, G.M. (2013) The importance of resolving mesoscale latent heating in the North Atlantic storm track. *Journal of Atmospheric Sciences*, 70, 2234–2250. <https://doi.org/10.1175/JAS-D-12-0226.1>.
- Yang, X., Vecchi, G.A., Gudgel, R.G., Delworth, T.L., Zhang, S., Rosati, A., Jia, L., Stern, W.F., Wittenberg, A.T., Kapnick, S., Msadek, R., Underwood, S.D., Zeng, F., Anderson, W. and Balaji, V. (2015) Seasonal predictability of extratropical storm tracks in GFDL’s high-resolution climate prediction model. *Journal of Climate*, 28, 3592–3611. <https://doi.org/10.1175/JCLI-D-14-00517.1>.
- Yao, J., Zhou, T., Guo, Z., Chen, X., Zou, L. and Sun, Y. (2017) Improved performance of high-resolution atmospheric models in simulating the East Asian summer monsoon rain belt. *Journal of Climate*, 30, 8825–8840. <https://doi.org/10.1175/JCLI-D-16-0372.1>.
- Yu, B. and Lin, H. (2016) Tropical atmospheric forcing of the wintertime North Atlantic Oscillation. *Journal of Climate*, 29, 1755–1772. <https://doi.org/10.1175/JCLI-D-15-0583>.
- Zhu, J., Huang, B., Cash, B., Kinter, J.L., Manganello, J., Barimalala, R., Altshuler, E., Vitart, F., Molteni, F. and Towers, P. (2015) ENSO prediction in project Minerva: sensitivity to atmospheric horizontal resolution and ensemble size. *Journal of Climate*, 28, 2080–2095.

How to cite this article: Scaife AA, Camp J, Comer R, *et al.* Does increased atmospheric resolution improve seasonal climate predictions? *Atmos Sci Lett.* 2019;e922. <https://doi.org/10.1002/asl.922>