

Fredholm theory of Toeplitz operators on doubling Fock Hilbert spaces

Article

Accepted Version

Al-Qabani, A., Hilberdink, T. and Virtanen, J. A. (2020)
Fredholm theory of Toeplitz operators on doubling Fock Hilbert
spaces. *Mathematica Scandinavica*, 126 (3). pp. 593-602.
ISSN 1903-1807 doi: 10.7146/math.scand.a-120920 Available
at <https://centaur.reading.ac.uk/85955/>

It is advisable to refer to the publisher's version if you intend to cite from the
work. See [Guidance on citing](#).

Published version at: <https://www.mscand.dk/forthcoming>

To link to this article DOI: <http://dx.doi.org/10.7146/math.scand.a-120920>

Publisher: *Mathematica Scandinavica*

All outputs in CentAUR are protected by Intellectual Property Rights law,
including copyright law. Copyright and IPR is retained by the creators or other
copyright holders. Terms and conditions for use of this material are defined in
the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

FREDHOLM THEORY OF TOEPLITZ OPERATORS ON DOUBLING FOCK HILBERT SPACES

AAMENA AL-QABANI, TITUS HILBERDINK AND JANI A. VIRTANEN

ABSTRACT. We study the Fredholm properties of Toeplitz operators acting on doubling Fock Hilbert spaces, and describe their essential spectra for bounded symbols of vanishing oscillation. We also compute the index of these Toeplitz operators in the special case when $\varphi(z) = |z|^\beta$ with $\beta > 0$. Our work extends the recent results on Toeplitz operators on the standard weighted Fock spaces to the setting of doubling Fock spaces.

1. INTRODUCTION

A positive Borel measure μ on the complex plane \mathbb{C} is said to be a *doubling measure* if there is a positive constant C such that

$$\mu(D(z, 2r)) \leq C\mu(D(z, r)),$$

for all $z \in \mathbb{C}$ and $r > 0$, where

$$D(z, r) = \{w \in \mathbb{C} : |w - z| < r\}.$$

We denote by dA the standard Lebesgue area measure on \mathbb{C} . Let φ be a subharmonic non-harmonic real-valued function of class C^2 on the complex plane \mathbb{C} such that $\Delta\varphi dA$ is a doubling measure, where $\Delta\varphi$ is the Laplacian of the function φ defined by

$$\Delta\varphi = \varphi_{xx} + \varphi_{yy}.$$

The *doubling Fock space* F_φ^2 is defined by

$$F_\varphi^2 = \left\{ f \in H(\mathbb{C}) : \|f\|_\varphi^2 = \int_{\mathbb{C}} |f(z)|^2 e^{-2\varphi(z)} dA(z) < \infty \right\},$$

where $H(\mathbb{C})$ is the set of all entire functions. These spaces were introduced in [6], where their sampling and interpolating sequences were described.

It is well known that F_φ^2 is a Hilbert space with inner product given by

$$\langle f, g \rangle = \int_{\mathbb{C}} f(w) \overline{g(w)} e^{-2\varphi(w)} dA(w).$$

We also note that the doubling Fock spaces include the standard weighted Fock spaces [14], the Fock-Sobolev space [3], the Fock spaces with weights $\varphi(z) = |z|^\beta$, where $\beta > 0$ (see [10]), and the generalized Fock spaces [11] with weights φ satisfying $0 < C_1 \leq \Delta\varphi(z) \leq C_2$, for all $z \in \mathbb{C}$, where C_1 and C_2 are positive constants.

2010 *Mathematics Subject Classification.* Primary 47B35; Secondary 30H20.

Key words and phrases. Toeplitz operators, Hankel operators, doubling Fock spaces, Fredholm properties, essential spectrum.

J. Virtanen was supported in part by Engineering and Physical Sciences Research Council grant EP/M024784/1.

When $\Delta\varphi dA$ is doubling, we say that a measurable function $f : \mathbb{C} \rightarrow \mathbb{C}$ is in L_φ^2 if

$$\|f\|_\varphi^2 = \int_{\mathbb{C}} |f(z)|^2 e^{-2\varphi(z)} dA(z) < \infty.$$

We denote by P the orthogonal projection of L_φ^2 onto F_φ^2 . It has the integral representation

$$Pf(z) = \int_{\mathbb{C}} f(w) \overline{K_z(w)} e^{-2\varphi(w)} dA(w),$$

where K_z is the reproducing kernel function; that is, for each $z \in \mathbb{C}$, K_z is the unique function in F_φ^2 for which

$$f(z) = \langle f, K_z \rangle,$$

for all $f \in F_\varphi^2$. Note that K_z depends on φ and we write

$$K_z(w) = K_\varphi(w, z) = K(w, z),$$

for $z, w \in \mathbb{C}$. For $z \in \mathbb{C}$, the *normalized reproducing kernel function* at z is defined by

$$k_z(w) = \frac{K_\varphi(w, z)}{\|K_\varphi(z, z)\|_\varphi},$$

for $w \in \mathbb{C}$.

For $f \in L^\infty = L^\infty(\mathbb{C})$, the *Toeplitz operator* T_f on F_φ^2 with *symbol* f is defined by

$$T_f(g) = P(fg).$$

It is clearly a bounded linear operator on F_φ^2 and $\|T_f\| \leq \|f\|_\infty$.

We say that a bounded linear operator T on a Banach space X is *Fredholm* if $\ker T$ and $X/T(X)$ are both finite-dimensional. In this case, the *index* of T is defined to be

$$\text{ind } T = \dim \ker T - \dim (X/T(X)).$$

The *essential spectrum* $\sigma_{\text{ess}}(T)$ is defined by

$$\sigma_{\text{ess}}(T) = \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not Fredholm}\},$$

where I denotes the identity operator on X . Clearly $\sigma_{\text{ess}}(T)$ is contained in the spectrum $\sigma(T)$ of T .

Our main results are concerned with the Fredholm properties and computation of the index of Toeplitz operators acting on doubling Fock spaces F_φ^2 with bounded symbols of vanishing oscillation. Before introducing the notion of vanishing oscillation, we make a few useful observations. Since $d\mu = \Delta\varphi dA$ is a doubling measure, by [12, §I.8.6], for any $z \in \mathbb{C}$ and $r > 0$, we have that

$$\mu(\partial(D(z, r))) = \mu(\{z\}) = 0$$

and $\mu(D(z, 2r)) \geq C\mu(D(z, r))$, where $C > 1$ is a constant. Moreover, our assumption that φ is non-harmonic implies that μ is a locally finite non-zero doubling measure on \mathbb{C} , so $0 < \mu(D(z, r)) < \infty$, for any $z \in \mathbb{C}$ and $r > 0$. It follows that, for each $z \in \mathbb{C}$, $\mu(D(z, r)) \rightarrow \infty$ as $r \rightarrow \infty$, and the function $r \mapsto \mu(D(z, r))$ is an increasing homeomorphism from the interval $(0, \infty)$ onto itself. Hence, for every $z \in \mathbb{C}$, there is a unique radius $\rho(z)$ such that

$$\mu(D(z, \rho(z))) = 1.$$

For $z \in \mathbb{C}$ and $r > 0$, we denote by $D^r(z)$ the disk $D(z, r\rho(z))$. Then we say that f is of *vanishing oscillation* and write $f \in VO$ if f is a continuous function on \mathbb{C} and

$$\omega_r(f)(z) = \sup_{w \in D^r(z)} |f(z) - f(w)| \rightarrow 0$$

as $z \rightarrow \infty$. Note that VO is independent of the choice of r .

2. MAIN RESULTS

The Fredholm properties of Toeplitz operators on the unweighted Fock space F^2 were described by Berger and Coburn [2] in 1987 using heavy machinery of C^* -algebras suitable for operators on Hilbert spaces. In 1992, Stroethoff [13] provided elementary proofs of their results, and very recently the theory was extended to the standard weighted Fock spaces F_α^p using elementary methods [1] and limit operator techniques [4]. In the following theorem, we extend the theory to doubling Fock Hilbert spaces F_φ^2 , which also sets the stage for further extensions to other doubling Fock spaces F_φ^p .

Theorem 1. *Let $f \in L^\infty \cap VO$. Then the Toeplitz operator T_f on F_φ^2 is Fredholm if and only if there is a radius $R > 0$ such that f is bounded away from zero on $\mathbb{C} \setminus D(0, R)$, that is, $\inf_{|z| \geq R} |f(z)| > 0$. In this case,*

$$\sigma_{\text{ess}}(T_f) = \bigcap_{R>0} \text{cl } f(\mathbb{C} \setminus D(0, R)),$$

where cl stands for the closure of the given set.

The difficulty with computing the index of Toeplitz operators on F_φ^2 is that doubling Fock spaces do not have simple bases unlike the standard weighted Fock spaces where index formulas can be obtained more easily; see [2] for Toeplitz operators on F^2 and [1] for these operators on F_α^p . The following result gives an index formula in the special case $\varphi(z) = \frac{1}{2}|z|^\beta$ with $\beta > 0$, which was introduced in [9] to study the properties of Hankel operators. For convenience, we write $F_{|z|^\beta}^2$ for $F_{\frac{1}{2}|z|^\beta}^2$. It is not difficult to see that $F_{|z|^\beta}^2$ is a doubling Fock space (and not a standard weighted Fock space).

Theorem 2. *Let $f \in L^\infty \cap VO$. Then the Toeplitz operator T_f on $F_{|z|^\beta}^2$ is Fredholm if and only if there is a radius $R > 0$ such that f is bounded away from zero on $\mathbb{C} \setminus D(0, R)$. In this case,*

$$\text{ind}(T_f) = -\text{wind}(f|_{\{|z|=R\}}),$$

where $\text{wind}(f|_{\{|z|=R\}})$ is the winding number of the curve $f(\{|z|=R\})$ around the origin.

The proofs of our main theorems are given in Section 4 below.

3. PRELIMINARIES

The following estimate for the Bergman kernel for F_φ^2 plays an important role in the study of concrete operators (such as Toeplitz and Hankel operators) on doubling Fock spaces (see [5, 8]).

Theorem 3 ([7, Theorem 1.1 and (3)]). *There are constants $C > 0$ and $\epsilon > 0$ such that*

$$|K(z_1, z_2)| \leq C \frac{1}{\rho(z_1)\rho(z_2)} \frac{e^{\varphi(z_1)+\varphi(z_2)}}{\exp(d_\varphi(z_1, z_2)^\epsilon)},$$

for any $z_1, z_2 \in \mathbb{C}$, where d_φ is the distance on \mathbb{C} induced by the metric $\varphi(z)^{-2}dz \otimes d\bar{z}$.

The previous theorem combined with the estimate (see [7, Proposition 2.10])

$$C^{-1} \frac{e^{\varphi(z)}}{\rho(z)} \leq \|K_\varphi(\cdot, z)\|_\varphi \leq C \frac{e^{\varphi(z)}}{\rho(z)}, \quad \text{for } z \in \mathbb{C}, \quad (1)$$

where C is a positive constant, gives the following result.

Lemma 4 ([8, Lemma 2.7]). *Let $r \geq 1$. For every $k \geq 0$ and $\epsilon > 0$ there exists a constant $C_{k,\epsilon}(r) > 0$ such that*

$$\int_{\mathbb{C} \setminus D^r(z_2)} \frac{|z_1 - z_2|^k}{\exp(d_\varphi(z_1, z_2)^\epsilon)} \frac{dA(z_1)}{\rho(z_1)^2} \leq C_{k,\epsilon}(r) \rho(z_2)^k, \quad \text{for any } z_2 \in \mathbb{C}.$$

Moreover, $C_{k,\epsilon}(r) \rightarrow 0$ as $r \rightarrow \infty$, for every $k \geq 0$ and $\epsilon > 0$.

For $f \in L^\infty$, the Berezin transform \tilde{f} of f is defined by

$$\tilde{f}(z) = \langle f k_z, k_z \rangle = \int_{\mathbb{C}} |k_z(w)|^2 f(w) e^{-2\varphi(w)} dA(w).$$

Lemma 5. *Let $f \in L^\infty \cap VO$. Then $\tilde{f}(z) - f(z) \rightarrow 0$ as $z \rightarrow \infty$.*

Proof. Let $\epsilon > 0$. By Theorem 3, (1) and Lemma 4, there is a radius $r \geq 1$ so that

$$\int_{\mathbb{C} \setminus D^r(z)} |k_z(w)|^2 e^{-2\varphi(w)} dA(w) < \frac{\epsilon}{\|f\|_\infty + 1},$$

for all $z \in \mathbb{C}$. Then

$$\begin{aligned} |\tilde{f}(z) - f(z)| &= \left| \int_{\mathbb{C}} (f(w) - f(z)) |k_z(w)|^2 e^{-2\varphi(w)} dA(w) \right| \\ &\leq \omega_r(f)(z) + 2\|f\|_\infty \int_{\mathbb{C} \setminus D^r(z)} |k_z(w)|^2 e^{-2\varphi(w)} dA(w) \\ &\leq \omega_r(f)(z) + 2\epsilon \leq 3\epsilon, \end{aligned}$$

when $|z|$ is large enough. Thus, $\tilde{f}(z) - f(z) \rightarrow 0$ as $|z| \rightarrow \infty$. □

For $f \in L^\infty$, the Hankel operator H_f on F_φ^2 with symbol f is defined by

$$H_f(g) = (I - P)(fg).$$

Note that H_f is a bounded linear operator from F_φ^2 into L_φ^2 and $\|H_f\| \leq \|f\|_\infty$. A useful relationship between Toeplitz and Hankel operators is the well-known multiplication formula

$$T_f T_g = T_{fg} - H_{\bar{f}}^* H_g, \quad (2)$$

which holds for $f, g \in L^\infty$.

In order to characterize compact Hankel operators, we state the definition of functions of vanishing mean oscillation. We say that $f \in L^2_{\text{loc}}(\mathbb{C})$ is in VMO if

$$\lim_{z \rightarrow \infty} MO(f)(z) = 0,$$

where

$$MO(f)(z) = \left(\frac{1}{A(D^1(z))} \int_{D^1(z)} |f - \hat{f}(z)|^2 dA \right)^{1/2}$$

and

$$\hat{f}(z) = \frac{1}{A(D^1(z))} \int_{D^1(z)} f dA.$$

The following two results can be found in [5].

Proposition 6. *We have that*

$$VMO = VO + VA,$$

where

$$VA = \{h \in L^2_{\text{loc}}(\mathbb{C}) : \lim_{z \rightarrow \infty} \widehat{|h|^2}(z) = 0\} = \{h \in L^2_{\text{loc}}(\mathbb{C}) : \lim_{z \rightarrow \infty} \widehat{|h|^2}(z) = 0\}.$$

Theorem 7. *Let $f \in L^\infty$. Then H_f and $H_{\bar{f}}$ are both compact operators from F^2_φ into L^2_φ if and only if $f \in VMO$.*

We finish this section with some basic properties of the Fock spaces $F^2_{|z|^\beta}$, where $\beta > 0$. For any non-negative integer n , we define e_n by

$$e_n(z) = \frac{z^n}{C_n}, \tag{3}$$

where

$$C_n^2 = \langle z^n, z^n \rangle_\beta = \int_{\mathbb{C}} |z^n|^2 e^{-|z|^\beta} dA(z) = \frac{2\pi}{\beta} \Gamma\left(\frac{2n+2}{\beta}\right). \tag{4}$$

Then $\{e_n\}_{n \geq 0}$ is an orthonormal basis for $F^2_{|z|^\beta}$, and it follows that the reproducing kernel function is given by

$$K_\beta(z, w) = \sum_{n=0}^{\infty} \frac{(z\bar{w})^n}{C_n^2},$$

for $z, w \in \mathbb{C}$, and the orthogonal projection has the following integral representation

$$\begin{aligned} P_\beta f(z) &= \int_{\mathbb{C}} f(w) K_\beta(z, w) e^{-|w|^\beta} dA(w) \\ &= \sum_{n=0}^{\infty} \frac{z^n}{C_n^2} \int_{\mathbb{C}} f(w) \bar{w}^n e^{-|w|^\beta} dA(w). \end{aligned} \tag{5}$$

4. PROOFS OF THE MAIN RESULTS

For the proof of the following approximation result, see [2, Lemma 17] and [1, Proposition 9].

Proposition 8. *Let $f : \mathbb{C} \rightarrow \mathbb{C}$ be a continuous function in A , where $A = L^\infty(\mathbb{C}) \cap VO$ or $A = L^\infty(\mathbb{C}) \cap VMO$. Then f is bounded away from zero on $\mathbb{C} \setminus D(0, R)$, for some $R > 0$, if and only if there is a continuous function $g \in A$ such that $f(z)g(z) \rightarrow 1$ as $z \rightarrow \infty$.*

Theorem 9. *If $f \in C(\mathbb{C})$ and $f(z) \rightarrow 0$ as $z \rightarrow \infty$, then T_f is compact on F_φ^2 .*

Proof. It directly follows from the proof of (2) \Rightarrow (1) in [8, Theorem 5.4]. \square

Proof of Theorem 1. Let $f \in L^\infty(\mathbb{C}) \cap VO$ and suppose that there are constants $R > 0$ and $m > 0$ such that $|f(z)| \geq m$ when $|z| > R$. By Proposition 8, there exists $g \in L^\infty(\mathbb{C}) \cap VO$ such that $fg - 1 \rightarrow 0$ as $|z| \rightarrow \infty$. Then

$$T_f T_g = I + T_{fg-1} - PM_f H_g,$$

where M_f is the multiplication operator on L_φ^2 defined by $M_f(h) = fh$. By Theorems 7 and 9, both T_{fg-1} and H_g are compact. Therefore, $T_f T_g = I + K$ for some compact operator K . Similarly, $T_g T_f = I + K_1$, where K_1 is a compact operator. Thus, T_f has a regularizer and hence it is Fredholm.

Conversely, suppose that there are $z_j \in \mathbb{C}$ such that $|z_j| \rightarrow \infty$ and $|f(z_j)| \rightarrow 0$. Then, by Lemma 5, we have that $\widetilde{|f|}(z_j) \rightarrow 0$. Since

$$\widetilde{|f|^2}(z) = \int_{\mathbb{C}} |f(w)|^2 |k_z(w)|^2 e^{-2\varphi(w)} dA(w) \leq \|f\|_\infty \widetilde{|f|}(z),$$

for $z \in \mathbb{C}$, we get that $\widetilde{|f|^2}(z_j) \rightarrow 0$. Now (2) gives that

$$\begin{aligned} \widetilde{|f|^2}(z_j) &= \langle T_{|f|^2} k_{z_j}, k_{z_j} \rangle = \langle (T_f T_f + H_f^* H_f) k_{z_j}, k_{z_j} \rangle \\ &= \langle T_f k_{z_j}, T_f k_{z_j} \rangle + \langle H_f k_{z_j}, H_f k_{z_j} \rangle = \|T_f k_{z_j}\|_\varphi^2 + \|H_f k_{z_j}\|_\varphi^2. \end{aligned}$$

Moreover, since $k_{z_j} \rightarrow 0$ weakly in F_φ^2 and H_f is compact on F_φ^2 (by Theorem 7), $\|H_f k_{z_j}\|_\varphi^2 \rightarrow 0$. Therefore, $\|T_f k_{z_j}\|_\varphi^2 \rightarrow 0$, and hence T_f is not invertible modulo compact operators, or equivalently, T_f is not Fredholm. \square

We now switch our focus and consider Toeplitz operators on $F_{|z|^\beta}^2$ with $\beta > 0$. To prove the index formula of Theorem 2, we consider first Toeplitz operators with symbols of the simple form $(z/|z|)^m$.

Lemma 10. *For $m \in \mathbb{N}$, let $\psi_m(z) = (z/|z|)^m$. Then T_{ψ_m} is an m -weighted shift operator on $F_{|z|^\beta}^2$, that is, there are numbers $\alpha_{m,n} > 0$ such that $T_{\psi_m} e_n = \alpha_{m,n} e_{n+m}$ for all $n \geq 0$, where $\{e_n\}_{n \geq 0}$ is defined by (3). Moreover, for any $m \in \mathbb{N}$, $\alpha_{m,n} \rightarrow 1$ as $n \rightarrow \infty$.*

Proof. By (5),

$$\begin{aligned} P_\beta(\psi_m e_n)(z) &= \frac{1}{C_n} \sum_{k=0}^{\infty} \frac{z^k}{C_k^2} \int_{\mathbb{C}} \bar{w}^k \frac{w^{n+m}}{|w|^m} e^{-|w|^\beta} dA(w) \\ &= \frac{1}{C_n} \frac{z^{n+m}}{C_{n+m}^2} \int_{\mathbb{C}} |w|^{2n+m} e^{-|w|^\beta} dA(w). \end{aligned}$$

Therefore,

$$P_\beta(\psi_m e_n) = \frac{e_{n+m}}{C_n C_{n+m}} \int_{\mathbb{C}} |w|^{2n+m} e^{-|w|^\beta} dA(w) = \frac{e_{n+m}}{C_n C_{n+m}} \frac{2\pi}{\beta} \Gamma\left(\frac{2n+m+2}{\beta}\right),$$

and so $T_{\psi_m} e_n = \alpha_{m,n} e_{n+m}$ with

$$\alpha_{m,n} = \frac{1}{C_n C_{n+m}} \frac{2\pi}{\beta} \Gamma\left(\frac{2n+m+2}{\beta}\right).$$

Then, by (4), Stirling's formula shows that, for any m , $\alpha_{m,n} \rightarrow 1$ as $n \rightarrow \infty$. \square

Proposition 11. *For every $m \in \mathbb{N}$, the Toeplitz operator $T_{\psi_m}^-$ is a Fredholm operator on $F_{|z|^\beta}^2$ of index m .*

Proof. Let

$$f(z) = \sum_{n=0}^{\infty} \lambda_n e_n$$

be a function in $F_{|z|^\beta}^2$. By Lemma 10,

$$T_{\psi_m} f = \sum_{n=0}^{\infty} \lambda_n T_{\psi_m} e_n = \sum_{n=0}^{\infty} \lambda_n \alpha_{m,n} e_{n+m},$$

where $\alpha_{m,n} \neq 0$, for any $n \geq 0$, and $\alpha_{m,n} \rightarrow 1$ as $n \rightarrow \infty$. It follows that $\dim \ker T_{\psi_m} = 0$ and $\dim(F_{|z|^\beta}^2 / T_{\psi_m}(F_{|z|^\beta}^2)) = m$. Therefore, T_{ψ_m} is a Fredholm operator of index $-m$, and hence $T_{\psi_m}^- = T_{\psi_m}^*$ is Fredholm of index m . \square

We can now prove the index formula of Toeplitz operators on $F_{|z|^\beta}^2$ with symbols in the class $L^\infty \cap VO$.

Proof of Theorem 2. Taking into account Proposition 8 and its proof, Theorem 9, (2), and Proposition 11, this proof is mutatis mutandis the proof of [1, Theorem 20]. \square

REFERENCES

- [1] A. Al-Qabani and J. A. Virtanen. Fredholm theory of Toeplitz operators on standard weighted Fock spaces. *Ann. Acad. Sci. Fenn. Math.*, 43(2):769–783, 2018.
- [2] C. A. Berger and L. A. Coburn. Toeplitz operators on the Segal-Bargmann space. *Trans. Amer. Math. Soc.*, 301(2):813–829, 1987.
- [3] H. R. Cho and K. Zhu. Fock-Sobolev spaces and their Carleson measures. *J. Funct. Anal.*, 263(8):2483–2506, 2012.
- [4] R. Fulsche and R. Hagger. Fredholmness of Toeplitz Operators on the Fock Space. *Complex Anal. Oper. Theory*, 13(2):375–403, 2019.
- [5] Z. Hu and X. Lv. Hankel operators on weighted Fock spaces (in Chinese). *Sci. China Math.*, 46(2):141–156, 2016.
- [6] N. Marco, X. Massaneda, and J. Ortega-Cerdà. Interpolating and sampling sequences for entire functions. *Geom. Funct. Anal.*, 13(4):862–914, 2003.
- [7] J. Marzo and J. Ortega-Cerdà. Pointwise estimates for the Bergman kernel of the weighted Fock space. *J. Geom. Anal.*, 19(4):890–910, 2009.
- [8] R. Oliver and D. Pascuas. Toeplitz operators on doubling Fock spaces. *J. Math. Anal. Appl.*, 435(2):1426–1457, 2016.
- [9] G. Schneider. Hankel operators with antiholomorphic symbols on the Fock space. *Proc. Amer. Math. Soc.*, 132(8):2399–2409, 2004.

- [10] G. Schneider and K. A. Schneider. Generalized Hankel operators on the Fock space. *Math. Nachr.*, 282(12):1811–1826, 2009.
- [11] A. P. Schuster and D. Varolin. Toeplitz operators and Carleson measures on generalized Bargmann-Fock spaces. *Integral Equations Operator Theory*, 72(3):363–392, 2012.
- [12] E. M. Stein. *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*. Princeton Mathematical Series, 43. Monographs in Harmonic Analysis, III. Princeton University Press, Princeton, NJ, 1993.
- [13] K. Stroethoff. Hankel and Toeplitz operators on the Fock space. *Michigan Math. J.*, 39(1):3–16, 1992.
- [14] K. Zhu. *Analysis on Fock spaces*. Graduate Texts in Mathematics, 263. Springer, New York, 2012.

DEPARTMENT OF MATHEMATICS AND COMPUTER APPLICATIONS, AL-NAHRAIN UNIVERSITY, BAGHDAD, IRAQ

E-mail address: aminarasim@hotmail.com

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF READING, WHITEKNIGHTS, READING RG6 6AX, ENGLAND

E-mail address: t.w.hilberdink@reading.ac.uk

E-mail address: j.a.virtanen@reading.ac.uk