

# *Stable isotope ratios - nutritional biomarkers of long-term intake?*

Article

Accepted Version

Kuhnle, G. G. C. ORCID: <https://orcid.org/0000-0002-8081-8931> (2019) Stable isotope ratios - nutritional biomarkers of long-term intake? American Journal of Clinical Nutrition, 110 (6). pp. 1265-1267. ISSN 0002-9165 doi: 10.1093/ajcn/nqz239 Available at <https://centaur.reading.ac.uk/86359/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1093/ajcn/nqz239>

Publisher: American Society for Nutrition

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

**CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

# 1 Stable isotope ratios – nutritional biomarker and more

2 Gunter G. C. Kuhnle

3 Department of Food and Nutritional Sciences

4 University of Reading, Reading, UK

5 email address: [g.g.kuhnle@reading.ac.uk](mailto:g.g.kuhnle@reading.ac.uk)

7 Sources of financial support: none

8 The author has no conflict of interest to disclose.

12 The accurate assessment of dietary intake is one of the biggest challenges in nutrition research. While  
13 there have been considerable advances in the development of new methods (1,2), many of the  
14 fundamental problems remain – in particular the bias introduced by misreporting and the limitations  
15 of food composition data (3). Nutritional biomarkers can address many of these problems as they  
16 measure actual intake and do not rely on self-reporting or food composition data to estimate intake  
17 (4). However, there is still a paucity of biomarkers, especially biomarkers that have been evaluated in  
18 controlled dietary intervention studies with known intakes.

19 Many biomarkers are based directly on the compounds of interest, such as micronutrients or  
20 fatty acids (5), or their metabolites (6). The inter-individual variability in metabolism, and more  
21 importantly the huge variability in food composition, make them generally unsuitable to estimate  
22 intake of foods or dietary patterns. They are also unable to distinguish between different sources of  
23 the self-same compound, for example added sugars from intrinsic sugars or different sources of fatty  
24 acids.

25 Yun and colleagues (7) have investigated a very different type of nutritional biomarker –  
26 natural abundance stable isotope ratios – which can be used to estimate intake of foods or dietary  
27 patterns. They are well established in ecological and archaeological research, where they are used to

reconstruct diet and food-web patterns (8). While it is often assumed that different isotopes behave in the same way, the small differences in the masses of the nuclides, as well as differences in quadrupole and magnetic moment (9), can result in a discrimination between different isotopes (*isotope effect*). While this isotopic fractionation is usually small (measured in *per mille* differences from a defined standard), it can be measured very reliably in bulk material such as blood or urine, or in individual compounds. The isotopic composition reflects the history of a molecule (9): increasing trophic levels for example result in an enrichment of  $^{15}\text{N}$ , and the differences in photosynthesis between C3 and C4 plants result in differences in the enrichment of  $^{13}\text{C}$  (11). This has been used extensively in Archaeology, for example to show the transition from fishing to farming during the Neolithic in Europe (10), investigate long-term dietary trends (13) or the introduction of maize in North America (11).

Despite the common use of stable isotope ratios in Archaeology and Ecology, they have been scarcely used in nutrition and nutritional epidemiology, and only very few studies have investigated their suitability as nutritional biomarker. The most common application so far has been the identification of dietary patterns, in particular the intakes of animal-derived foods like meat and fish. Petzke and colleagues (12) have used samples from a German nutritional survey (VERA) to show that carbon and nitrogen stable isotope ratios ( $\delta^{13}\text{C}$  and  $\delta^{15}\text{C}$ ) can be used to identify the intake of animal derived food. O'Brien and colleagues (13) could demonstrate that  $\delta^{15}\text{N}$  is a marker of fish and fish-derived fatty acid (EPA and DHA) intake and can therefore provide an alternative to laborious and expensive fatty acid analysis. In a small-scale feeding study, we could show that carbon and nitrogen stable isotope ratios can be used to distinguish between a vegan or vegetarian diet, and high meat or fish intake (14). However, many of these studies relied on extremes of intake and Hülsemann and colleagues showed that they are less sensitive to smaller changes in intake (15).

Another application is the use as biomarker of sugars intake, which is very difficult to assess from dietary data alone (16). In North America, the majority of sugars, especially in sugar-sweetened beverages (SSB), are derived from C4 plants, corn and sugar cane, whereas most other plants in the food supply are C3 plants. Foods containing sweeteners derived from corn or sugar cane therefore

have a distinct range of  $\delta^{13}\text{C}$  (17) and their consumption affects serum (21), whole blood (18) and hair stable isotope ratios (19) sufficiently to identify consumers. This method has been refined by using  $\delta^{13}\text{C}$  of glucose (20) and alanine (21) in blood, which is more specific for sugars and sugar-sweetened beverages.

There have been only few applications of stable isotope ratios to investigate associations between diet and health. In one study, Williams and O'Connell could show a positive association between  $\delta^{15}\text{N}$  and cognition in patients with Alzheimer's disease (22). In a series of studies in the Yup'ik population in Alaska,  $\delta^{15}\text{N}$  was used as biomarker of marine food intake to investigate gene  $\times$  diet interactions and DNA methylation patterns (8). In a case-cohort study of type 2 diabetes, Patel and colleagues (23) found positive associations between  $\delta^{15}\text{N}$  and incident diabetes, but inverse associations for  $\delta^{13}\text{C}$ . In particular the results of the last study highlight the need for a better evaluation of stable isotope ratios as nutritional biomarkers, as they can not only be affected by diet but also other factors.

The study by Yun and colleagues (7) in this issue is therefore very topical and provides important data for the evaluation of stable isotope ratios as nutritional biomarkers, not only because of the study size, but in particular because of the study design. The evaluation of nutritional biomarkers requires reliable data of actual intake, which cannot be obtained from self-reported dietary data. Yet despite the limitations of self-reporting, many studies rely on it to estimate actual intake. However, without reliable data on actual intake, the evaluation of a candidate biomarker is not possible, and any observed associations are likely to be biased. In this study, like in previous studies in the Women's Health Initiative (5), participants were therefore provided with their habitual diet, ensuring that the actual diet consumed was known. This should be the standard study design for biomarker evaluation, as it is the only method to obtain reliable data. The results of this evaluation clearly show an association between  $\delta^{15}\text{N}$  and the dietary intake of fish and seafood in a population with moderate intake, and of a combination of  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  with total animal protein intake. Interestingly, no associations between  $\delta^{13}\text{C}$  and added sugars or SSB were found, presumably due to the low intake in the study population.

The results also highlight one of the main challenges of using stable isotope ratios as nutritional biomarkers. Isotopic fractionation does not end when the food is consumed, but continues (24), and there are considerable differences between different tissues and thus the samples used for analysis (19,25,26). Isotopic fractionation of nitrogen depends on the availability of dietary nitrogen, especially from protein, and can be affected by factors such as pregnancy (27), nutritional stress (28) and changes in body mass (8). Lipids are generally depleted in  $^{13}\text{C}$  due to the preferences of enzymes in the biosynthetic pathways for  $^{12}\text{C}$  (34), thus having a lower  $\delta^{13}\text{C}$  than other tissues. The results of the study by Yun and colleagues (7), as well as data from the case-cohort study in EPIC Norfolk (23), show that other factors such as physical activity, BMI, smoking status and sex can also affect stable isotope ratios. They therefore not only provide information about dietary data, but also metabolic processes and subsequently the fate of dietary constituents.

The intricate nature of stable isotope ratios makes them a promising tool for future nutritional research. Yun and colleagues have demonstrated that the comparatively inexpensive bulk analysis of stable isotope ratios can be used as nutritional biomarker of fish/seafood and animal protein intake. Compound specific isotope ratio analysis can provide considerably more information: serum fatty acids reflect dietary fatty acid intake (35), and it is possible to distinguish between dairy and adipose fat of different animals using isotope ratio analysis of individual fatty acids (36). It is also possible to distinguish between endogenously and exogenously formed compounds, such as PUFAs (37), or to infer information on protein sources by determining the stable isotope ratios of individual amino acids (38). Many other methods have been developed for the analysis of specific compounds with applications in nutrition research, for example  $\delta^{15}\text{N}$  and  $\delta^{18}\text{O}$  of nitrate (39) to elucidate the role of exogenous and endogenous NO.

Stable isotope ratios have been used very successfully in other disciplines but have been underused in nutritional research for too long. The study by Yun and colleagues provides more evidence that this technique has an important place in nutritional research and should be used much more commonly.

## Acknowledgements

The sole author had responsibility for all parts of the manuscript.

## Notes

The author has no conflict of interest to disclose.

## References

1. Freedman LS, Schatzkin A, Midthune D, Kipnis V. Dealing With Dietary Measurement Error in Nutritional Cohort Studies. *J Natl Cancer Inst.* 2011;103:1086–92.
2. Subar AF, Freedman LS, Tooze JA, Kirkpatrick SI, Boushey C, Neuhouser ML, Thompson FE, Potischman N, Guenther PM, Tarasuk V, et al. Addressing Current Criticism Regarding the Value of Self-Report Dietary Data. *J Nutr.* 2015;145:2639–45.
3. Kuhnle GGC. Nutrition epidemiology of flavan-3-ols: The known unknowns. *Mol Aspects Med.* 2018;61:2–11.
4. Jenab M, Slimani N, Bictash M, Ferrari P, Bingham SA. Biomarkers in nutritional epidemiology: applications, needs and new horizons. *Hum Genet.* 2009;125:507–25.
5. Lampe JW, Huang Y, Neuhouser ML, Tinker LF, Song X, Schoeller DA, Kim S, Raftery D, Di C, Zheng C, et al. Dietary biomarker evaluation in a controlled feeding study in women from the Women's Health Initiative cohort. *Am J Clin Nutr.* 2017;105:466–75.
6. Ottaviani JJ, Fong R, Kimball J, Ensunsa JL, Britten A, Lucarelli D, Luben R, Grace PB, Mawson DH, Tym A, et al. Evaluation at scale of microbiome-derived metabolites as biomarker of flavan-3-ol intake in epidemiological studies. *Sci Rep.* 2018;8:9859.
7. Yun HY, Lampe JW, Tinker LF, Neuhouser ML, Beresford SAA, Niles KR, Mossavar-Rahmani Y, Snetselaar LG, Horn LV, Prentice RL, et al. Serum nitrogen and carbon stable isotope ratios meet biomarker criteria for fish and animal protein intake in a controlled feeding study of a Women's Health Initiative cohort. *Journal of Nutrition.* 2018.
8. O'Brien DM. Stable Isotope Ratios as Biomarkers of Diet for Health Research. *Annual Review of Nutrition.* 2015;35:565–94.
9. Isotope Fractionation: Why Aren't We What We Eat? 1999;26:667–73.
10. Richards MP, Schulting RJ, Hedges R. Archaeology: sharp shift in diet at onset of Neolithic. *Nature.* 2003.
11. van der Merwe NJ, Vogel J.  $^{13}\text{C}$  content of human collagen as a measure of prehistoric diet in woodland North America. *Nature.* 1978;276:815–6.

- 143 12. Petzke KJ, Boeing H, Klaus S, Metges CC. Carbon and nitrogen stable isotopic composition of  
144 hair protein and amino acids can be used as biomarkers for animal-derived dietary protein  
145 intake in humans. *J Nutr.* 2005;135:1515–20.
- 146 13. O'brien DM, Kristal AR, Jeannet MA, Wilkinson MJ, Bersamin A, Luick B. Red blood cell  
147  $\delta^{15}\text{N}$ : a novel biomarker of dietary eicosapentaenoic acid and docosahexaenoic acid  
148 intake. 2009 ed. 2009;89:913–9.
- 149 14. Kuhnle GGC, Joosen AMCP, Kneale CJ, O'Connell TC. Carbon and nitrogen isotopic ratios of  
150 urine and faeces as novel nutritional biomarkers of meat and fish intake. *Eur J Nutr.* 2012 ed.  
151 Springer-Verlag; 2013;52:389–95.
- 152 15. Huelsemann F, Flenker U, Koehler K, Schaenzer W. Effect of a controlled dietary change on  
153 carbon and nitrogen stable isotope ratios of human hair. 2009;23:2448–54.
- 154 16. Tasevska N. Urinary Sugars—A Biomarker of Total Sugars Intake. *Nutrients.*  
155 Multidisciplinary Digital Publishing Institute; 2015;7:5816–33.
- 156 17. Jahren AH, Saudek C, Yeung EH, Kao WH, Kraft RA, Caballero B. An isotopic method for  
157 quantifying sweeteners derived from corn and sugar cane. *Am J Clin Nutr.* 2006;84:1380–4.
- 158 18. Davy BM, Jahren AH, Hedrick VE, Comber DL. Association of  $\delta^{13}\text{C}$  in fingerstick blood with  
159 added-sugar and sugar-sweetened beverage intake. *J Am Diet Assoc.* 2011;111:874–8.
- 160 19. Nash SH, Kristal AR, Hopkins SE, Boyer BB, O'Brien DM. Stable Isotope Models of Sugar  
161 Intake Using Hair, Red Blood Cells, and Plasma, but Not Fasting Plasma Glucose, Predict  
162 Sugar Intake in a Yup'ik Study Population. *Journal of Nutrition.* 2014;144:75–80.
- 163 20. Cook CM, Alvig AL, Liu YQ, Schoeller DA. The Natural  $^{13}\text{C}$  Abundance of Plasma Glucose  
164 Is a Useful Biomarker of Recent Dietary Caloric Sweetener Intake. *Journal of Nutrition.*  
165 2010;140:333–7.
- 166 21. Kristal AR, Choy K, Nash SH, Hopkins S, Boyer BB, O'Brien DM. The carbon isotope ratio  
167 of alanine in red blood cells is a new candidate biomarker of sugar-sweetened beverage intake.  
168 *Journal of Nutrition.* American Society for Nutrition; 2013;143:878–84.
- 169 22. Williams JH, O'Connell TC. Differential relations between cognition and  $^{15}\text{N}$  isotopic content  
170 of hair in elderly people with dementia and controls. *J Gerontol A Biol Sci Med Sci.*  
171 2002;57:M797–802.
- 172 23. Patel PS, Cooper AJM, O'Connell TC, Kuhnle GGC, Kneale CK, Mulligan AM, Luben RN,  
173 Brage S, Khaw K-T, Wareham NJ, et al. Serum carbon and nitrogen stable isotopes as  
174 potential biomarkers of dietary intake and their relation with incident type 2 diabetes: the  
175 EPIC-Norfolk study. 2014;100:708–18.
- 176 24. O'Connell TC, Kneale CJ, Tasevska N, Kuhnle GG. The diet-body offset in human nitrogen  
177 isotopic values: a controlled dietary study. *American Journal of Physical Anthropology.*  
178 2012;149:426–34.
- 179 25. Kraft RA, Jahren AH, Saudek CD. Clinical-scale investigation of stable isotopes in human  
180 blood:  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  from 406 patients at the Johns Hopkins Medical Institutions.  
181 2008;22:3683–92.
- 182 26. Nash SH, Kristal AR, Boyer BB, King IB, Metzgar JS, O'Brien DM. Relation between stable  
183 isotope ratios in human red blood cells and hair: implications for using the nitrogen isotope



- 184 ratio of hair as a biomarker of eicosapentaenoic acid and docosahexaenoic acid. American  
185 Society for Nutrition; 2009;90:1642–7.
- 186 27. O'Connell TC. Nitrogen balance and  $\delta^{15}\text{N}$ : why you're not what you eat during pregnancy.  
187 2004;18:2889–96.
- 188 28. O'Connell TC. Nitrogen balance and  $\delta^{15}\text{N}$ : why you're not what you eat during nutritional  
189 stress. 2005;19:2497–506.
- 190