

Oxidative properties of Moringa oleifera kernel oil from different extraction methods during storage

Article

Supplemental Material

Mat Yusoff, M., Niranjan, K., Mason, O. A. and Gordon, M. H. (2020) Oxidative properties of Moringa oleifera kernel oil from different extraction methods during storage. *Journal of the Science of Food and Agriculture*, 100 (4). pp. 1588-1597. ISSN 1097-0010 doi: <https://doi.org/10.1002/jsfa.10167>
Available at <http://centaur.reading.ac.uk/87755/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1002/jsfa.10167>

Publisher: Wiley

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in


the [End User Agreement](#).

www.reading.ac.uk/centaur


CentAUR

Central Archive at the University of Reading


Reading's research outputs online



(a) storage temperature = 13 °C




(b) storage temperature = 25 °C




(c) storage temperature = 37 °C


Fig. 1. Effect of different extraction methods on the peroxide values of *Moringa oleifera* kernel oil stored for 140 days at different temperatures of (a) 13 °C; (b) 25 °C; and (c) 37 °C. SE, solvent extraction; AEE, aqueous enzymatic extraction; B-AEE, aqueous enzymatic extraction with boiling pre-treatment; HPP-AEE, aqueous enzymatic extraction with high pressure processing pre-treatment.



(a) storage temperature = 13 °C




(b) storage temperature = 25 °C




(c) storage temperature = 37 °C


Fig. 2. Effect of different extraction methods on the free fatty acids of *Moringa oleifera* kernel oil stored for 140 days at different temperatures of (a) 13 °C; (b) 25 °C; and (c) 37 °C. SE, solvent extraction; AEE, aqueous enzymatic extraction; B-AEE, aqueous enzymatic extraction with boiling pre-treatment; HPP-AEE, aqueous enzymatic extraction with high pressure processing pre-treatment.



(a) storage temperature = 13 °C




(b) storage temperature = 25 °C




(c) storage temperature = 37 °C


Fig. 3. Effect of different extraction methods on the alpha tocopherol content in *Moringa oleifera* kernel oil stored for 140 days at different temperatures of (a) 13 °C; (b) 25 °C; and (c) 37 °C. SE, solvent extraction; AEE, aqueous enzymatic extraction; B-AEE, aqueous enzymatic extraction with boiling pre-treatment; HPP-AEE, aqueous enzymatic extraction with high pressure processing pre-treatment.



(a) storage temperature = 13 °C



(b) storage temperature = 25 °C



(c) storage temperature = 37 °C

Fig. 4 Effect of different extraction methods on the gamma tocopherol content in *Moringa oleifera* kernel oil stored for 140 days at different temperatures of (a) 13 °C; (b) 25 °C; and (c) 37 °C. SE, solvent extraction; AEE, aqueous enzymatic extraction; B-AEE, aqueous enzymatic extraction with boiling pre-treatment; HPP-AEE, aqueous enzymatic extraction with high pressure processing pre-treatment.