

# *Sustainable intensification? Increased production diminishes omega-3 content of sheep milk*

Article

Supplemental Material

Open Access

Voutzourakis, N., Tzanidakis, N., Stergiadis, S. ORCID: <https://orcid.org/0000-0002-7293-182X>, Rempelos, L., Eyre, M., Atsali, I., Franceschin, E., Leifert, C., Stefanakis, A., Sotiraki, S. and Butler, G. (2020) Sustainable intensification? Increased production diminishes omega-3 content of sheep milk. Sustainability, 12 (3). 1228. ISSN 2071-1050 doi: 10.3390/su12031228 Available at <https://centaur.reading.ac.uk/88905/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.3390/su12031228>

Publisher: MPDI

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

## **CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

**Supplementary tables and figures** (not appearing with the paper when published but accessible if necessary)

**Table S1.** monthly temperature and rainfall during the study. (mean values  $\pm$ sem).

[illegible]

**Table S2.** Grazing regimes, supplementary concentrate, and conserved forages used in different management systems (production intensity),by sampling months and year. (mean values  $\pm$ sem).

Factors	grazing time on pasture:		Concentrate feeds g/ewe/day	conserved forage used:		
	hours/day	hours/day		total	oat	alfalfa
<u>Management System</u>						
Semi-intensive	3.5 $\pm$ 0.4	2.6 $\pm$ 0.1	934 $\pm$ 31	183 $\pm$ 21	68 $\pm$ 13	115 $\pm$ 17
Extensive	6.6 $\pm$ 0.4	0.4 $\pm$ 0.1	712 $\pm$ 40	267 $\pm$ 29	25 $\pm$ 9	242 $\pm$ 28
<u>Sampling Months</u>						
January	2.0 $\pm$ 0.4 <sup>e</sup>	1.2 $\pm$ 0.2 <sup>cd</sup>	1036 $\pm$ 40 <sup>a</sup>	473 $\pm$ 37 <sup>a</sup>	126 $\pm$ 27 <sup>a</sup>	347 $\pm$ 45 <sup>a</sup>
February	2.4 $\pm$ 0.5 <sup>de</sup>	1.7 $\pm$ 0.3 <sup>abc</sup>	1017 $\pm$ 42 <sup>a</sup>	423 $\pm$ 4 <sup>a</sup>	104 $\pm$ 26 <sup>a</sup>	319 $\pm$ 45 <sup>ab</sup>
March	3.4 $\pm$ 0.5 <sup>d</sup>	2.2 $\pm$ 0.3 <sup>a</sup>	942 $\pm$ 52 <sup>a</sup>	263 $\pm$ 39 <sup>b</sup>	23 $\pm$ 17 <sup>b</sup>	241 $\pm$ 38 <sup>b</sup>
April	5.5 $\pm$ 0.6 <sup>c</sup>	2.0 $\pm$ 0.3 <sup>ab</sup>	738 $\pm$ 61 <sup>b</sup>	103 $\pm$ 26 <sup>c</sup>	21 $\pm$ 13 <sup>b</sup>	82 $\pm$ 24 <sup>c</sup>
May	7.7 $\pm$ 0.6 <sup>b</sup>	1.3 $\pm$ 0.3 <sup>bcd</sup>	584 $\pm$ 68 <sup>c</sup>	58 $\pm$ 34 <sup>d</sup>	0 $\pm$ 0 <sup>b</sup>	58 $\pm$ 34 <sup>c</sup>
June	9.6 $\pm$ 0.6 <sup>a</sup>	0.6 $\pm$ 0.2 <sup>d</sup>	551 $\pm$ 66 <sup>c</sup>	10 $\pm$ 7 <sup>e</sup>	3 $\pm$ 3 <sup>b</sup>	7 $\pm$ 7 <sup>c</sup>
<u>Year</u>						
First	4.6 $\pm$ 0.4	1.5 $\pm$ 0.2	952 $\pm$ 34	290 $\pm$ 26	79 $\pm$ 14	212 $\pm$ 25
Second	5.5 $\pm$ 0.4	1.5 $\pm$ 0.2	698 $\pm$ 36	159 $\pm$ 23	14 $\pm$ 6	144 $\pm$ 22
<u>ANOVA (P-values)</u>						
<u>Main Effects</u>						
Management systems (MS)	***	***	***	**	***	***
Sampling Months (SM)	***	***	***	***	***	***
Year (Y)	**	ns	***	***	***	**
<u>Interactions</u>						
MS x SM	***	*	*** <sup>1</sup>	ns	** <sup>1</sup>	** <sup>1</sup>
MS x Y	ns	ns	ns	ns	* <sup>2</sup>	ns
SM x Y	ns	ns	ns	ns	*** <sup>3</sup>	* <sup>3</sup>
MS x SM x Y	ns	ns	ns	ns	*	ns

<sup>a-e</sup> Means within a column with different superscripts are significantly different ( $P$ -values < 0.05).

<sup>1</sup> see **Figure1** for interaction means/SE

<sup>2</sup> **Table S4** reports interaction means/SE.

<sup>3</sup> **Table S5** reports interaction means/SE.

**Table S3.** Interaction means±SEs for milk lactose content for the management systems and sampling months.

Sampling Months	January	February	March	April	May	June
Factors						
	lactose (g/100ml milk)					
semi-intensive system	4.77±0.04A <sup>ab</sup>	4.79±0.03A <sup>ab</sup>	4.72±0.07A <sup>b</sup>	4.82±0.05A <sup>ab</sup>	4.91±0.04A <sup>a</sup>	4.85±0.04A <sup>ab</sup>
extensive system	4.86±0.05A <sup>a</sup>	4.83±0.05A <sup>a</sup>	4.78±0.06A <sup>ab</sup>	4.74±0.08A <sup>ab</sup>	4.65±0.05B <sup>b</sup>	4.25±0.15B <sup>c</sup>
<sup>a-c</sup> Means within a row with different superscripts are significantly different ( <i>P</i> -values < 0.05).						
A-B Means within a column with different superscripts are significantly different ( <i>P</i> -values < 0.05).						

**Table S4.** Interaction means/SEs for total PUFA, omega-3 PUFA, lauric acid (C12:0), palmitic acid (C16:0), oleic acid (C18:1 *cis* 9), a-Linolenic acid (C18:3*cis*9.*cis*12.*cis*15), and eicosapentaenoic acid (EPA) in milk fat from two management systems and two years.

Parameter assessed	Year 1		Year 2	
	Semi intensive	Extensive	Semi intensive	Extensive
Wholecrop oat (g/ewe/day)	112 ±22 <sup>a</sup>	45 ±16 <sup>b</sup>	24 ±10 <sup>b</sup>	4 ±4 <sup>b</sup>
	FA (g/100g total FA)			
PUFA	5.13 ±0.11 <sup>c</sup>	5.00 ±0.13 <sup>c</sup>	6.33 ±0.09 <sup>b</sup>	6.69 ±0.19 <sup>a</sup>
n-3PUFA	0.74 ±0.04 <sup>c</sup>	0.83 ±0.07 <sup>c</sup>	1.20 ±0.04 <sup>b</sup>	1.50 ±0.09 <sup>a</sup>
C12:0	5.70 ±0.15 <sup>a</sup>	4.62 ±0.18 <sup>b</sup>	4.15 ±0.16 <sup>c</sup>	3.55 ±0.13 <sup>d</sup>
C16:0	28.28 ±0.26 <sup>b</sup>	29.24 ±0.31 <sup>a</sup>	26.06 ±0.24 <sup>c</sup>	26.14 ±0.22 <sup>c</sup>
C18:1 <i>cis</i> 9	17.69 ±0.32 <sup>c</sup>	20.41 ±0.53 <sup>b</sup>	20.87 ±0.51 <sup>ab</sup>	22.08 ±0.44 <sup>a</sup>
C18:3 <i>cis</i> 9. <i>cis</i> 12. <i>cis</i> 15	0.48 ±0.03 <sup>d</sup>	0.59 ±0.06 <sup>c</sup>	0.61 ±0.03 <sup>b</sup>	0.89 ±0.07 <sup>a</sup>
EPA	0.03 ±0.00 <sup>b</sup>	0.06 ±0.00 <sup>a</sup>	0.06 ±0.00 <sup>a</sup>	0.07 ±0.00 <sup>a</sup>

<sup>a-c</sup> Means within a row with different superscripts are significantly different (*P*-values < 0.05).

**Table S5.** Interaction mean values±SEs for whole-crop oat and alfalfa hay fed and palmitic acid and conjugated linoleic acid (CLA) concentrations in milk fat from two years and sampling months.


Sampling Months	January	February	March	April	May	June
Factors						
Oat hay (g /ewe/day)						
Year 1	196 ±40A <sup>a</sup>	183 ±45A <sup>a</sup>	45 ±33A <sup>b</sup>	43 ±25A <sup>b</sup>	0 ±0B <sup>c</sup>	6 ±6B <sup>c</sup>
Year 2	55 ±27B <sup>a</sup>	30 ±17B <sup>a</sup>	0 ±0B <sup>b</sup>	0 ±0B <sup>b</sup>	0 ±0B <sup>b</sup>	0 ±0B <sup>b</sup>
Alfalfa hay(g/ewe/day)						
Year 1	326 ±60A <sup>a</sup>	299 ±66A <sup>a</sup>	353 ±56A <sup>a</sup>	148 ±43A <sup>b</sup>	115 ±67A <sup>bc</sup>	14 ±14A <sup>c</sup>
Year 2	369 ±69A <sup>a</sup>	338 ±63A <sup>a</sup>	129 ±38B <sup>b</sup>	15 ±10B <sup>c</sup>	0 ±0B <sup>c</sup>	0 ±0A <sup>c</sup>
C16:0 (g/100g total FA)						
Year 1	27.67 ±0.32A <sup>b</sup>	27.66 ±0.41A <sup>b</sup>	28.24 ±0.32A <sup>b</sup>	27.97 ±0.31A <sup>b</sup>	30.43 ±0.66A <sup>a</sup>	30.87 ±0.37A <sup>a</sup>
Year 2	26.14 ±0.36B <sup>ab</sup>	25.65 ±0.50B <sup>ab</sup>	25.87 ±0.34B <sup>ab</sup>	25.55 ±0.52B <sup>b</sup>	26.74 ±0.31B <sup>a</sup>	26.68 ±0.24B <sup>a</sup>
CLAc9:111 (g/100g total FA)						
Year 1	0.79 ±0.06B <sup>c</sup>	0.79 ±0.06B <sup>c</sup>	0.96 ±0.05B <sup>ab</sup>	0.97 ±0.06B <sup>ab</sup>	1.06 ±0.10A <sup>a</sup>	0.86 ±0.04B <sup>bc</sup>
Year 2	0.95 ±0.07A <sup>c</sup>	1.17 ±0.07A <sup>ab</sup>	1.17 ±0.06A <sup>ab</sup>	1.29 ±0.07A <sup>a</sup>	1.09 ±0.05A <sup>bc</sup>	1.06 ±0.05A <sup>bc</sup>
<sup>a-c</sup> Means within a row with different superscripts are significantly different ( $P$ -values < 0.05). A-B Means within a column with different superscripts are significantly different ( $P$ -values < 0.05).						

**Table S6.** Effect of management systems (production intensity), sampling month, and year on concentrations of odd chain fatty acids in milk fat. (mean values  $\pm$ sem).

	decapentanoic acid C15:0	heptapentanoic acid C17:0	Total odd chain fatty acids	C15:0 : C17:0 ratio
Factors	g/100 g total FA			
<u>Management System</u>				
Semi-intensive	1.05 ±0.02	0.50 ±0.01	2.31 ±0.04	0.004 ±0.0003
Extensive	0.97 ±0.02	0.49 ±0.01	2.12 ±0.03	0.006 ±0.0004
<u>Sampling Months</u>				
January	0.92 ±0.03 <sup>ab</sup>	0.48 ±0.02 <sup>ab</sup>	2.16 ±0.06 <sup>ab</sup>	0.005 ±0.001 <sup>ab</sup>
February	0.95 ±0.03 <sup>b</sup>	0.49 ±0.02 <sup>ab</sup>	2.21 ±0.06 <sup>ab</sup>	0.004 ±0.001 <sup>a</sup>
March	1.03 ±0.03 <sup>c</sup>	0.46 ±0.02 <sup>b</sup>	2.30 ±0.06 <sup>a</sup>	0.004 ±0.001 <sup>a</sup>
April	1.06 ±0.03 <sup>c</sup>	0.49 ±0.02 <sup>ab</sup>	2.27 ±0.06 <sup>a</sup>	0.006 ±0.001 <sup>b</sup>
May	1.08 ±0.03 <sup>c</sup>	0.52 ±0.03 <sup>a</sup>	2.25 ±0.06 <sup>ab</sup>	0.005 ±0.001 <sup>a</sup>
June	1.01 ±0.04 <sup>cb</sup>	0.53 ±0.03 <sup>a</sup>	2.09 ±0.07 <sup>b</sup>	0.004 ±0.000 <sup>a</sup>
<u>Milking Season</u>				
First	1.02 ±0.02	0.40 ±0.01	2.15 ±0.04	0.005 ±0.0003
Second	1.00 ±0.02	0.58 ±0.01	2.28 ±0.03	0.005 ±0.003
<u>ANOVA (P-values)<sup>1</sup></u>				
<u>Main effects</u>				
Management (MS)	***	ns	***	**
Sampling months (SM)	***	*	ns	ns
Milking year (Y)	ns	***	**	ns
<u>Interactions</u>				
MS x SM	**	*	*	ns
MS x Y	ns	ns	ns	ns
SM x Y	**	***	*	*
MS x SM x Y	ns	ns	ns	ns


<sup>a-e</sup> Means within a column with different superscripts are significantly different according to Tukey's honestly significant difference test ( $P$ -values < 0.05).

<sup>1</sup> \*\*\*:  $p < 0.001$ , \*\*:  $p < 0.01$ ; \*:  $p < 0.05$ , t:  $p < 0.1$ , ns:  $p > 0.1$ .




**Figure S1.** Interaction means for the concentration of saturated fatty acids (SFA), Lauric acid (C12:0), Myristic acid (C14:0), Palmitic acid (C16:0) in milk fat and the respective Athrogenicity (AI) and Thrombogenicity (TI) indices of milk for from different management systems and sampling months. Semi-intensive management system is represented by (◆) and extensive management system by (◇). J:January, F:February, M:March, A:April, M:May, J:June. Values with different, capitalized letters represent statistically significant differences between the two management systems ( $P$ -value<0.05). Values with different lowercase letters represent statistically significant differences between months within the same system.





**Figure S2.** Interaction means the concentration of monounsaturated fatty acids (MUFA), polyunsaturated fatty acids (PUFA), Vaccenic acid (C18:1*trans*-11) and CLA*cis*9*trans*-11 in milk and oleic acid (C18:1 *cis*9) fat for the different management systems and sampling months. Semi-intensive management system is represented by (◆) and extensive management system by (◇). J:January, F:February, M:March, A:April, M:May, J:June. Values with different capital letters represent statistically significant differences between the two management systems ( $P$ -value < 0.05). Values with different lowercase letters represent statistically significant differences between months within the same system.



**Figure S3.** Interaction means for omega-3 PUFA (omega-3), omega-3 to omega-6 ratio, a-Linoleic acid (C18:3cis9.cis12.cis15; ALA), Eicosapentaenoic acid (C20:5omega-3; EPA), Docosapentaenoic acid (C22:5omega-3; DPA) and Docosahexaenoic acid (C22:6omega-3; DHA) for the different management systems and sampling months. Semi-intensive management system is represented by (♦) and extensive management system by (◆). J:January, F:February, M:March, A:April, M:May, J:June. Values with different capitalized letters represent statistically significant differences between the two management systems ( $P$ -value<0.05). Values with different lowercase letters represent statistically significant differences between months within the same system.