What the fluff is this? - Gammarus pulex prefer food sources without plastic microfibers

Article

Accepted Version

Yardy, L. and Callaghan, A. (2020) What the fluff is this? - Gammarus pulex prefer food sources without plastic microfibers. Science of the Total Environment, 715. 136815. ISSN 0048-9697 doi: <https://doi.org/10.1016/j.scitotenv.2020.136815> Available at <https://centaur.reading.ac.uk/89509/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1016/j.scitotenv.2020.136815>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

1 What the fluff is this? - *Gammarus pulex* prefer food sources without
2 plastic microfibers.

3 Lewis Yardy, Amanda Callaghan

4 Ecology and Evolutionary Biology, School of Biological Sciences, University of Reading,
5 Harborne Building, Reading RG6 6AS, UK

6 Microplastics, Microfibres, Pollution, Amphipoda

Abstract

Investigations into the impact of micro plastics (MP) and microfibers (MFs) upon the freshwater aquatic environment are still in their infancy despite our growing awareness of their importance. *Gammarus pulex* have long been used as a study organism for ecotoxicology and several studies have already used them to investigate the impact of MFs. One area of research which has not been exploited is the extent to which *G.pulex* can detect MFs and whether or not they avoid eating them. To answer this question we developed a reliable and accurate method of exposing *Gammarus* to known amounts of MF embedded in algal wafers. Here we show that when given the choice between control wafers and those contaminated with 2% or 3% MF *Gammarus* ingest fewer MF than would be expected if a random choice was made (2% W=7 P=0.01698, 3% W=13 P=0.03397). Their feeding behaviour also changes, with a significant reduction in time feeding ($F_{1,18}=21.3$ P=0.0002) as well as significantly fewer visits to contaminated wafers ($F_{1,18}= 5.312$ P=0.0333). This suggests that *G.pulex* are able to detect MF in the 200-500 μ m range and are partially repelled by them.

24 Introduction

25 Approximately 70% to 80% of microplastics (MPs) in marine environments are thought to
26 originate from inland sources and be transported out from rivers to the oceans (Andrady,
27 2011). Microplastics are defined as diverse plastics, including polyethylene and polystyrene,
28 whose fragments are smaller than 5 mm in size, they can be particles or fibres, fibres being
29 more than twice as long as they are thick and generally thinner than human hair (Cole,
30 Lindeque, Halsband, & Galloway, 2011). They can be produced by the degradation of larger
31 particles, for example through clothes washing (Browne et al., 2011; Napper & Thompson,
32 2016), or are manufactured as microbeads for use in personal care products including
33 toothpaste, sunscreen and facial scrubs (Duis & Coors, 2016; Fendall & Sewell, 2009;
34 Kalčíková, Alič, Skalar, Bundschuh, & Gotvajn, 2017; Leslie, 2014).

35 The highest volumes of MP pollution have been found in the Northern Hemisphere at water
36 fronts and in enclosed waters near to urban areas (Cózar et al., 2014; Barnes et al., 2009). As
37 well as accumulation in the environment (Cózar et al., 2014), MPs can accumulate in
38 individuals (Browne et al., 2008) and they have even been found in human stools (Schwabl et
39 al., 2018). Their size results in them being easily ingested by many aquatic organisms at
40 various trophic levels and stages of development, including freshwater invertebrates (Cole et
41 al., 2013; Scherer et al., 2017; Al-Jaibachi et al., 2018a, 2018b,; Aljaibachi and Callaghan,
42 2018). By entering the food chain MPs can be readily transferred between trophic levels (Chua
43 et al., 2014; Betts, 2008; Farrell and Nelson, 2013; Setälä et al., 2014; Davarpanah and
44 Guilhermino, 2015).

45 Studies to determine the impact of ingested MPs in smaller invertebrates such as copepods,
46 isopods and zooplankton have concluded that MPs have no detrimental effect following

47 ingestion, possibly because the MPs were too large to cross the midgut wall and were
48 eliminated in faeces (Cole et al., 2013; Cole, 2015). This was found in the isopod *Idotea*
49 *emarginata* (Hämer et al., 2014), cladoceran *Daphnia magna* (Al-Jaibachi and Callaghan,
50 2017) and dipteran mosquito *Culex pipiens* (Al-Jaibachi et al., 2018a, 2018b;; Aljaibachi and
51 Callaghan, 2018). In studies using the larger *Gammarus fossarum*, the impact of MP
52 ingestion varied depending on the type of plastic (Straub et al. 2017). Petroleum-based MPs
53 significantly reduced the assimilation efficiency of MP contaminated food in the long-term,
54 whereas biodegradable plastic did not, although ingestion of both types of plastic led to
55 significantly reduced growth compared to the control (Straub et al. 2017). In other studies,
56 Irregular MP fragments of polyethylene terephthalate (PET) had no negative effects on
57 feeding in *Gammarus pulex* (Weber et al. 2019).

58 *A meta-analysis on the impact of MP on the aquatic environment revealed that most studies*
59 *had focussed on particles rather than fibres (Foley, Feiner, Malinich, & Höök, 2018).*

60 Microfibres (MFs) have been investigated in several marine crustaceans, including Sand
61 Hoppers (*Orchestia gammarellus*), Shore Crabs (*Carcinus maenas*, *Carcinus aestuarii*) and
62 Langoustine (*Nephrops norvegicus*) concluding that MF between 1-5mm were ingested
63 (Piarulli et al., 2019; Watts, Urbina, Corr, Lewis, & Galloway, 2015; Welden & Cowie, 2016).
64 Welden & Cowie (2016) found that the number and length of MF retained in the digestive
65 tract of *N. norvegicus* was related to the gastric mill, an organ used to grind food in the
66 upper gut, larger specimens had larger gaps and so more and larger fibres could pass
67 through the gut and be excreted. They found that the only way for these trapped fibres to
68 be lost was through moulting, where their gut lining and gastric mill was shed.

69 Most studies into MF have focussed on the marine environment and have found that the
70 majority of fibres from the deep sea benthos were of cellulose origin (80%) with the
71 remainder being polyester or acrylic. Degradation in the ocean is linked to UV action, so that
72 plastic MFs in the deep sea tend to persist for hundreds if not thousands of years (Browne
73 et al., 2011; Sanchez-Vidal, Thompson, Canals, & De Haan, 2018). As the UV absorbance of
74 freshwater is greater than saltwater, and there is likely to be turbidity, there is likely to be a
75 similar problem in deeper river and lakes (Markager & Vincent, 2000).

76 The freshwater shrimp *G. pulex* has been used as a model organism for investigating a range
77 of topics within ecotoxicology, for example hormonal responses (Gismondi, 2018),
78 metabolic responses (Lebrun, Perret, Geffard, & Gourlay-Francé, 2012), the effect of
79 pesticides (Auber, Roucaute, Togola, & Caquet, 2011), and heavy metals (Duddridge &
80 Wainwright, 1980). *Gammarus pulex* are especially useful for investigating the impact of MP
81 because of their variable diet (Bloor, 2010, 2011; Kunz, Kienle, & Gerhardt, 2010). While
82 predominantly shredders feeding on leafy detritus, they will predate several invertebrate
83 taxa as well as feed upon carrion. In addition they are an essential food source for many
84 small fish (Kunz et al., 2010; MacNeil, Dick, & Elwood, 1999) and represent a vector for
85 plastics to enter the vertebrate food chain. *Gammaridae* are a diverse family of amphipod
86 crustaceans with representatives in freshwater, brackish and marine environments.
87 Therefore conclusions drawn from studying them are applicable all over the globe (Costa,
88 Neuparth, Correia, & Helena Costa, 2005; Kunz et al., 2010).

89 No recent studies have investigated how MP may affect feeding behaviour and may cause
90 selective feeding in *G. pulex*, nor have *G. pulex* been exposed to MF. Previous studies have
91 shown that several macroinvertebrates, including *G. pulex*, will ingest MP in a variety of

presentations, from as a suspension that settles on food (Weber, Scherer, Brennholt, Reifferscheid, & Wagner, 2018).

One difficulty in many studies into MFs has been that they are often studied without being incorporated into food sources and in concentrations well above environmentally relevant levels (Hanvey et al., 2017; Wagner et al., 2014). While some studies have produced a method for exposing invertebrates to a reliable dose of MP alongside plant matter (Straub, Hirsch, & Burkhardt-Holm, 2017), it is unknown how well they work with MF or larger MPs. It has been shown that algae and grasses provide a vector for MP into taxa not obviously at risk of MP ingestion (Goss, Jaskiel, & Rotjan, 2018; Gutow, Eckerlebe, Giménez, & Saborowski, 2016), therefore this relationship must be thoroughly investigated.

In this study we have adapted a method for dosing food with MFs that was originally developed to study plant litter decomposition and invertebrate consumption (Kampfraath et al., 2012). Our new method permits a reliable quantifiable method for exposing benthic macroinvertebrates to MFs. We used the method to identify whether *G.pulex* show any preference or repellence towards MF when they are part of a food source. This understanding is of utmost importance because it gives an idea as to the potential for environmental MF to enter the food chain. In order to gain a greater understanding behaviour must be investigated, previous studies have suggested that chronic exposure to MP impacts growth (Straub, Hirsch, & Burkhardt-Holm, 2017), thus making it less nutritious and could be a driver for food choice (Carrasco et al., 2019). However, if such avoidance is detected during the first exposure to MP then avoidance cannot be due to the lower nutritional value, as this has not yet been learned by individual organisms.

Materials and Methods

G. pulex Collection Site

The *G. pulex* were collected from Emm Brook, a tributary of the River Lodden, within Dinton Country Park in Reading, between the points (Decimal Degrees 51.440494, -0.874373 to 51.442274, -0.874359). This site was chosen for its good population of *G. pulex*, ease of access and because of its relatively shallow depth of <90cm. Animals over 12mm in length were collected by kick sampling using a hessian net, placed in plastic bottles filled with stream water and transported to the laboratory. The animals were briefly rinsed with reverse osmosis (RO) water in the laboratory to remove silt and river water and then species confirmed using a key (Eggers et al., 2016) .

In the laboratory *G. pulex* were placed in 45L plastic tanks (150 per tank) containing 40L aerated Organisation for Economic Co-operation and Development (OECD) reconstituted water (Hooper et al., 2006), maintained at 17°C with 12:12 light to dark ratio and fed algae wafers (Wafer Algae Eater Fish Food, API).

Microfibre Preparation

Black 100% acrylic wool (Hayfield Bonus DK product code 5723101001, Hobbycraft, Farnborough) was used to generate MFs. The wool was cut into pieces to generate lengths of <5mm by wrapping a length 5 times around two nails placed into a piece of wood 10 cm apart to generate ten parallel lengths. The wool was sprayed with RO water until it was saturated and then frozen at -80°C for 1 hour. After an hour the wool was removed and the first and last cm removed using a metal scalpel (Swann-Morton No 11 blade) and then cut

into 5cm lengths which were stored on ice until ready to be used. The wool lengths were further sliced into <500µm lengths and dried on a hot plate.

Wafer Production

Algae wafers, were ground using a mortar and pestle for 1 min until they were powder and stored in an airtight lidded glass beaker to prevent contamination. To make the wafers, 1g of the algae powder was added to 0.5ml of RO water and mixed to form a paste. The paste was shaped into a flat cake 5mm thick and placed on a hot plate at 70°C for 2 hours to dry. Test wafers were prepared by adding 0.5%, 1%, 1.5%, 2%, 2.5% and 3% MF fibres by weight to the powder and then homogenized by grinding for a further 1 min before adding the RO water.

Once dried each cake was cut up into 0.05 g wafers with a scalpel and placed in a separate lidded container to prevent contamination. To test the accuracy of this method for exposure of animals to known amounts of fibre, ten of each nominal concentration of test wafer were cut into quarters. Each quarter was crushed with a spatula and placed under a 10x binocular microscope (Optech Microtech) for counting.

Execution of Tests

Eight individual *Gammarus* 12-20mm in length were placed in a 5L aquarium filled with aerated 2L reconstituted water and starved for 24h. The *Gammarus* were then individually placed into an aerated 5L aquarium filled with 2L reconstituted water along with one 0.05g wafer (either control or treatment) and left for 4 hours to feed. After 4 hours each *Gammarus* was removed from its tank, placed in a 5ml beaker and killed with 50°C water. Eight tanks were used per day for 5 days, with concentrations distributed randomly across

the period, resulting in 10 replicates per treatment. Each day the aquariums were rotated in order to ensure that there was no impact from position.

Guts were removed from dead *Gammarus* under a binocular dissection microscope at 10X magnification. To remove the gut, the telson was removed with a second cut immediately behind the eyes. The gut was then pulled whole from the body using fine point forceps and picked through, counting the number of fibres.

Choice experiments were conducted using the same protocol, except each test aquarium had one 0.05g control wafer as well as a 0.05g test concentration wafer. The amount of time each *G.pulex* spent feeding on each wafer and the number of visits to each were recorded over four hours, this was referred to as behavioural data.

Data Analysis

All data analysis was conducted using R and R Studio. Shapiro-Wilkes tests were used to test for normality. The wafer data met the assumptions for normality and Two Way Analyses of Variance were conducted to see if there was any significant difference between wafers or wafer quadrants within each concentration. ANOVA was conducted between the concentrations in order to confirm significant difference in the number of MF between the concentrations.

The ingestion data met assumptions for normality therefore ANOVA was conducted to identify the relationship between the number of MF ingested and the concentration of MF in wafers.

The choice data did not meet the assumptions for normality, therefore Kruskal-Wallis tests were used in place of ANOVAs to investigate MF ingestion between concentrations. It was expected that the number of MF ingested would be half that of the non-choice experiment, however it was found that approximately half *G.pulex* ingested no MF, these were ignored and Wilcoxon Rank tests were used to investigate the difference between the treatments of choice and no choice of those *G.pulex* which did ingest MF.

Behaviour data fit the assumptions for normality and so ANOVAs were used to identify the functional response.

Results

Wafers

All wafers dried and set as expected and were easily dissected. There was no significant difference in acrylic fibre counts between wafers or wafer quadrants within each concentration (Table S1).

The number of fibres were directly proportional to the % of MF by mass added (Fig 1), and significantly different between concentrations $F_{1,118}=14766$ $P<0.0001$.

Figure 1. The number of fibres per quadrant of algae wafers made using different percentages (by mass) of 200-500µm Acrylic fibres, N at each concentration = 40.

Ingestion

The *G.pulex* readily fed on the test wafers and ingested MFs. Thirty percent of the 1% treatment and 10% of the 2% treatment ingested no MF. There was a direct relationship between wafer concentration and the number of MF eaten (Fig 2), with a significant difference between test concentrations $F_{1,28}=54.21$ $P<0.0001$.

Figure 2. The number of 200-500µm Acrylic fibres ingested by *G.pulex* in 4 hours at 3 test concentrations. N for each concentration = 10.

Choice experiments

Gammarus ingestion of MF approximately halved when animals were given a choice between contaminated and uncontaminated food (Fig 3). There was no significant difference in the number of MFs ingested between the concentrations when given a choice of uncontaminated food $H(2)=3.028$ $P=0.22$. Of the 12 *G. pulex* at each concentration, 4 of the 1%, 6 of the 2% and 5 of the 3% had ingested no MF, equating to approximately half of each concentration. When those that had ingested no MF were removed from the data and the remaining results were compared to the no-choice data, those *G. pulex* with a choice ingested significantly fewer MF than those without a choice Fig 4 (2% $W=7$ $P=0.017$, 3% $W=13$ $P=0.034$).

Fig 3. Linear Regressions for the ingestion of 200-500µm Acrylic fibres by *G. pulex*, with and without the choice of non-contaminated food. N for each concentration = 12.

A

Fig 4. The ingestion of 200-500µm Acrylic fibres by *G. pulex* with and without the choice of uncontaminated food at fibre concentrations (by mass) of 2% (A) and 3% (B)

The observation tests revealed that *G. pulex* spent significantly less time feeding ($F_{1,18}=21.3$ $P=0.0002$) on and significantly fewer visits ($F_{1,18}= 5.312$ $P=0.0333$) to contaminated wafers (Figure 5).

235 A

236

237

238 B

239

240 Fig 5. The amount of time in minuets *G.pulex* spend feeding from uncontaminated wafers and wafers
241 contaminated with 200-500µm Acrylic fibres (A) and the number of visits to each type of wafer (B).

242

243

Discussion

We have developed an accurate, cheap and easy method to produce wafers to investigate the impact of MFs on aquatic invertebrates based on the method of Straub et al., (2017). The wafers produced were homogenous within a concentration and MF counts were directly proportional to the % of MF used to produce the wafer. Therefore we can be confident that this method allows reliable dosing of MF which show a tendency to clump together without a solid matrix. *G.pulex* ingest plastic MFs in lengths up to at least 500µm in proportion to the concentration present.

This method allows researchers to instigate worst case scenarios where invertebrates may be unable to avoid MF and can be used to study preference between different MFs. This method would work for smaller MF and MP, and should be suitable for other organisms which will feed upon algae wafers, enabling a standardised method for understanding the impact of various MPs upon a range of environments.

There are several reasons why invertebrates may detect and avoid plastics in food, there could be chemical cues (De Lange, Sperber, & Peeters, 2006) or it could be they can physically feel their presence (Carrasco et al., 2019). If the main driving factor is the difference in texture between food and MP then the main food media texture should match the natural food texture as much as possible. An agar based gelatinous food source such as is used by Straub et al., (2017) produces a greater contrast between the food and the MP texture compared to this new method or natural food sources.

When given a choice of contaminated vs uncontaminated food, *Gammarus* significantly avoided eating food with MFs, with fewer visits to the food and a reduction in time feeding.

266 These observations were supported by quantitative data demonstrating a significant
267 difference in MFs ingested. *Gammarus* have previously avoided eating contaminated food
268 including when chemical cues to bacteria and fungi are present (De Lange, Lüring, Van Den
269 Borne, & Peeters, 2005; De Lange, Sperber, & Peeters, 2006). Furthermore there is evidence
270 that animals can detect and avoid MPs. Carrasco et al(2019) exposed *Orchestoidea*
271 *tuberculata* to artificial food containing 8 µm particles of polystyrene MP spheres at 3
272 different concentrations (0%, 5% and 10%). The animals consumed significantly more food
273 when no MPs were present compared to food contaminated with 10% MPs. As this study
274 was a relatively short exposure (15 days) it is possible that the avoidance mechanism is
275 physical rather than biochemical.

276 In the current study contaminated wafers were eaten with no evident repulsion when no
277 uncontaminated food was available. This is in line with other studies which have recorded
278 MF ingestion of fibres of up to 5mm in length by taxa larger than *Gammarus*, including
279 crustaceans, molluscs, annelids and fish, (Farrell & Nelson, 2013; Foley et al., 2018; Straub et
280 al., 2017; Watts et al., 2015). Similar results have been found in the smaller *Daphnia magna*
281 with many studies showing that there is a positive relationship between concentration of
282 MP and the number ingested (Canniff & Hoang, 2018; Jemec, Horvat, Kunej, Bele, & Kržan,
283 2016; Rehse, Kloas, & Zarfl, 2016). However, Aljaibachi & Callaghan (2018) found that
284 *Daphnia* seemed to be able to selectively ingest algal cells and avoid 2µm MP particles.

285 These results are important in understanding the risk to the environment. It suggests that,
286 at least *Gammarus* is able to avoid MF contaminated food, meaning that as long as their
287 environment is not totally saturated with MF they could be ingested in rates lower than one
288 might assume given environmental concentrations. As macroinvertebrates are the main

vector for MP entering the higher trophic levels (Foley et al., 2018), including vertebrates and ultimately humans, their ability to limit MP ingestion would in turn limit the amount entering higher trophic levels. There is already a highlighted knowledge gap in this area (Horton, Walton, Spurgeon, Lahive, & Svendsen, 2017) and its understanding would help direct mitigation processes.

Gammarus produce copious amounts of faecal pellets which are eaten by other freshwater macroinvertebrates and are important sources of organic matter for bacteria (Joyce, Warren, & Wotton, 2007) . Microfibres were clearly observed in faecal pellets with no evidence of being shortened which means that not only could *G.pulex* act as a vector for MP to enter higher trophic levels if they are eaten by fish or other invertebrates, but their faeces provide a source of MP to enter lower trophic levels through faecal ingestion (Kelly, Dick, & Montgomery, 2002; Ladle & Griffiths, 1980) (Kelly, Dick, & Montgomery, 2002).

Despite their apparent ability to avoid ingesting MF contaminated wafers, it remains to be seen whether *G. pulex* predation on differentially contaminated prey would vary.

- Al-Jaibachi, R., Cuthbert, R. N., & Callaghan, A. (2018a). Examining effects of ontogenic microplastic transference on *Culex* mosquito mortality and adult weight. *Science of The Total Environment*, 651, 871–876. <https://doi.org/10.1016/j.scitotenv.2018.09.236>
- Al-Jaibachi, R., Cuthbert, R. N., & Callaghan, A. (2018b). Up and away: ontogenic transference as a pathway for aerial dispersal of microplastics. *Biology Letters*, 14(9).
- Aljaibachi, R., & Callaghan, A. (2018a). Impact of polystyrene microplastics on *Daphnia magna* mortality and reproduction in relation to food availability. *PeerJ*, 6, e4601. <https://doi.org/10.7717/peerj.4601>
- Aljaibachi, R., & Callaghan, A. (2018b). Impact of polystyrene microplastics on *Daphnia magna* mortality and reproduction in relation to food availability. *PeerJ*, 6, e4601. <https://doi.org/10.7717/peerj.4601>
- Andrady, A. L. (2011). Microplastics in the marine environment. *Marine Pollution Bulletin*, 62(8), 1596–1605. <https://doi.org/10.1016/j.marpolbul.2011.05.030>
- Auber, A., Roucaute, M., Togola, A., & Caquet, T. (2011). Structural and functional effects of conventional and low pesticide input crop-protection programs on benthic macroinvertebrate communities in outdoor pond mesocosms. *Ecotoxicology*, 20(8), 2042–2055. <https://doi.org/10.1007/s10646-011-0747-5>
- Besseling, E., Wang, B., Lürling, M., & Koelmans, A. a. (2014). Nanoplastic Affects Growth of *S. obliquus* and Reproduction of *D. magna*. *Environmental Science & Technology*, 48(20), 12336–12343. <https://doi.org/10.1021/es503001d>
- Bloor, M. C. (2010). Animal standardisation for mixed species ecotoxicological studies : Establishing a laboratory breeding programme for *Gammarus pulex* and *Asellus aquaticus*. *Zool. Baetica*, 21, 179–190.
- Bloor, M. C. (2011). Dietary preference of *Gammarus pulex* and *Asellus aquaticus* during a laboratory breeding programme for ecotoxicological studies. *International Journal of Zoology*. <https://doi.org/10.1155/2011/294394>
- Browne, M. A., Crump, P., Niven, S. J., Teuten, E., Tonkin, A., Galloway, T., & Thompson, R. (2011). Accumulation of microplastic on shorelines worldwide: sources and sinks. *Environmental Science & Technology*, 45(21), 9175–9179. <https://doi.org/10.1021/es201811s>
- Canniff, P. M., & Hoang, T. C. (2018). Microplastic ingestion by *Daphnia magna* and its enhancement on algal growth. *Science of the Total Environment*, 633, 500–507. <https://doi.org/10.1016/j.scitotenv.2018.03.176>
- Carrasco, A., Pulgar, J., Quintanilla-Ahumada, D., Perez-Venegas, D., Quijón, P. A., & Duarte, C. (2019). The influence of microplastics pollution on the feeding behavior of a prominent sandy beach amphipod, *Orchestoidea tuberculata* (Nicolet, 1849). *Marine Pollution Bulletin*, 145, 23–27. <https://doi.org/10.1016/j.marpolbul.2019.05.018>
- Cole, M., Lindeque, P., Fileman, E., Halsband, C., Goodhead, R., Moger, J., & Galloway, T. S. (2013). Microplastic ingestion by zooplankton. *Environmental Science and Technology*, 47(12), 6646–6655. <https://doi.org/10.1021/es400663f>
- Cole, M., Lindeque, P., Halsband, C., & Galloway, T. S. (2011). Microplastics as contaminants in the

351 marine environment: A review. *Marine Pollution Bulletin*.
 352 <https://doi.org/10.1016/j.marpolbul.2011.09.025>

353 Costa, F. O., Neuparth, T., Correia, A. D., & Helena Costa, M. (2005). Multi-level assessment of
 354 chronic toxicity of estuarine sediments with the amphipod *Gammarus locusta*: II. Organism and
 355 population-level endpoints. *Marine Environmental Research*, 60, 93–110.
 356 <https://doi.org/10.1016/j.marenvres.2004.08.005>

357 De Lange, H. J., Lürling, M., Van Den Borne, B., & Peeters, E. T. H. M. (2005). Attraction of the
 358 amphipod *Gammarus pulex* to water-borne cues of food. *Hydrobiologia*, 544(1), 19–25.
 359 <https://doi.org/10.1007/s10750-004-7896-y>

360 De Lange, H. J., Sperber, V., & Peeters, E. T. H. M. (2006). Avoidance of polycyclic aromatic
 361 hydrocarbon-contaminated sediments by the freshwater invertebrates *Gammarus pulex* and
 362 *Asellus aquaticus*. *Environmental Toxicology and Chemistry*, 25(2), 452–457.
 363 <https://doi.org/10.1897/05-413.1>

364 Duddridge, J. E., & Wainwright, M. (1980). Heavy metal accumulation by aquatic fungi and reduction
 365 in viability of *Gammarus pulex* fed Cd²⁺ contaminated mycelium. *Water Research*, 14(11),
 366 1605–1611. [https://doi.org/10.1016/0043-1354\(80\)90065-2](https://doi.org/10.1016/0043-1354(80)90065-2)

367 Duis, K., & Coors, A. (2016). Microplastics in the aquatic and terrestrial environment: sources (with a
 368 specific focus on personal care products), fate and effects. *Environmental Sciences Europe*,
 369 28(1), 2. <https://doi.org/10.1186/s12302-015-0069-y>

370 Eggers, T. O., Martens, A., Hanselmann, A. J., Kenna, D., Fincham, W. N. W., Dunn, A. M., ... Pöckl, M.
 371 (2016). Bestimmungsschlüssel der Süßwasser-Amphipoda (Crustacea) Deutschlands.
 372 *Lauterbornia*, 42(1), 1–68. <https://doi.org/10.1007/s00442-016-3796-x>

373 Farrell, P., & Nelson, K. (2013). Trophic level transfer of microplastic: *Mytilus edulis* (L.) to *Carcinus*
 374 *maenas* (L.). *Environmental Pollution*, 177, 1–3. <https://doi.org/10.1016/j.envpol.2013.01.046>

375 Fendall, L. S., & Sewell, M. a. (2009). Contributing to marine pollution by washing your face:
 376 microplastics in facial cleansers. *Marine Pollution Bulletin*, 58(8), 1225–1228.
 377 <https://doi.org/10.1016/j.marpolbul.2009.04.025>

378 Foley, C. J., Feiner, Z. S., Malinich, T. D., & Höök, T. O. (2018). A meta-analysis of the effects of
 379 exposure to microplastics on fish and aquatic invertebrates. *Science of the Total Environment*.
 380 <https://doi.org/10.1016/j.scitotenv.2018.03.046>

381 Gismondi, E. (2018). Identification of molt-inhibiting hormone and ecdysteroid receptor cDNA
 382 sequences in *Gammarus pulex*, and variations after endocrine disruptor exposures.
 383 *Ecotoxicology and Environmental Safety*, 158, 9–17.
 384 <https://doi.org/10.1016/j.ecoenv.2018.04.017>

385 Goss, H., Jaskiel, J., & Rotjan, R. (2018). *Thalassia testudinum* as a potential vector for incorporating
 386 microplastics into benthic marine food webs. *Marine Pollution Bulletin*, 135, 1085–1089.
 387 <https://doi.org/10.1016/j.marpolbul.2018.08.024>

388 Gutow, L., Eckerlebe, A., Giménez, L., & Saborowski, R. (2016). Experimental Evaluation of Seaweeds
 389 as a Vector for Microplastics into Marine Food Webs. *Environmental Science and Technology*,
 390 50(2), 915–923. <https://doi.org/10.1021/acs.est.5b02431>

391 Hanvey, J. S., Lewis, P. J., Lavers, J. L., Crosbie, N. D., Pozo, K., & Clarke, B. O. (2017). A review of
 392 analytical techniques for quantifying microplastics in sediments. *Anal. Methods*, 9(9), 1369–
 393 1383. <https://doi.org/10.1039/C6AY02707E>

394 Hooper, H. L., Connon, R., Callaghan, A., Maund, S. J., Liess, M., Duquesne, S., ... Sibly, R. M. (2006).
395 The use of image analysis to estimate population growth rate in *Daphnia magna*. *Journal of*
396 *Applied Ecology*, 43(4), 828–834. <https://doi.org/10.1111/j.1365-2664.2006.01180.x>

397 Horton, A. A., Walton, A., Spurgeon, D. J., Lahive, E., & Svendsen, C. (2017). Microplastics in
398 freshwater and terrestrial environments: Evaluating the current understanding to identify the
399 knowledge gaps and future research priorities. *Science of the Total Environment*.
400 <https://doi.org/10.1016/j.scitotenv.2017.01.190>

401 Jemec, A., Horvat, P., Kunej, U., Bele, M., & Kržan, A. (2016). Uptake and effects of microplastic
402 textile fibers on freshwater crustacean *Daphnia magna*. *Environmental Pollution*, 219, 201–209.
403 <https://doi.org/10.1016/j.envpol.2016.10.037>

404 Joyce, P., Warren, L. L., & Wotton, R. S. (2007). Faecal pellets in streams: Their binding, breakdown
405 and utilization. *Freshwater Biology*, 52(10), 1868–1880. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2427.2007.01828.x)
406 [2427.2007.01828.x](https://doi.org/10.1111/j.1365-2427.2007.01828.x)

407 Kalčíková, G., Alič, B., Skalar, T., Bundschuh, M., & Gotvajn, A. Ž. (2017). Wastewater treatment plant
408 effluents as source of cosmetic polyethylene microbeads to freshwater. *Chemosphere*, 188, 25–
409 31. <https://doi.org/10.1016/j.chemosphere.2017.08.131>

410 Kelly, D. W., Dick, J. T. A., & Montgomery, W. I. (2002). The functional role of *Gammarus* (Crustacea,
411 Amphipoda): Shredders, predators, or both? *Hydrobiologia*, 485, 199–203.
412 <https://doi.org/10.1023/A:1021370405349>

413 Kunz, P. Y., Kienle, C., & Gerhardt, A. (2010). *Gammarus* spp. in aquatic ecotoxicology and water
414 quality assessment: Toward integrated multilevel tests. *Reviews of Environmental*
415 *Contamination and Toxicology*, 205, 1–76. https://doi.org/10.1007/978-1-4419-5623-1_1

416 Ladle, M., & Griffiths, B. S. (1980). A study on the faeces of some chalk stream invertebrates.
417 *Hydrobiologia*, 74(2), 161–171. <https://doi.org/10.1007/BF00014568>

418 Lebrun, J. D., Perret, M., Geffard, A., & Gourlay-Francé, C. (2012). Modelling copper bioaccumulation
419 in *Gammarus pulex* and alterations of digestive metabolism. *Ecotoxicology*, 21(7), 2022–2030.
420 <https://doi.org/10.1007/s10646-012-0955-7>

421 Leslie, H. A. IVM Institute for Environmental Studies Review of Microplastics in Cosmetics (2014).

422 Liu, Z., Yu, P., Cai, M., Wu, D., Zhang, M., & Huang, Y. (2019). Polystyrene nanoplastic exposure
423 induces immobilization, reproduction, and stress defense in the freshwater cladoceran *Daphnia*
424 *pulex*. *Chemosphere*, 215, 74–81. <https://doi.org/10.1016/j.chemosphere.2018.09.176>

425 MacNeil, C., Dick, J. T. A., & Elwood, R. W. (1999). The dynamics of predation on *Gammarus* spp.
426 (Crustacea: Amphipoda). *Biological Reviews*, 74, 375–395.
427 <https://doi.org/doi:10.1017/S0006323199005368>

428 Markager, S., & Vincent, W. F. (2000). Spectral light attenuation and the absorption of UV and blue
429 light in natural waters. *Limnology and Oceanography*, 45(3), 642–650.
430 <https://doi.org/10.4319/lo.2000.45.3.0642>

431 Napper, I. E., & Thompson, R. C. (2016). Release of synthetic microplastic plastic fibres from
432 domestic washing machines: Effects of fabric type and washing conditions. *Marine Pollution*
433 *Bulletin*, 112(1–2), 39–45. <https://doi.org/10.1016/j.marpolbul.2016.09.025>

434 Piarulli, S., Scapinello, S., Comandini, P., Magnusson, K., Granberg, M., Wong, J. X. W., ... Airoldi, L.
435 (2019). Microplastic in wild populations of the omnivorous crab *Carcinus aestuarii*: A review
436 and a regional-scale test of extraction methods, including microfibrils. *Environmental Pollution*,

- 117–127. <https://doi.org/10.1016/j.envpol.2019.04.092>
- Rehse, S., Kloas, W., & Zarfl, C. (2016). Short-term exposure with high concentrations of pristine microplastic particles leads to immobilisation of *Daphnia magna*. *Chemosphere*, 153, 91–99. <https://doi.org/10.1016/j.chemosphere.2016.02.133>
- Sanchez-Vidal, A., Thompson, R. C., Canals, M., & De Haan, W. P. (2018). The imprint of microfibrils in Southern European deep seas. *PLoS ONE*, 13(11). <https://doi.org/10.1371/journal.pone.0207033>
- Scherer, C., Brennholt, N., Reifferscheid, G., & Wagner, M. (2017). Feeding type and development drive the ingestion of microplastics by freshwater invertebrates. *Scientific Reports*, 7(1). <https://doi.org/10.1038/s41598-017-17191-7>
- Schwabl, P., Liebmann, B., Köppel, S., Königshofer, P., Bucsics, T., Trauner, M., & Reiberger, T. (2018). Assessment of microplastic concentrations in human stool - Preliminary results of a prospective study. *Assessment of Microplastic Concentrations in Human Stool - Preliminary Results of a Prospective Study*.
- Straub, S., Hirsch, P. E., & Burkhardt-Holm, P. (2017). Biodegradable and petroleum-based microplastics do not differ in their ingestion and excretion but in their biological effects in a freshwater invertebrate *Gammarus fossarum*. *International Journal of Environmental Research and Public Health*, 14(7). <https://doi.org/10.3390/ijerph14070774>
- Wagner, M., Scherer, C., Alvarez-Muñoz, D., Brennholt, N., Bourrain, X., Buchinger, S., ... Reifferscheid, G. (2014). Microplastics in freshwater ecosystems: what we know and what we need to know. *Environmental Sciences Europe*, 26(1), 1–9. <https://doi.org/10.1186/s12302-014-0012-7>
- Watts, A. J. R., Urbina, M. A., Corr, S., Lewis, C., & Galloway, T. S. (2015). Ingestion of Plastic Microfibers by the Crab *Carcinus maenas* and Its Effect on Food Consumption and Energy Balance. *Environmental Science and Technology*, 49(24), 14597–14604. <https://doi.org/10.1021/acs.est.5b04026>
- Weber, A., Scherer, C., Brennholt, N., Reifferscheid, G., & Wagner, M. (2018). PET microplastics do not negatively affect the survival, development, metabolism and feeding activity of the freshwater invertebrate *Gammarus pulex*. *Environmental Pollution*, 234, 181–189. <https://doi.org/10.1016/j.envpol.2017.11.014>
- Welden, N. A. C., & Cowie, P. R. (2016). Environment and gut morphology influence microplastic retention in langoustine, *Nephrops norvegicus*. *Environmental Pollution*, 214, 859–865. <https://doi.org/10.1016/j.envpol.2016.03.067>