

Investigation into the antimicrobial activity of fumarate against Listeria monocytogenes and its mode of action under acidic conditions

Article

Accepted Version

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Barnes, R. H. and Karatzas, K. A. G. (2020) Investigation into the antimicrobial activity of fumarate against *Listeria monocytogenes* and its mode of action under acidic conditions. *International Journal of Food Microbiology*, 324. 108614. ISSN 0168-1605 doi: <https://doi.org/10.1016/j.ijfoodmicro.2020.108614> Available at <https://centaur.reading.ac.uk/91061/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

Published version at: <http://dx.doi.org/10.1016/j.ijfoodmicro.2020.108614>

To link to this article DOI: <http://dx.doi.org/10.1016/j.ijfoodmicro.2020.108614>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

1 **Investigation into the antimicrobial activity of fumarate against *Listeria monocytogenes***
2 **and its mode of action under acidic conditions**

3

4 Ruth H. Barnes^a and Kimon Andreas G. Karatzas^{a*}

5 ^a Department of Food and Nutritional Sciences, School of Chemistry, Food and Pharmacy,
6 Department of Food & Nutritional Sciences, University of Reading, Reading RG6 6AD UK.

7

8 *Corresponding author.

9

10 Corresponding author address:

11 School of Chemistry, Food and Pharmacy

12 Department of Food & Nutritional Sciences,

13 University of Reading,

14 Reading

15 RG6 6AD

16 UK

17

18 Tel. +44 118 378 6678

19 Fax. +44 118 931 0080

20 Email: k.karatzas@reading.ac.uk

21

22 **Keywords:** *Listeria monocytogenes*, fumaric acid, glutamate decarboxylase, inhibition, GAD,
23 biofilm

24

25

26

27 **Abstract**

28 Organic acids such as fumarate are commonly used as antimicrobials in foods. Apart from the
29 classical mechanism of intracellular dissociation, weak acids are active through important
30 additional mechanisms which are not well-defined. Fumarate, based on its low dissociation
31 constants is expected to have a low antimicrobial activity which is not the case, suggesting
32 additional antimicrobial effects. Previously, fumarate has been shown to inhibit the GAD
33 system of *E. coli* and therefore, we investigated for first time how it affects this system in
34 *Listeria monocytogenes*. We found that fumarate is highly antimicrobial towards *L.*
35 *monocytogenes* under acidic conditions. We also show that in cell lysates and similarly to *E.*
36 *coli*, fumarate inhibits the GAD system of *L. monocytogenes*. However, despite the inhibition
37 and in contrast to *E. coli*, *L. monocytogenes* is able to counteract this and achieve a higher
38 extracellular GAD output (measured by GABA export) in the presence of fumarate compared
39 to its absence. The latter is achieved by a dramatic 9.44-fold increase in the transcription of
40 *gadD2* which is the main component of the extracellular GAD system. Interestingly, although
41 maleate, the cis-isomer of fumarate results in a more dramatic 48.5-fold *gadD2* upregulation
42 than that of fumarate, the final GAD_e output is lower suggesting that maleate might be a stronger
43 inhibitor of the GAD system. In contrast, the GAD_e removes more protons in the presence of
44 fumarate than in the presence of HCl at the same pH. All the above suggest that there are
45 additional effects by fumarate which might be associated with the intracellular GAD system
46 (GAD_i) or other acid resistance systems. We assessed the GAD_i output by looking at the
47 intracellular GABA pools which were not affected by fumarate. However, there are multiple
48 pathways (e.g. GABA shunt) that can affect GABA_i pools and we can not conclusively suggest
49 that GAD_i is affected. Furthermore, similarly to maleate, fumarate is able to eliminate *L.*
50 *monocytogenes* in biofilms under acidic conditions. Overall, fumarate is a good candidate for

51 *L. monocytogenes* decontamination and biofilm removal which is not toxic compared to the
52 toxic maleate.

53

54 **1. Introduction**

55 Foodborne illness is a significant public health problem both in the UK and globally. The
56 World Health Organisation (WHO) estimates that foodborne illness is responsible for 2.2
57 million deaths annually (Food standards agency, 2011). The majority of this illness in the UK
58 is caused by *Campylobacter* and *Norovirus* whilst most deaths are due to *Listeria*
59 *monocytogenes* and *Escherichia coli* infections. (Food standards agency, 2011)

60 Various strategies are employed to eliminate these pathogens in foods aiming to reduce the
61 incidence of foodborne illness. One such strategy is the addition of organic acids which have
62 been used for centuries to prevent the growth of pathogenic and spoilage bacteria (Ricke, 2003).

63 Organic acids are believed to affect microorganisms through diffusion of undissociated
64 molecules across the cell membrane followed by intracellular dissociation and release of
65 protons causing death or growth inhibition (Comes and Beelman, 2002; Lambert and Stratford,
66 1999; Podolak et al., 1996).

67 One of the well-known organic acids with antimicrobial activity is fumaric acid, which is a
68 food grade, dicarboxylic acid found widely in nature and active against a number of foodborne
69 pathogens including *E. coli*, *L. monocytogenes* and *Salmonella* sp. (Comes and Beelman, 2002;
70 Kim et al., 2009; Kondo et al., 2006; Miller and Kaspar, 1994; Pérez - Díaz and McFeeters,
71 2010; Podolak et al., 1996). In the EU and the US besides as an antimicrobial is also used as an
72 acidulant, and a flavour enhancer (Lee, 2014; Saltmarsh et al., 2013). Fumaric acid is regularly
73 used in various products including baked goods, confectionery, juices and dried powdered
74 foods as well as in animal feed (Lee, 2014). Fumaric acid is considered as one of the relatively
75 strongest among the weak organic acids, which however has low solubility in aqueous solutions

76 (Arnold et al., 2001; Roa Engel et al., 2013) while its salts are highly soluble (Zhou et al.,
77 2002). Based on its low dissociation constants ($pK_{a1} = 3.02$ and $pK_{a2} = 4.38$; Lohbeck et al.,
78 2000; Okuyama and Maskill, 2013; Szalka et al., 2013) it should be expected that fumarate has
79 low antimicrobial activity although this is not the case, if compared to other organic acids
80 including acetic and lactic acid (Podolak et al., 1996). This additional antimicrobial activity of
81 fumarate, beyond what could be explained by the intracellular dissociation theory of weak
82 acids, is normally attributed to unknown factors such as interference with metabolic activities
83 stress mechanisms or other cellular functions. Understanding these additional effects could
84 increase our knowledge and allow us to enhance the antimicrobial activity of these compounds
85 and consequently achieve higher levels of hygiene or develop novel and improved
86 antimicrobial regimes. Furthermore, it is important to understand in detail against which
87 organisms this additional effect occurs and what the mode of action is.

88 The current study focuses on the foodborne pathogen *L. monocytogenes*, a Gram positive,
89 facultative anaerobic bacterium that is ubiquitous in the environment (Posfay-Barbe and Wald,
90 2009) causing listeriosis, that mainly affects pregnant women, neonates and
91 immunocompromised individuals (Posfay-Barbe and Wald, 2009). The organism is capable of
92 surviving a wide range of environmental conditions and can grow under refrigeration
93 temperatures affecting ready-to-eat products (Liu et al., 2002; O'Driscoll et al., 1996). It is also
94 able to survive extreme acidic environments such as the stomach or acidic foods through the
95 use key mechanisms of acid resistance of which the main one is the GAD system (Davis et al.,
96 1996; Foster, 2004).

97 The GAD system converts glutamate to γ -amino butyric acid (GABA) with the removal of a
98 proton resulting in an increase in the intracellular pH (Cotter et al., 2001; Karatzas et al., 2012).
99 The architecture of the GAD system is highly variable and in *L. monocytogenes* it typically
100 comprises two antiporters, GadT1 and GadT2 and three decarboxylases GadD1, GadD2 and

101 GadD3. The GadD1T1 operon is typically associated with growth under mild acidic conditions,
102 the GadT2D2 promoting survival under extreme acidic conditions (Cotter et al., 2005) while
103 the GadD3 is the main part of the intracellular GAD system (GAD_i) utilising solely intracellular
104 glutamate to produce intracellular GABA which is catabolised to glutamate by the GABA shunt
105 (Cotter and Hill, 2003; Feehily et al., 2014; Feehily and Karatzas, 2013).

106 Given that fumarate has previously been described as an inhibitor of the *E. coli* GAD enzyme
107 (Fonda, 1972) we investigate here the antimicrobial activity of fumarate on *L. monocytogenes*
108 under acidic conditions and the possibility that this stems from effects on the GAD system and
109 possibly other aminoacid decarboxylase systems (Grobelny, 1995). Furthermore, we look at
110 the ability of fumarate to remove biofilms of *L. monocytogenes* and investigate further the
111 effects of the cis-isomer of fumarate, maleic acid on the GAD system which has also been
112 previously shown to affect it in *L. monocytogenes* (Paudyal et al., 2018).

113

114 **2. Materials and methods**

115 **2.1 Bacterial strains and growth conditions**

116 All strains (Table 1) were stored in 2 ml cryovials with a 7% dimethyl sulfoxide (DMSO) at -
117 80°C. *L. monocytogenes* 10403S and EGD-e were cultured onto Brain Heart Infusion (BHI)
118 agar (LABM, Lancashire UK) and *L. monocytogenes* LO28 onto Tryptic Soy Broth (Oxoid,
119 UK) supplemented with 5% yeast extract (TSBY; Oxoid, UK) and incubated at 37°C overnight.
120 Three colonies from each plate were transferred, with an inoculation loop, into BHI and TSBY
121 broth respectively in 10 ml Sterilin polystyrene tubes and incubated at 37°C with shaking (150
122 rpm) for 18 h. These overnight cultures were used to inoculate 20 ml cultures of the
123 corresponding media (1% inoculum) in 250 ml conical flasks which then were subsequently
124 incubated overnight at 37°C with shaking at (150 rpm) for 18 h.

125

126 **2.2 Calculation of undissociated acids using Henderson-Hasselbalch equation**

127 The percentage of undissociated acid present at pH 3 was determined using the Henderson-
128 Hasselbalch equation as adapted by Wemmenhove et al., (2016) and presented at Table 2

129
$$[\text{Undissociated acid}] = [\text{Total acid}] / 1 + 10^{(\text{pH}-\text{pK}_a)}$$

130

131 **2.3 Determination of minimum inhibitory concentration**

132 A range of concentrations (0, 1, 2, 4, 8, 16 and 32 mg/ml) of selected organic acids (Table 3)
133 were used in BHI inoculated at 1% with overnight culture of *L. monocytogenes* 10403S WT.
134 Cultures were then placed into 96 well plates and had their OD_{620nm} analysed over a 24 h period
135 using a Sunrise plate reader machine (Tecan, Mannedorf, Switzerland) operated by Magellan
136 software (Tecan, Mannedorf, Switzerland) with 20 min intervals between measurements. The
137 plates were kept at 37°C with shaking to assess the MIC.

138

139 **2.4 Survival under acidic conditions**

140 Survival experiments were undertaken with *L. monocytogenes* 10403S WT. Twenty ml
141 cultures were prepared in BHI, using stock cultures prepared as described previously, and
142 grown in 250 ml Erlenmeyer flasks at 37°C with agitation at 150 rpm. Acid challenge took
143 place with the addition of 8.6 mM fumaric acid, and a variety of organic acids (Table 3). Control
144 experiments were performed in the absence of any chemicals in overnight cultures. The pH of
145 the cultures was then adjusted to pH 3.0 using 1 M HCl.

146 One hundred µl samples were obtained prior to the acid challenge and every 20 min for 1 h and
147 placed in 900 µl Maximum Recovery Diluent (MRD; Oxoid Limited, Hampshire UK). Ten-
148 fold serial dilutions were prepared and 10 µl of each dilution were placed onto BHI agar plates
149 using the spot plate method and incubated at 37°C overnight. Subsequently, colonies were

150 counted to assess the cell concentration in the culture at each time point. All experiments were
151 performed in triplicate.

152

153 **2.5 Survival in the presence of sodium fumarate**

154 Following initial survival experiments, further survival experiments were performed focusing
155 on the effect of fumaric acid and its salt sodium fumarate on *L. monocytogenes* 10403S WT,
156 and its isogenic mutants $\Delta gadD1$, $\Delta gadD2$, $\Delta gadD3$, on EGD-e WT and its isogenic mutants
157 $\Delta gadD1$, $\Delta gadD2$, $\Delta gadD3$ and on LO28 WT with its isogenic mutants $\Delta gadD1$, $\Delta gadD2$,
158 $\Delta gadD1/2$.

159 Cultures were prepared in BHI or TSBY for LO28, using stock cultures, prepared as described
160 previously and grown in 250 ml Erlenmeyer flasks at 37°C with agitation at 150 rpm. Due to
161 the significantly differences in strain sensitivity, different concentrations of sodium fumarate
162 were used for 10403S (8.6 mM) and for EGD-e and LO28 (4.3 mM). Control cultures were
163 also prepared containing no additional antimicrobials. Subsequently, all *L. monocytogenes*
164 10403S cultures had their pH adjusted to 3.0 and *L. monocytogenes* EGD-e and LO28 to pH
165 3.3.

166 One hundred μ l samples were taken immediately prior to the acid challenge and every 20 or 5
167 min thereafter for 10403S or EGD-e and LO28 respectively. Samples were subsequently added
168 in 900 μ l MRD (Oxoid Limited, Hampshire UK) to prepare decimal serial dilutions and 10 μ l
169 of each dilution was plated onto BHI agar or TSBY agar respectively and incubated at 37°C
170 overnight. Following incubation, colonies were counted to assess the cell concentration at every
171 time point.

172

173 **2.6 GABAse assay**

174 GABase assay was used to determine the concentrations of intracellular (GABA_i) and
175 extracellular (GABA_e) GABA in 10403S and LO28 WT as described by O'Byrne et al., (2011).
176 *E. coli* K-12 samples were assayed following the same methodology with the modification of
177 the initial culture being grown in Lysogeny broth (LB LAB M, Lancashire, UK) supplemented
178 with 10 mM monosodium glutamate (MSG; Steinheim, Germany), GABA_e was quantified
179 according to Tsukatani et al. (2005) as modified by Karatzas (2010).

180

181 **2.7 GAD activity in protein lysates**

182 WT 10403S cultures were prepared in 20 ml BHI in 250 ml Erlenmeyer as described previously
183 by Paudyal et al (2018), while for LO28 and EGD-e 40 ml cultures were used. All cultures
184 were then transferred to 50 ml falcon tubes (VWR, Leighton Buzzard UK) with 10 µg/ml
185 chloramphenicol (Sigma-Aldrich, Steinheim, Germany). The samples were then centrifuged at
186 12,000 x g for 15 min and washed with a buffer solution, as described previously (Abrams et
187 al., 2008; Boura et al., 2016). Suspensions were then incubated at 37°C in an orbital shaker at
188 150 rpm (Gallenkamp, Germany).

189 A 2 ml cryovial (Sarstedt, Germany) was filled with 0.07g acid washed glass beads (< 106 µm
190 diameter Sigma-Aldrich, Steinheim, Germany) together with 1 ml cell suspension. The sample
191 was then agitated using a Mini-Beadbeater (Biospec, Bartesville, USA), thrice for 1 min,
192 followed by 1 min on ice. DNase I (Thermo Fisher Scientific, California, USA) was then added
193 up to 0.1% in the cell lysates and were then incubated at 37°C in an orbital shaker at 150 rpm
194 for 30 min. One ml sample was then transferred to an Eppendorff tube and centrifuged at 5,000
195 x g for 15 min. The supernatant was then transferred to an Eppendorf tube and the pellet
196 discarded. Subsequently, 100 µl of the supernatant was added to a pyridine hydrochloride buffer
197 (Fonda, 1972) supplemented with 30 mM MSG and with or without 20 mM sodium fumarate.
198 All samples were then adjusted to a pH of 4.5. The GABase assay was then used to assess

199 GABA levels. It had previously been established using standard concentrations of GABA, that
200 the presence of sodium fumarate does not affect the accuracy of this assay.

201

202 **2.8 Amino acid analysis by GC-MS**

203 As the activity of the GABase enzyme could be affected by the presence of other molecules,
204 GABA concentrations were also assessed in the supernatant or the bacterial lysates with the
205 use of gas chromatography mass spectrometry (Elmore et al., 2005). The method also assessed
206 the concentration of a wide range of aminoacids. Intra- and extra-cellular samples taken from
207 *L. monocytogenes* 10403S were assessed in the presence and absence of sodium fumarate.
208 Previous work by Paudyal et al. 2018 has shown that in similar conditions to those described
209 here, GABA levels quantified by GC-MS and GABase were always within $\pm 5\%$.

210

211 **2.9 Real-time PCR determination of GAD gene expression**

212 The transcription of the *gad* genes in the presence and absence of sodium fumarate was assessed
213 in *L. monocytogenes* 10403S WT (*gadD1*, *gadD2* and *gadD3*) using real time reverse
214 transcription-PCR (RT-PCR) as previously described by Karatzas et al., (2010). The
215 transcription of the antiporter-encoding genes (*gadT1* and *gadT2*) was not examined as it has
216 previously been demonstrated that it is similar to the corresponding glutamate decarboxylases
217 (*gadD1* and *gadD2*) belonging to the same corresponding operon (Karatzas et al., 2012).
218 Overnight cultures of *L. monocytogenes* 10403S WT grown for 24 h until stationary phase in
219 BHI were treated with 10 mM of either sodium fumarate or maleic acid for 40 min. Samples
220 were taken and prepared as previously described by Karatzas et al (2010). Relative expression
221 of the data was calculated as a ratio between expression of each of the target genes and the
222 expression of the 16S rRNA which was used as the reference gene for each cDNA sample.
223 Calculations were carried out following the advanced relative quantification settings of the

224 Light Cycler 480 SW 1.5.1 software programme, with PCR efficiencies of the primer pairs
225 gadD1F-gadD1R, gadD2F-gadD2R, gadD3F-gadD3R and 16SF-16SR being 2.12, 2.09, 2.03
226 and 2.27 respectively (Karatzas et al., 2010).

227

228 **2.10 Biofilm removal by sodium fumarate**

229 Biofilm formation was assessed using *L. monocytogenes* 10403S WT, EGD-e WT and LO28
230 WT overnight cultures grown in their corresponding BHI or TSBY agar as described previously
231 and then inoculated to 1% in a 2 ml of BHI broth or TSBY broth. The broth was mixed
232 thoroughly and placed in a 24-flat-bottom-well Corning Costar cell culture plate and sealed
233 using petrifilm. Following incubation at 37°C for 48h, the culture was removed and the wells
234 washed thrice with sterile water. Subsequently a fourth treatment was applied using 2.5 ml of
235 either water, 100 ppm free chlorine from calcium hyperchlorite, HCl (pH 2.4), HCl (pH 2.4)
236 with 25 mM sodium fumarate, AM (an organic acid disinfectant) at pH 2.4 and AM at pH 2.4
237 with 25 mM of sodium fumarate. The biofilm was exposed to these solutions in the well for 5
238 min and then the supernatants were discarded and wells were rinsed with 2.5 ml deionized
239 water. Subsequently, 500 µl MRD was placed in the well and the bottom of the well was scraped
240 using a 200 µl pipette tip for 30 s in a pattern covering the whole well bottom. This was repeated
241 4 times to provide a total volume of 2 ml which was serially diluted 10-fold and then 10 µl was
242 plated onto BHI or TSBY agar and incubated at 37°C for 24 h and then growth was assessed
243 (Ramírez et al., 2015). The impact of the treatments on the biofilm was assessed using the
244 following calculation (Hamilton, 2003; Heersink J., 2003; Ramírez et al., 2015).

245 Density = (Average count/Volume plated) * Dilution * Volume of MRD scraped into *(1/well surface
246 area).

247

248 **2.11 Statistical analysis**

249 In all cases all experiments were run in triplicate unless stated otherwise. Subsequently results
250 were assessed using paired Student t-tests. A P value below 0.05 indicated a statistically
251 significant result accompanied by an asterisk.

252

253 **3 Results**

254 **3.1 Calculation of the percentage of undissociated acid**

255 The pKa of sodium fumarate is low compared to the other acids tested (maleic acid, tartaric
256 acid and oxaloacetic acid; Table 2). This suggest that fumaric acid has a lower level of
257 undissociated acid (51.7 %) and therefore, lower antimicrobial activity.

258

259 **3.2 Growth in the presence of selected acids**

260 The MICs of a variety of organic acids on *L. monocytogenes* 10403S WT were assessed (Table
261 3). However, tartaric acid seemed to be the most bacteriostatic as it had the lowest MIC (14.9
262 mM). Sodium fumarate as a salt did not inhibit *L. monocytogenes* under the conditions of the
263 current experiment.

264

265 **3.3 Acid survival of *L. monocytogenes* 10403S, LO28 and EGD-e in the presence of** 266 **different organic acids.**

267 Under acidic conditions (pH 3), 8.6 mM sodium fumarate showed a significant bactericidal
268 effect on survival of the 10403S (Fig. 1A). Similar results were obtained with both EGD-e and
269 LO28 (Fig. 1B and 1C) assessed at pH 3.3 as more acid-sensitive than 10403S (Karatzas et al.,
270 2012). Based on previous work, it was expected that EGD-e might be the most sensitive strain,
271 however it displayed a similar response with LO28 to sodium fumarate (Fig. 1B and 1C;
272 Karatzas et al., 2012). Furthermore, we tested survival of 10403S against all compounds
273 mentioned in Table 1, and fumaric acid, sodium fumarate and maleic acid were the most

274 bactericidal at 8.6 mM and pH 3.3 with all other compounds hardly conferring any inactivation
275 (data not shown).

276

277 **3.4 Survival of *L. monocytogenes* 10403S LO28 and EGD-e and their isogenic mutants**
278 **under acidic conditions in the presence and absence of sodium fumarate.**

279 Once it was determined that sodium fumarate conferred the highest bactericidal activity, the
280 role of the GAD genes in the presence of sodium fumarate was assessed. In all cases, the
281 presence of sodium fumarate resulted in significant increase in the log reduction in all WT
282 strains and mutants (Fig. 2).

283 In 10403S the absence of sodium fumarate at pH 3 with HCl, minor log reductions in survival
284 occurred with $\Delta gadD2$ being the most sensitive. In the presence of 8.6 mM sodium fumarate
285 (pH 3) a significantly higher log reduction occurred for all strains while a similar trend
286 occurred, with all mutants except $\Delta gadD2$, behaving similarly to the WT (10403S WT, $\Delta gadD1$
287 and $\Delta gadD3$ showed 2.29, 1.99, and 2.56 log reduction of CFU/ml respectively; Fig. 2A).
288 $\Delta gadD2$ was the most sensitive strain and impacted more by the presence of sodium fumarate
289 (5.21 log reduction of CFU/ml).

290 In LO28 the effect of sodium fumarate showed a similar trend to 10403S, although this strain
291 was more sensitive and the effect was significantly more pronounced. Also, in this case the
292 addition of 4.3 mM sodium fumarate significantly affected survival at pH 3.3 and $gadD2$ was
293 also in this case the main determinant either with HCl alone or with sodium fumarate (Fig. 2B).

294 In the presence of 4.3 mM sodium fumarate (pH 3.3), LO28 WT and $\Delta gadD1$, showed a 4.80
295 and 5.11 log reduction CFU/ml respectively while that of $\Delta gadD2$ and $\Delta gadD1/2$ was higher
296 than the maximum of 6 logs that could be determined with this protocol (Fig. 2B).

297 In EGD-e the major difference compared to the other two strains was that removal of $gadD2$
298 did not result in increased sensitivity. In the presence of 4.3 mM sodium fumarate (pH 3.3),

299 EGD-e WT, $\Delta gadD1$, $\Delta gadD2$ and $\Delta gadD3$ showed 4.62, 2.65, 2.60 and 5.07 log reduction
300 CFU/ml respectively (Fig. 2C). In the presence of sodium fumarate, $\Delta gadD3$ was the most
301 sensitive strain, while $\Delta gadD1$ and $\Delta gadD2$ appeared significantly more resistant than the WT
302 probably due to the activation of another acid resistance mechanism. This trend had also been
303 observed in the presence of maleic acid the GAD_i system may play a survival role in the
304 presence of sodium fumarate (Paudyal et al., 2018).

305

306 **3.5 Extracellular GABA of *L. monocytogenes* 10403S, LO28 and *E. coli* K-12.**

307 To assess the possible influence of sodium fumarate on the activity of the GAD system an
308 examination of the effect of this compound on the levels of GABA_e was undertaken while the
309 cells were in stationary phase. In *L. monocytogenes* 10403S WT the presence of sodium
310 fumarate resulted in significant higher GABA_e levels from (4.11 mM GABA) compared to its
311 absence at pH 4.2 (2.01 mM GABA; 2.04-fold higher GABA levels P <0.05; paired T-test, Fig.
312 3A). Similarly, higher levels were obtained with all isogenic mutants except $\Delta gadD2$ (data not
313 shown). Similarly, LO28 WT also showed higher GABA_e levels in the presence of sodium
314 fumarate (2.89 mM GABA_e) compared to 1.24 mM GABA_e in its absence (2.33-fold higher
315 GABA_e levels Fig. 3B) although this result was not statistically significant. Also, its isogenic
316 mutants followed the same pattern (data not shown). These experiments were not performed
317 with EGD-e as it is defective in extracellular GABA production.

318 Finally, when a similar experiment was undertaken using *E. coli* K-12 WT (pH 4) in contrast
319 to the above, GABA_e levels were significantly lower (4.5 mM GABA_e) in the presence of
320 sodium fumarate compared to its absence (9.2mM GABA_e; 2.01-fold lower levels, P <0.05;
321 paired T-test; Fig. 3C). GABA_i was also examined however, no significant difference in any
322 of the strains tested was observed in the presence of sodium fumarate.

323

324 **3.6 10403S WT GAD activity in protein lysates**

325 The GAD activity of protein lysates was assessed by monitoring GABA production in the
326 presence of MSG (Sigma-Aldrich, Steinheim, Germany). The results indicate that, sodium
327 fumarate inhibited the GAD system activity in 10403S WT, resulting in reduced levels of
328 GABA from 3.4 mM to 2.7 mM (0.79-fold reduced levels; $P < 0.05$; Fig. 4). A similar protocol
329 was attempted with EGD-e and LO28 however, GABA levels were below the detection limit
330 of the GABase assay and despite protocol alterations in the pH, the buffer used, higher
331 glutamate supplementation or increasing the volume of culture utilised no improvement
332 occurred (Fig. 5).

333

334 **3.7 Real-time PCR determination of GAD gene expression.**

335 Real time quantitative Polymerase Chain Reaction (RT-qPCR) was used to quantify the
336 transcription of the *L. monocytogenes* 10403S WT GAD system genes in the presence of
337 sodium fumarate and its cis-isomer maleic which has previously been shown to inhibit the
338 listerial GAD system (Paudyal et al., 2018). Transcription of *gadD1* was very low and not
339 affected by the presence of sodium fumarate or maleic acid (Fig. 6A). In contrast, both sodium
340 fumarate and maleic acid resulted in a significant upregulation ($P < 0.05$) of the main component
341 of the GAD_e system, *gadD2* by 9.44- and 48.51-fold respectively (Fig. 6B). The latter gene also
342 showed the highest expression compared to the other two decarboxylases. Regarding *gadD3*,
343 expression was not affected by the presence of sodium fumarate although that of maleic acid
344 showed to result in an increase of 22.33-fold which however, was not statistically significant
345 (Fig. 6C).

346

347 **3.8 Biofilm formation**

348 The survival of *L. monocytogenes* biofilms was assessed after the application of various
349 antimicrobial treatments including 100 ppm chlorine and an acidic disinfectant (AM).

350 When *L. monocytogenes* 10403S was assessed, all treatments did not affect the survival in the
351 biofilm with the exception of those with 25 mM sodium fumarate, either alone at pH 2.4 (1M
352 HCl; 1.49 log reduction of CFU/cm²) or in combination with AM at the same pH (1.98 log
353 reduction of CFU/cm²; Fig. 7A).

354 In *L. monocytogenes* EGD-e three treatments achieved a statistically significant reduction of
355 CFU/ml. Those were the same ones that affected 10403S and the AM disinfectant alone (2.35
356 log reduction of CFU/cm²; Fig. 7B). Sodium fumarate alone at pH 2.4 resulted in a significant
357 3.72 log reduction of CFU/cm² while in combination with AM resulted in a 4.7 log reduction
358 of CFU/cm².

359 In *L. monocytogenes* LO28 all treatments resulted in a significant reduction in biofilm survival.
360 However, this was due to a lower variability between the replicates and overall the results were
361 similar to EGD-e with the exception of the AM treatment which seemed to be highly effective
362 against this strain. In this case also the two treatments with sodium fumarate were the most
363 effective along with 100 ppm chlorine (2.96 log reduction of CFU/cm²). Sodium fumarate alone
364 at pH 2.4 resulted in a significant 2.67 log reduction of CFU/cm² while in combination with
365 AM resulted in a 3.40 log reduction of CFU/cm². The AM disinfectant treatment at pH 2.4
366 resulted in a 2.23 log reduction of CFU/cm² while pH 2.4 alone resulted in 1.13 log reduction
367 of CFU/cm² (Fig. 7C).

368

369 **4. Discussion**

370 The antimicrobial effects of organic acids are mainly explained by the theory of passive
371 diffusion of undissociated molecules and their intracellular dissociation (Foster, 2004) which
372 also dictates that organic acids that dissociate more (higher Ka) are less antimicrobial, while

373 those that dissociate less (lower K_a) more antimicrobial. However, there are major deviations
374 to this rule (Ricke, 2003) suggesting the existence of additional effects which are highly
375 important for our understanding of the mode of action of various organic acids and the
376 behaviour of microorganisms in foods and many other environments.

377 The present work focuses on fumaric acid which is widely used in foods while it is present in
378 all foods, all living organisms and various environments as it is a key component of the TCA
379 cycle. It is one of the relatively strongest among the weak organic acids and based on its low
380 pKa of 3.02 it should have low antimicrobial activity as it dissociates more (Table 2).
381 **Therefore**, as predicted, the MIC of fumaric acid was among the highest (34 mM) with only
382 oxaloacetic acid (60.5 mM) and sodium fumarate having higher MICs (Table 3).

383 However, in contrast to the above fumarate showed high bactericidal activity against all three
384 strains of *L. monocytogenes* used (Fig. 1A, B & C). This is a clear deviation from the behaviour
385 that would be predicted by the intracellular dissociation theory of weak acids suggesting
386 additional antimicrobial effects which however, only occur at highly acidic conditions. This
387 high antimicrobial activity of fumarate has been noted previously (Chikthimmah et al., 2003;
388 Comes and Beelman, 2002; Podolak et al., 1996; Kondo et al., 2006) and our aim was to
389 identify the complementary mode of action of fumarate which does not stem from the theory
390 of intracellularly dissociation of organic acids.

391 Fumarate is an inhibitor of the *E. coli* GAD system (Fonda, 1972) and based on this, an obvious
392 hypothesis to explain the above effects, is the inhibition of the *L. monocytogenes* GAD system.
393 To investigate this hypothesis and in parallel identify which component of the GAD system
394 might be affected, deletion mutants in GAD decarboxylase genes for all three strains of *L.*
395 *monocytogenes* were used. In 10403S, removal of *gadD1* and *gadD3* resulted in similar
396 population reduction when compared to the WT in presence of sodium fumarate. However,
397 removal of *gadD2* caused the greatest death both in the presence and in the absence of sodium

398 fumarate as expected since most strains rely on the GadD2 for the operation of the dominant
399 GAD_e system (Fig. 2A; Karatzas, Brennan et al. 2010). Similarly, to 10403S, in LO28 the
400 removal of the key *gadD2* significantly reduced survival under acidic conditions (Fig. 2B).
401 However, EGD-e does not possess a GAD_e system associated with GadD2, and only utilises
402 the GAD_i which is mediated by GadD3 (Feehily et al., 2014; Karatzas et al., 2012). In this
403 strain, Δ *gadD3* was the most sensitive either in the presence or absence of sodium fumarate,
404 without statistical significance though, whereas *gadD1* and *gadD2* removal significantly
405 increased resistance but only in the presence of sodium fumarate (Fig. 2C). Overall inactivation
406 was much higher in the presence of sodium fumarate and more exaggerated for the more
407 sensitive mutants that showed higher inactivation in the absence of sodium fumarate.
408 Furthermore, it should be noted that removal of *gadD1* from all strains did not significantly
409 influence their ability to survive acidic conditions. This gene has previously been linked to
410 increased growth under mild acidic conditions based on observations with WT strains that do
411 not possess it (e.g. serotype 4b strains; Cotter et al., 2005). However, mutants in this operon
412 have never been shown to possess an acid sensitive phenotype, or evidence showing GadD1T1
413 exporting GABA neither in the literature or in our experiments, raising questions over its
414 function as a glutamate decarboxylase. Overall, GadD2 and GadD3, depending on the strain,
415 were significant for survival against sodium fumarate under acid conditions (Fig. 2A, B & C).
416 Sodium fumarate resulted in significant increase in GABA_e exported by *L. monocytogenes*
417 10403S (2.04-fold increase) at pH 4.2 (P <0.05; Fig. 3A). Interestingly, this increased GAD_e
418 output did not confer increased survival under acidic conditions (Fig. 2A). This is highly
419 interesting as each GABA molecule exported, removes one intracellular proton and based on
420 the above, GAD_e removes twice more protons in the presence than in the absence of fumarate,
421 but instead of conferring higher resistance the opposite occurs (Fig. 2A & B). A similar trend
422 was observed with LO28 although it was not statistically significant (P >0.05; Fig. 3B). It

423 should be stated that as we have shown previously, maleic acid, the cis-isomer of fumaric acid
424 works in an opposite fashion than fumarate, as it reduces GAD_e output (Paudyal et al., 2018).
425 However, in contrast to *L. monocytogenes*, when we challenged *E. coli* K12, sodium fumarate
426 resulted in a significant -2.01-fold decrease in GABA_e export (P <0.05; Fig. 3C). The latter was
427 expected as fumarate is an inhibitor of *E. coli* GAD enzyme (Fonda, 1972) and it demonstrates
428 a different GAD system behaviour between these two organisms. Further work should
429 investigate if this is a different feature between Gram-positive and Gram-negative bacteria.
430 To further assess the effect of sodium fumarate on the GAD system, we measured GABA_i
431 levels, and found no significant effect of sodium fumarate. At first glance, this might suggest
432 no effect of sodium fumarate however, GABA_i levels are also affected by its metabolic flux
433 through the GABA shunt pathway and therefore the above results are not conclusive.
434 Surprisingly, we observed that sodium fumarate significantly inhibited GAD activity as
435 measured through GABA levels in *L. monocytogenes* 10403S cell lysates (P <0.05; Fig. 4).
436 This coincides with its role as GAD inhibitor in *E. coli* (Fonda, 1972) and in plants (Ohno and
437 Okunuki, 1962). We further investigated these inhibitory effects of sodium fumarate in lysates
438 of LO28 and EGD-e but unfortunately, we were not able to get measurable GABA levels (Fig.
439 5) even in the absence of sodium fumarate, despite various protocol modifications (usage of
440 higher cell numbers, higher levels of glutamate, different buffer pH values). This might be
441 related to lower GAD activity or a different optimal pH of the GAD enzymes in these strains.
442 Further we looked at the effect of fumarate on transcription of GAD genes *gadD1*, *gadD2* and
443 *gadD3* including maleic acid which is a cis-isomer of fumarate and we have previously shown
444 that reduces GAD output and activity in *L. monocytogenes* enhancing its acid sensitivity
445 (Paudyal et al., 2018). RT-qPCR showed no effect of fumarate or maleate on *gadD1* and *gadD3*
446 (Fig. 6A and C) however, *gadD2*, the key component of GAD_e system in *L. monocytogenes*
447 10403S WT (Cotter et al., 2001; Cotter et al., 2005) was upregulated by sodium fumarate and

448 even more by sodium maleate ($P < 0.05$, paired t-test, Fig. 6 B). The above suggest that *L.*
449 *monocytogenes* tries to counteract the inhibitory effects of fumarate and maleate on GAD
450 activity by increasing *gadD2* transcription and the final result of these opposing actions in the
451 case of fumarate is increased GAD_e system output which however, does not enhance acid
452 resistance. However, in the case of maleate, higher increase in *gadD2* transcription is unable to
453 increase GAD_e system output (main difference with fumarate) but similarly to fumarate, the
454 acid resistance is reduced.

455 The explanation for the antilisterial effects of fumarate might lie in the effects on the GAD_i
456 system (Feehily and Karatzas, 2013), or other possible effects on other acid resistance systems
457 or on cell metabolism that in its turn could affect acid resistance. For example, fumarate is
458 highly antimicrobial against organisms such as *Salmonella* (Kondo et al., 2006) that lack GAD
459 system suggesting these additional effects (Park et al., 1996). To assess this, we first looked at
460 the aminoacid profile in presence or absence of fumarate and the only difference found was the
461 increased GABA_e levels in presence of fumarate confirming the GABase results. This suggests
462 that other aminoacid decarboxylase systems are possibly not affected and the above effects of
463 fumarate are on GAD_i system or possibly another non-amino acid decarboxylase system.

464 Furthermore, the ability of sodium fumarate to act on cells in a biofilm was examined. It has
465 previously been shown that maleic acid can act on biofilms of *L. monocytogenes* (Paudyal et
466 al., 2018) and *E. faecalis* (Ferrer-Luque et al., 2010). Due to these properties it has been
467 suggested that maleic acid could be an effective alternative to the more toxic EDTA commonly
468 used to remove biofilms from the oral cavity and dental equipment (Ballal et al., 2009; Ferrer-
469 Luque et al., 2010). However, fumarate has no toxicity and therefore further work could
470 investigate other potential applications. Our results showed the striking ability of sodium
471 fumarate (25 mM) to eliminate cells of three different strains of *L. monocytogenes* in a biofilm,
472 which was significantly higher than that of hypochlorite and a commonly used organic acid

473 disinfectant AM at pH 2.4 (Fig. 7). Furthermore, the addition of fumarate together with the AM
474 disinfectant increased significantly the ability of the disinfectant to eliminate cells in biofilm.
475 Our results also show that the more acid resistant strain 10403S survived the treatments better
476 than the other two acid sensitive strains (EGD-e and LO28), underpinning the important role
477 of acid resistance and GAD system in survival in a biofilm. Furthermore, we also found that
478 LO28 was highly sensitive to chlorine. This coincides with previous reports suggesting a high
479 variation in resistance to chlorine-based sanitisers among different strains (Brackett, 1987;
480 Jacquet and Reynaud, 1994) and that mixed culture strains of *L. monocytogenes* are better able
481 to resist chlorine treatments (Vaid et al., 2010). Our results suggest that fumarate has a great
482 potential for removal of biofilms of *L. monocytogenes* while it is also nontoxic.

483

484 Conclusions

485 Overall, we investigated the effect of fumarate on *L. monocytogenes* under acidic conditions
486 showing that although it is a GAD_e inhibitor, the bacterium is able to counteract this with
487 increased transcription, being able to increase its overall GAD_e output, which however does
488 not translate into increased acid resistance. We also show that there is a difference between
489 fumarate and maleate as although the first increases GAD output, the latter reduces it, but both
490 significantly enhance death under acidic conditions. The antimicrobial activity of fumarate
491 might be related to reduced GAD_i or other systems. Further work is required to elucidate the
492 full extent of the antimicrobial activity of fumarate on *L. monocytogenes* and other organisms.
493 Such work will allow us to successfully eliminate this pathogen in food and food preparation
494 environments but also explain its behaviour in environments where fumarate is present.

495

496 Acknowledgements

497 The authors would like to thank all colleagues at the Microbiology research team and the
498 Department of Food and Nutritional Sciences, University of Reading. The help of Foteini Pavli
499 and Prof. Nychas (Agricultural University of Athens, Greece) on setting up the biofilm
500 experiments has been invaluable. The work was supported by a Biotechnology and Biological
501 Sciences Research Council doctoral training partnership (BBSRC-DTP Grant number
502 RE04907), awarded to Dr. K.A.G. Karatzas and funds from the University of Reading and by
503 AgriCoat NatureSeal Ltd.

504

505

506

507

508 **References**

- 509 Abrams, F., Wan-Lin, S., Wiedmann, M., Boor, K.J., Coote, P.J., Botting, C.H., Karatzas,
510 K.A.G., O'Byrne, C.P., (2008). Proteomic analysis of a *Listeria monocytogenes*
511 mutant lacking sigma B (σ^B) identify new components of the σ^B regulon and highlight
512 a role for sigma B in the utilization of glycerol. *Appl. Environ. Microbiol.* 74, 594-
513 604.
- 514
- 515 Arnold, C.N., McElhanon, J., Lee, A., Leonhart, R., Siegele, D.A., (2001). Global analysis of
516 *Escherichia coli* gene expression during the acetate-induced acid tolerance response.
517 *J. Bacteriol.* 183.
- 518
- 519 Ballal, N.V., Kandian, S., Mala, K., Bhat, K.S., Acharya, S., (2009). Comparison of the
520 efficacy of maleic acid and ethylenediaminetetraacetic acid in smear layer removal
521 from instrumented human root canal: a scanning electron microscopic study. *J.*
522 *Endod.* 35, 1573-1576.
- 523
- 524 Boura, M., Keating, C., Royet, K., Paudyal, R., O'Donoghue, B., O'Byrne, C.P., Karatzas,
525 K.A., (2016). Loss of SigB in *Listeria monocytogenes* strains EGD-e and 10403S
526 confers hyperresistance to hydrogen peroxide in stationary phase under aerobic
527 conditions. *Appl. Environ. Microbiol.* 82, 4584-4591.
- 528
- 529 Brackett, R., (1987). Antimicrobial effect of chlorine on *Listeria monocytogenes*. *J. Food*
530 *Prot.* 50, 999-1003.
- 531
- 532 Chikthimmah, N., Laborde, L., Beelman, R., (2003). Critical factors affecting the destruction
533 of *Escherichia coli* O157: H7 in apple cider treated with fumaric acid and sodium
534 benzoate. *J. Food Sci.* 68, 1438-1442.

535
536 Comes, J.E., Beelman, R.B., (2002). Addition of fumaric acid and sodium benzoate as an
537 alternative method to achieve a 5-log reduction of *Escherichia coli* O157: H7
538 populations in apple cider. *J. Food Prot.* 65, 476-483.
539
540 Cotter, P.D., Gahan, C.G., Hill, C., (2001). A glutamate decarboxylase system protects
541 *Listeria monocytogenes* in gastric fluid. *Mol. Microbiol.* 40, 465-475.
542
543 Cotter, P.D., Hill, C., (2003). Surviving the acid test: Responses of Gram-positive bacteria to
544 low pH. *Microbiol. Mol. Biol. Rev.* 67, 429-429.
545
546 Cotter, P.D., Ryan, S., Gahan, C.G., Hill, C., (2005). Presence of GadD1 glutamate
547 decarboxylase in selected *Listeria monocytogenes* strains is associated with an ability
548 to grow at low pH. *Appl. Environ. Microbiol.* 71, 2832-2839.
549
550 Davis, M.J., Coote, P.J., O'Byrne, C.P., (1996). Acid tolerance in *Listeria monocytogenes*: the
551 adaptive acid tolerance response (ATR) and growth-phase-dependent acid resistance.
552 *Microbiol.* 142 (Pt 10), 2975-2982.
553
554 Elmore, J.S., Koutsidis, G., Dodson, A.T., Mottram, D.S., Wedzicha, B.L., (2005).
555 Measurement of acrylamide and its precursors in potato, wheat, and rye model
556 systems. *J. Agric. Food Chem.* 53, 1286-1293.
557
558 Feehily, C., Finnerty, A., Casey, P.G., Hill, C., Cormac, G.M.G., O'Byrne, C.P., Karatzas, K.-
559 A.G., (2014). Divergent evolution of the activity and regulation of the glutamate
560 decarboxylase systems in *Listeria monocytogenes* EGD-e and 10403S: Roles in
561 virulence and acid tolerance: *PLOS* 9 (11), e112649.
562
563 Feehily, C., Karatzas, K.A.G., (2013). Role of glutamate metabolism in bacterial responses
564 towards acid and other stresses. *J. Appl. Microbiol.* 114, 11-24.
565
566 Feehily, C., O'Byrne, C.P., Karatzas, K.A.G., (2013) Functional γ -Aminobutyrate shunt in
567 *Listeria Monocytogenes*: role in acid tolerance and succinate biosynthesis. *Appl.*
568 *Environ. Microbiol.*(79), 74-80.
569
570 Ferrer-Luque, C.M., Arias-Moliz, M.T., Gonzalez-Rodriguez, M.P., Baca, P., (2010).
571 Antimicrobial activity of maleic acid and combinations of cetrime with chelating
572 agents against *Enterococcus faecalis* biofilm. *J. Endod.* 36, 1673-1675.
573
574 Fonda, M.L., (1972). Glutamate decarboxylase. Substrate specificity and inhibition by
575 carboxylic acids. *Biochemistry (Easton)* 11, 1304-1309.
576
577 Food Standards Agency, (2011). Foodborne disease strategy 2010-15, in: agency, F.s. (Ed.).
578
579 Foster, J.W., (2004). *Escherichia coli* acid resistance: tales of an amateur acidophile. *Nat.*
580 *Rev. Microbiol.* 2, 898-907.
581
582 Grobelny, J., (1995). N.m.r. study of maleate (cis)—fumarate (trans) isomerism in
583 unsaturated polyesters and related compounds. *Polymer.* 36, 4215-4222.
584

- 585 Heersink J., (2003). Basic biofilm analytical methods. In: Hamilton, M., Heersink, J.,
586 Buckingham-Meyer, K. Goeres, D. (Ed 4th.) The biofilms laboratory: Step-by-step
587 protocols for experimental design, analysis, and data interpretation. Bozeman: Cytergy
588 Publishing, pp. 16-23.
- 589
- 590 Jacquet, C., Reynaud, A., (1994). Differences in the sensitivity to eight disinfectants of
591 *Listeria monocytogenes* strains as related to their origin (short note). *Int. J. Food*
592 *Microbiol.* 22, 79-83.
- 593
- 594 Karatzas, K.-A.G., Suur, L., O'Byrne, C.P., (2012). Characterization of the intracellular
595 glutamate decarboxylase system: Analysis of its function, transcription, and role in the
596 acid resistance of various strains of *Listeria monocytogenes*. *Appl. Environ.*
597 *Microbiol.* 78, 3571-3579.
- 598
- 599 Karatzas, K.A., Brennan, O., Heavin, S., Morrissey, J., O'Byrne, C.P., (2010). Intracellular
600 accumulation of high levels of gamma-aminobutyrate by *Listeria monocytogenes*
601 10403S in response to low pH: uncoupling of gamma-aminobutyrate synthesis from
602 efflux in a chemically defined medium. *Appl. Environ. Microbiol.* 76, 3529-3537.
- 603
- 604 Kim, Y.J., Kim, M.H., Song, K.B., (2009). Efficacy of aqueous chlorine dioxide and fumaric
605 acid for inactivating pre-existing microorganisms and *Escherichia coli* O157:H7,
606 *Salmonella* Typhimurium, and *Listeria monocytogenes* on broccoli sprouts. *Food*
607 *Control.* 20, 1002-1005.
- 608
- 609
- 610 Kondo, N., Murata, M., Isshiki, K., (2006). Efficiency of sodium hypochlorite, fumaric acid,
611 and mild heat in killing native microflora and *Escherichia coli* O157: H7, *Salmonella*
612 Typhimurium DT104, and *Staphylococcus aureus* attached to fresh-cut lettuce. *J.*
613 *Food Prot.* 69, 323-329.
- 614
- 615 Lambert, R., Stratford, M., (1999). Weak-acid preservatives: modelling microbial inhibition
616 and response. *J. Appl. Microbiol.* 86, 157-164.
- 617
- 618 Lee, B.H., (2014). Fundamentals of Food Biotechnology 2nd edn. John Wiley & Sons,
619 Incorporated, New York, NY, 1413 pp.
- 620
- 621 Liu, S., Graham, J.E., Bigelow, L., Morse, P.D., Wilkinson, B.J., (2002). Identification of
622 *Listeria monocytogenes* genes expressed in response to growth at low temperature.
623 *Appl. Environ. Microbiol.* 68, 1697-1705.
- 624
- 625 Lohbeck, K., Haferkorn, H., Fuhrmann, W., Fedtke, N., (2000). Maleic and fumaric acids.
626 Ullmann's encyclopedia of industrial chemistry Wiley-VCH, Weinheim.
- 627
- 628 Miller, L.G., Kaspar, C.W., (1994). *Escherichia coli* O157: H7 acid tolerance and survival in
629 apple cider. *J. Food Prot.* 57, 460-464.
- 630
- 631 O'Byrne, C.P., Feehily, C., Ham, R., Karatzas, K.A.G., (2011). A modified rapid enzymatic
632 microtiter plate assay for the quantification of intracellular γ -aminobutyric acid and
633 succinate semialdehyde in bacterial cells. *J. Microbiol. Meth.* 84, 137-139.
- 634

- 635 O'Driscoll, B., Gahan, C.G., Hill, C., (1996). Adaptive acid tolerance response in *Listeria*
636 *monocytogenes*: isolation of an acid-tolerant mutant which demonstrates increased
637 virulence. *Appl. Environ. Microbiol.* 62, 1693-1698.
638
- 639 Ohno, M., Okunuki, K., (1962). Studies on the inhibition of glutamic acid decarboxylase of
640 squash. *J. Biochem.* 51, 313-316.
641
- 642 Okuyama, T., Maskill, H., (2013). Organic chemistry: a mechanistic approach. Oxford
643 University Press.
644
- 645 Park, Y.K., Bearson, B., Bang, S.H., Bang, I.S., Foster, J.W., (1996). Internal pH crisis, lysine
646 decarboxylase and the acid tolerance response of *Salmonella Typhimurium*. *Mol*
647 *Microbiol.* 20, 605-611.
648
- 649 Paudyal, R., Barnes, R.H., Karatzas, K.A.G., (2018). A novel approach in acidic disinfection
650 through inhibition of acid resistance mechanisms; Maleic acid-mediated inhibition of
651 glutamate decarboxylase activity enhances acid sensitivity of *Listeria monocytogenes*.
652 *Food Microbiol.* 69, 96-104.
653
- 654 Pérez-Díaz, I., McFeeters, R., (2010). Preservation of acidified cucumbers with a natural
655 preservative combination of fumaric acid and allyl isothiocyanate that target lactic
656 acid bacteria and yeasts. *J. Food Sci.* 75 (4). doi: 10.1111/j.1750-3841.2010.01587.x.
657
- 658 Podolak, R., Zayas, J., Kastner, C., Fung, D., (1996). Inhibition of *Listeria monocytogenes*
659 and *Escherichia coli* O157: H7 on beef by application of organic acids. *J. Food Prot.*
660 59, 370-373.
661
- 662 Posfay-Barbe, K.M., Wald, E.R., (2009). Listeriosis. *Semin. Fetal Neonatal Med.* 14, 228-
663 233.
664
- 665 Ramírez, M.D.F., Smid, E.J., Abee, T., Groot, M.N.N., (2015). Characterisation of biofilms
666 formed by *Lactobacillus plantarum* WCFS1 and food spoilage isolates. *Int. J. Food*
667 *Microbiol.* 207, 23-29.
668
- 669 Ricke, S., (2003). Perspectives on the use of organic acids and short chain fatty acids as
670 antimicrobials. *Poult. Sci.* 82, 632-639.
671
- 672 Roa Engel, C.A., ter Horst, J.H., Pieterse, M., van der Wielen, L.A., Straathof, A.J., (2013).
673 Solubility of fumaric acid and its monosodium salt. *Ind. Eng. Chem. Res.* 52, 9454-
674 9460.
675
- 676 Saltmarsh, M., Barlow, S., Richardson, V., Robin, A.-L., Jukes, D.J., (2013). Essential guide
677 to food additives. Royal Society of Chemistry, Cambridge.
678
- 679 Szalka, M., Rokaszewski, E., Kaczmarek, K., (2013). Kinetics of hydrolysis of bisoprolol
680 hemifumarate in aqueous acidic solutions. *Int. J. Chem. Kinet.* 45, 744-754.
681
- 682 Vaid, R., Linton, R.H., Morgan, M.T., (2010). Comparison of inactivation of *Listeria*
683 *monocytogenes* within a biofilm matrix using chlorine dioxide gas, aqueous chlorine
684 dioxide and sodium hypochlorite treatments. *Food Microbiol.* 27, 979-984.

685
686 Wemmenhove, E., van Valenberg, H.J.F., Zwietering, M.H., van Hooijdonk, T.C.M., Wells-
687 Bennik, M.H.J., (2016). Minimal inhibitory concentrations of undissociated lactic,
688 acetic, citric and propionic acid for *Listeria monocytogenes* under conditions relevant
689 to cheese. *Food Microbiol.* 58, 63-67.
690
691 Zhou, Y., Du, J., Tsao, G., (2002). Comparison of fumaric acid production by *Rhizopus*
692 *oryzae* using different neutralizing agents. *Bioprocess Biosyst. Eng.* 25, 179-181.
693

694

695

696

697

698

699 **Legends to the figures**

700 **Fig. 1.** Survival of *L. monocytogenes* (A) 10403S WT in the presence (black circles) and
701 absence (black triangles) of 8.6 mM sodium fumarate adjusted to pH 3 using 1 M HCl (B)
702 EGD-e WT and (C) LO28 WT in the presence (black circles) and absence (black triangles) of
703 4.3 mM sodium fumarate at pH 3.3 using 1 M HCl. Asterisks represent statistically significant
704 result ($P < 0.05$ paired student T-test) while D.L denotes detection limit of the experimental
705 setup.

706

707 **Fig. 2.** Survival of *L. monocytogenes* and its GAD mutants in the (A) 10403S background in
708 the presence and absence of 8.6 mM of sodium fumarate adjusted to pH 3 for 60 min, in the (B)
709 LO28 and (C) EGD-e background in the presence and absence of 4.3 mM of sodium fumarate
710 adjusted to pH 3.3 for 15 min. Adjustment of pH was done using 1 M HCl. Asterisks represent
711 statistically significant result as assessed with paired student T-test ($P < 0.05$) and M.L. denotes
712 the maximum log reduction could be recorded with the current protocol.

713

714 **Fig. 3** GABA_e levels of overnight cultures grown to stationary phase (~18 h at 37°C) with
715 shaking in the presence or absence of 10 mM sodium fumarate (SF) for (A) *L. monocytogenes*
716 10403 WT at pH 4.2, (B) *L. monocytogenes* LO28 WT at pH 4.2 and (C) *E. coli* K-12 WT at
717 pH 4. pH was adjusted with the addition of 1 M HCl. Asterisk represents statistically significant
718 result. $P < 0.05$ paired student T-test.

719

720

721 **Fig. 4.** GAD activity in cell lysates of *L. monocytogenes* 10403S WT cells grown overnight
722 until stationary phase (~18h) at 37°C with agitation (150 rpm) in the presence or absence of 20
723 mM sodium fumarate at pH 4.2. Lysates were prepared and then levels of GAD activity were
724 assessed using GC-MS. Asterisk represents statistically significant result. P <0.05 paired
725 student T-test.

726

727 **Fig. 5** GAD activity in cell lysates of *L. monocytogenes* 10403S WT, EGD-e WT and LO28
728 WT cells grown overnight until stationary phase (~18h) at 37°C with agitation (150 rpm).
729 Lysates were produced and then levels of GAD activity were assessed using the GABase
730 enzymatic assay. Asterisks represents statistically significant result. P <0.05 paired student T-
731 test.

732

733 **Fig. 6** Expression of (A) *gadD1* (B) *gadD2* and (C) *gadD3* gene in *L. monocytogenes* 10403S
734 WT in the absence or presence of 10 mM sodium fumarate or 10 mM maleic acid. Relative
735 expression of each gene was calculated by comparing expression relative to 16S rRNA gene in
736 each strain. Numbers above the bars represent fold difference in relative expression compared
737 to control. Markers represent an average of triplicate measurements and error bars represent
738 standard deviations. Asterisks * denote statistical significant difference compared to the control
739 (P <0.05 paired student T-test).

740

741 **Fig. 7** Survival of cells in biofilms of *L. monocytogenes* (A) 10403S WT (B) *L. EGD-e* WT
742 and (C) LO28 WT following no treatment (water) or treatment with an acidic disinfectant (AM),
743 AM together with 25 mM sodium fumarate (SF), HCl and HCl together with 25 mM SF. All
744 treatments were at pH 2.4. Asterisks represent statistically significant difference between no
745 treatment and a treatment (P <0.05; paired student T-test).

746

747

748

749

750

751

752

753

754

755

756

757

758

759

760

761 **Tables**

762

763

Table 1. list of strains used in these experiments

Strains	Relevant properties	Source
<i>L. monocytogenes</i> 10403S	Serotype 1/2a, WT	Karatzas et al., 2010
<i>L. monocytogenes</i> 10403S Δ <i>gadD1</i>	10403S with <i>gadD1</i> deleted	Karatzas et al., 2010
<i>L. monocytogenes</i> 10403S Δ <i>gadD2</i>	10403S with <i>gadD2</i> deleted	Karatzas et al., 2010
<i>L. monocytogenes</i> 10403S Δ <i>gadD3</i>	10403S with <i>gadD3</i> deleted	Karatzas et al., 2010
<i>L. monocytogenes</i> EGD-e	Serotype 1/2a, WT	Feehily et al., 2013
<i>L. monocytogenes</i> EGD-e Δ <i>gadD1</i>	EGD-e with <i>gadD1</i> deleted	Feehily et al., 2013
<i>L. monocytogenes</i> EGD-e Δ <i>gadD2</i>	EGD-e with <i>gadD2</i> deleted	Feehily et al., 2013
<i>L. monocytogenes</i> EGD-e Δ <i>gadD3</i>	EGD-e with <i>gadD3</i> deleted	Feehily et al., 2013
<i>L. monocytogenes</i> LO28	Serotype 1/2c, WT	Cotter et al., 2001
<i>L. monocytogenes</i> LO28 Δ <i>gadD1</i>	LO28 with <i>gadD1</i> deleted	Cotter et al., 2001
<i>L. monocytogenes</i> LO28 Δ <i>gadD2</i>	LO28 with <i>gadD2</i> deleted	Cotter et al., 2001
<i>L. monocytogenes</i> LO28 Δ <i>gadD1/2</i>	LO28 with <i>gad D1/2</i> deleted	Cotter et al., 2001

<i>E. coli</i> K-12	WT	KEIO collection
764		
765		
766		
767		
768		
769		
770		
771		
772		
773	Table 2. Percentage of undissociated acids at pH 3.	

Compound	pKa ₁ 1	pKa ₂ 2	pKa ₃ 3	%undissociated /total acid (pKa ₁)	%undissociated /total acid (pKa ₂)	%undissociated/ total acid (pKa ₃)
Maleic acid	1.9	6.07		7.35	99.91	
Fumaric acid	3.03	4.44		51.72	96.49	
Sodium fumarate	3.55			78.01		
Pimlic acid	4.71	5.58		98.08	99.73	
Valeric acid	4.82			98.50		
Adipic acid	4.43	5.41		96.41	99.61	
Glutaric acid	4.34	5.22		95.62	99.40	
Malic acid	3.4	5.44		71.52	99.63	
Citric acid	3.13	4.76	6.39	57.42	98.29	99.95
Tartaric acid	2.98	4.34		48.84	95.62	
Oxaloacetic acid	2.22	3.89		14.23	88.58	

Alpha ketoglutaric acid	3.08	54.59
Valeric acid	4.82	98.50
Levulinic acid	4.59	97.49

774

775

776

777

778

779

780

781

782

Table 3. MICs of compounds tested

Potential inhibitor	MIC
Maleic acid	34.4 mM
Fumaric acid	34.4 mM
Sodium fumarate	Above solubility threshold 0.22 mg/ml
Glutaric acid	30mM
Pimelic acid	24.9 mM
Adipic acid	27.3 mM
Malic acid	14.9 mM
Citric acid	20.8 mM
Tartaric acid	26.6 mM
Oxaloacetic acid	60.5 mM
α -Ketoglutaric acid	27.3 mM
Valeric acid	13 mM
Levulinic acid	60 mM

783