

# *Field validation of senesced banana leaf extracts for trapping banana weevils on smallholder banana/plantain farms*

Article

Accepted Version

Abagale, S. A., Braimah, H., Osafo-Acquaah, S., Powers, S. J., van Emden, H. F., Birkett, M. A., Pickett, J. A., Sanda, U. L. and Vuts, J. (2020) Field validation of senesced banana leaf extracts for trapping banana weevils on smallholder banana/plantain farms. *Journal of Applied Entomology*. ISSN 0931-2048 doi: <https://doi.org/10.1111/jen.12838> Available at <https://centaur.reading.ac.uk/93985/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1111/jen.12838>

Publisher: Wiley-Blackwell

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

**CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

**Table 1** Total numbers of adult banana weevils, *Cosmopolites sordidus*, captured at five field locations in Ashanti region, Ghana, using aggregation pheromone, senesced banana leaf palm alcohol extract, pseudostem, and combinations thereof, in type TAL and Voltic traps. Overall total and percentage of overall total capture for two groups (with/without pseudostem) of the five treatments are also shown.

<b>Field (Type of trap)</b>	<b>Duration of experiment (Weeks)</b>	<b>Aggregation pheromone plus banana leaf extract</b>	<b>Palm alcohol extract of banana leaf</b>	<b>Aggregation pheromone</b>	<b>Pseudostem plus banana leaf extract</b>	<b>Pseudostem</b>
1 (TAL)	12	151	20	109	10	5
2 (Voltic)	7	87	19	32	17	7
3 (TAL)	12	180	30	86	29	9
4 (Voltic)	12	433	51	197	15	7
5 (Voltic)	5	24	3	11	4	1
<b>Total</b>		<b>875</b>	<b>123</b>	<b>435</b>	<b>75</b>	<b>29</b>
<b>Overall total</b>		<b>1433</b>			<b>104</b>	
<b>Percent weevil capture (%)</b>		<b>61.1</b>	<b>8.5</b>	<b>30.4</b>	<b>72.1</b>	<b>27.9</b>

**Table 2** Mean ( $\pm$  SE) number of adult banana weevils, *Cosmopolites sordidus*, caught per week in field trapping experiments in Ashanti region, Ghana, using aggregation pheromone, senesced banana leaf palm alcohol extract, pseudostem, and combinations thereof, in type TAL and Voltic traps. These are the data analysed in Table 3.

Field (Type of trap)	Aggregation pheromone plus banana leaf extract	Palm alcohol extract of banana leaf	Aggregation pheromone	Pseudostem plus banana leaf extract	Pseudostem
1 (TAL)	12.58 $\pm$ 4.9	2.22 $\pm$ 0.4	9.08 $\pm$ 2.8	0.83 $\pm$ 0.6	0.42 $\pm$ 0.3
2 (Voltic)	12.43 $\pm$ 5.4	2.71 $\pm$ 1.6	4.57 $\pm$ 1.3	2.43 $\pm$ 1.1	1.00 $\pm$ 0.4
3 (TAL)	15.00 $\pm$ 3.2	3.33 $\pm$ 0.8	7.17 $\pm$ 2.6	2.42 $\pm$ 1.4	0.75 $\pm$ 0.6
4 (Voltic)	36.08 $\pm$ 9.3	5.67 $\pm$ 2.0	16.42 $\pm$ 5.7	1.25 $\pm$ 0.4	0.58 $\pm$ 0.4
5 (Voltic)	4.80 $\pm$ 1.5	0.60 $\pm$ 0.3	2.20 $\pm$ 0.7	0.80 $\pm$ 0.6	0.20 $\pm$ 0.2
<b>Sum of means</b>	<b>80.89</b>	<b>14.53</b>	<b>39.44</b>	<b>7.73</b>	<b>2.95</b>
<b>Overall mean <math>\pm</math> SE</b>	<b>16.2 <math>\pm</math> 5.3</b>	<b>2.9 <math>\pm</math> 0.8</b>	<b>7.9 <math>\pm</math> 2.4</b>	<b>1.5 <math>\pm</math> 0.4</b>	<b>0.6 <math>\pm</math> 0.1</b>

**Table 3** Analysis of variance (ANOVA) of the mean weevil catches per week data (data in Table 2 transformed to natural logarithms). The table details the sources of variation, the degrees of freedom (df), sums of squares (ss), mean squares (ms) (*i.e.* the variances), variance ratios (vr) and the *P*-values for the F-tests of the sources of variation. The ANOVA factors are denoted: *TrapType*, for Voltic vs. TAL type of trap; *ExtractOnly*, for the palm alcohol extract of senesced banana leaves treatment vs. the factorial set of four treatments involving aggregation pheromone or pseudostem; *Extract*, for the main effect of presence or absence of palm alcohol extract of senesced banana leaves; and *Treatment*, for the main effect aggregation pheromone vs. pseudostem. The dot indicates the interaction between factors. Significant ( $P < 0.05$ ) ANOVA factors of interest are given in bold.

Source of variation	df	ss	ms	vr	<i>P</i> -value
<b>Field stratum</b>					
<i>TrapType</i>	1	0.054	0.054	0.00	0.957
Residual	3	47.720	15.907	7.93	
<b>Field.Trap stratum</b>					
<i>ExtractOnly</i>	1	0.432	0.432	0.22	0.651
<i>ExtractOnly.Extract</i>	1	31.456	31.456	15.68	<b>0.002</b>
<i>ExtractOnly.Treatment</i>	1	304.937	304.9371	51.99	<b>&lt;0.001</b>
<i>ExtractOnly.TrapType</i>	1	0.007	0.007	0.00	0.954
<i>ExtractOnly.Extract.Treatment</i>	1	0.685	0.685	0.34	0.570
<i>ExtractOnly.Extract.TrapType</i>	1	0.310	0.310	0.15	0.701
<i>ExtractOnly.Treatment.TrapType</i>	1	0.130	0.130	0.06	0.803
<i>ExtractOnly.Extract.Treatment.TrapType</i>	1	0.289	0.289	0.14	0.711
Residual	12	24.075	2.006		
Total	24	410.096			