

2-oxoglutarate-dependent dioxygenases: a renaissance in attention for ascorbic acid in plants

Article

Published Version

Creative Commons: Attribution 4.0 (CC-BY)

Open Access

Mahmood, A. M. and Dunwell, J. M. ORCID:
<https://orcid.org/0000-0003-2147-665X> (2020) 2-oxoglutarate-dependent dioxygenases: a renaissance in attention for ascorbic acid in plants. PLoS ONE, 15 (12). e0242833. ISSN 1932-6203 doi: 10.1371/journal.pone.0242833 Available at <https://centaur.reading.ac.uk/94829/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1371/journal.pone.0242833>

Publisher: Public Library of Science

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online


RESEARCH ARTICLE

2-oxoglutarate-dependent dioxygenases: A renaissance in attention for ascorbic acid in plants

Asaad M. Mahmood^{1*}, Jim M. Dunwell²

1 Department of Biology, College of Education, University of Garmian, Kalar, KRG/Iraq, **2** School of Agriculture, Policy and Development, University of Reading, Reading, Berkshire, United Kingdom

* asaad@garmian.edu.krd



Abstract

L-Ascorbic acid (ascorbate, Vitamin C) is an essential human micronutrient that is predominantly obtained from plants. It is known to work as the major antioxidant in plants, and it underpins several environmentally induced stresses due to its use as a co-factor by certain 2-oxoglutarate-dependent (2-OG) dioxygenases [2(OG)-dioxygenases]. It is important to understand the role of 2(OG)-dioxygenases in the biosynthesis of ascorbate. The present study examined contents of ascorbate and protein-protein interaction in nine T-DNA mutants of *Arabidopsis* containing an insert in their respective (2-OG) dioxygenase genes (*At1g20270*, *At1g68080*, *At2g17720*, *At3g06290*, *At3g28490*, *At4g35810*, *At4g35820*, *At5g18900*, *At5g66060*). In this study, the amount of ascorbate in five of the mutants was shown to be almost two-fold or more than two-fold higher than in the wild type. This result may be a consequence of the insertion of the T-DNA. The prediction of possible protein interactions between 2(OG)-dioxygenases and relevant ascorbate-function players may indicate the oxidative effects of certain dioxygenase proteins in plants. It is expected that certain dioxygenases are actively involved in the metabolic and biosynthetic pathways of ascorbate. This involvement may be of importance to increase ascorbate amounts in plants for human nutrition, and to protect plant species against stress conditions.

OPEN ACCESS

Citation: Mahmood AM, Dunwell JM (2020) 2-oxoglutarate-dependent dioxygenases: A renaissance in attention for ascorbic acid in plants. PLoS ONE 15(12): e0242833. <https://doi.org/10.1371/journal.pone.0242833>

Editor: Mayank Gururani, United Arab Emirates University, UNITED ARAB EMIRATES

Received: July 29, 2020

Accepted: November 9, 2020

Published: December 8, 2020

Copyright: © 2020 Mahmood, Dunwell. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its [Supporting Information](#) files.

Funding: This work was financially supported by Kurdistan Regional Government (KRG)/Iraq

Competing interests: The authors have declared that no competing interests exist.

Introduction

Ascorbate is an essential human micronutrient that is predominantly obtained from plants. Early use of large dose of ascorbate is currently proven to provide a useful adjuvant therapy in moderate-severe patients with coronavirus (SARS-CoV-2) infection [1]. The potential benefit of increasing the level of L-ascorbic acid (ascorbate) in plants may enhance human nutrition, allow new metabolic engineering and improve the tolerance of crops under stress conditions [2,3]. Numerous pathways of ascorbate synthesis have been elucidated, and the details of such pathways are becoming more complete [4]. However, regulation of such pathways to control levels of ascorbate is still under investigation. Recent studies have discovered that a translational feedback mechanism can control ascorbate levels in plants; this discovery demonstrates

a new insight into regulation, and represents an additional method to increase ascorbate levels [5]. It is known that ascorbate-related genes (*vtc1-5*) encode pivotal enzymes, which play important roles in the biosynthesis pathways of ascorbate in plants [6]. For example, a mutation in *vtc1-1*, results in defective hormonal homeostasis (auxin/ethylene homeostasis) and nitric oxide signalling. Such mutants are expected to be more tolerant to abiotic stresses as a consequence of containing higher amount of ascorbate [2,7].

Due to the protective role of ascorbate as an antioxidant [8,9], and the use of this antioxidant as a co-factor by 2(OG)-dioxygenases [10] these enzymes might contribute in several physiological responses to unfavourable conditions in plants [11,12]. It is known that ascorbate serves as a cofactor for 2(OG)-dioxygenases, enzymes that are involved in several physiological activities [3,13]. For example, they play an important role in the synthesis of hormones including, gibberellins, abscisic acid, ethylene and the degradation of hormones such as indole acetic acid (IAA). These enzymes are also involved in the synthesis of a wide range of secondary compounds, including glucosinolates and anthocyanins [14]. In light of this evidence, mutants containing a T-DNA insert in their respective 2(OG) dioxygenase genes might demonstrate inactivation of these genes. Such inactivation is expected to lead to the subsequent inhibition/activation of other genes that encode enzymes that are actively involved in the biosynthesis of ascorbate. This involvement might increase ascorbate levels and also improve the tolerance of plants against stress conditions. To examine this prediction, the present study was designed to determine the content of ascorbate in WT and nine homozygous *Arabidopsis* mutants containing a T-DNA insert in their respective 2(OG) dioxygenase genes.

Material and methods

Plant materials

Seeds of WT and nine T-DNA insert mutant lines [N671573 (insert in *At1g20270*), N668172 (*At1g68080*), N652869 (*At2g17720*), N679576 (*At3g06290*), N678627 (*At3g28490*), N338446 (*At4g35810*), N683883 (*At4g35820*), N666896 (*At5g18900*), N598611 (*At5g66060*)] of *A. thaliana* Columbia ecotype were obtained from the European *Arabidopsis* Stock Centre (<http://arabidopsis.info/BasicForm>). Prior to germination seeds were soaked in water and kept in the fridge at 4°C for 4 d (stratified) in order to break seed dormancy.

Seed germination

Stratified seeds were grown and germinated in 6.5 x 6 cm mini-flower pots (2 seeds in each pot) containing potting growing medium (supplied from <http://www.sinclairpro.com/>) and topped with vermiculite, pots were maintained under controlled environmental conditions at 22°C and 60% Relative Humidity (RH) with a day/night cycle of 16/8 h (at 100 $\mu\text{mol m}^{-2} \text{s}^{-1}$ light intensity) in the growth chambers (Fitotron plant growth chambers, Weiss Gallenkamp, UK).

This study was conducted using a complete randomised design (CRD) with three replicates (used as biological replicates) and plants from each replicate (used as technical replicates).

Confirmation of genetic status of mutant lines

To confirm the genetic status (homozygous, heterozygous or wild type) of the underlying mutant lines, their genomic DNA (5 plants/mutant) was extracted using a DNeasy Plant Mini kit (<http://www.qiagen.com>). The PCR mixture (25 μl) contained 2 μl of DNA, 0.75 μl (0.3 μM) of each forward primer (or left border primer LBB1.3, 5' -ATTTTGCCGATTTCGG AAC-3' and NR80 5' -GCTGATACAAAACAAAACAACGA-3' were used with SALK and GK insertions respectively) and reverse primer (S1 Table), 12.5 μl 2X BioMix PCR master mix

(Bioline, UK,) and 9 μ l TE water. The reaction mixtures were amplified in a GeneAmp PCR system 9700 (Applied Biosystems). The PCR products were separated using electrophoresis in 1% agarose gels supplemented with ethidium bromide 3 μ l/70 ml. PCR products were visualized on a GelDoc-ItTS2 Imager (UVP) followed by capturing a clear image through GeneSnap (version 6.00.19) system (SynGene, UK). Finally, each homozygous plant/line was determined, labelled and seeds were collected from those plants for further investigations [15].

Sample preparation

Seeds from each homozygous plant/line were grown and germinated according to the protocol stated above. Then, samples of fresh leaves were taken from 3-week old plants (mutants and wild type) of similar morphology. 1 ml of a solution containing 0.1 M HCl and 1 mM EDTA was added to the fine-ground powder of each sample. This mixture was incubated for 5 minutes, after which the mixture was thawed and then centrifuged at 13 000 rpm for 10 min. The supernatant was filtered and transferred to a new 1.5 ml Eppendorf tube. All steps were conducted at 4°C in a dark room.

Determination of ascorbate levels

Prepared samples were immediately used to measure total content of ascorbate using Capillary Electrophoresis (EC). Linear standards (0.1, 0.2, 0.3, 0.4, and 0.5) mM of L-ascorbate were dissolved in the same solvent and run in parallel with three technical replicates of each sample. All reads (Peak areas) were measured with the absorbance at 254 nm (S1 Fig). The measured area of each sample was converted to concentration using a linear equation generated from the intercept of standard values. The standard curve for ascorbate was generated with a linear intercept starting from zero over a broad range of concentrations up to 0.5 μ Mol. The value of coefficient of determination ($R^2 = 0.9958$) was highly significant and the generated linear equation was used to calculate and normalise the amount of total ascorbate in WT and the nine homozygous mutant lines (Fig 1).

Furthermore, to quantify the amount of ascorbate, the measured peak area of each replicate/mutant and WT was converted to concentration using the linear equation ($Y = 231.93X$) in which Y refers to the concentration of ascorbate and X indicates area measurement of each experimental unit (replicate). The amount of ascorbate was adjusted to be expressed as μ Molar/g fresh weight. Results were analysed using a CRD with one-way ANOVA and means compared using the Duncan's multiple range test.




Fig 1. Standard curve for ascorbate dissolved in 1 ml of 0.1 M HCl and 1 mM EDTA.

<https://doi.org/10.1371/journal.pone.0242833.g001>

Protein-protein interactions

Protein-protein interactions in association with physical and biological activities were investigated between underlying dioxygenases and enzymes involved in ascorbate biosynthesis of ascorbate. We used the latest version of String (11.0) protein-protein interaction networks (<http://string-db.org/>). This software is regarded as a valuable means to study comprehensive functional associations and integrations between proteins of 5090 organisms including *A. thaliana* [16].

Results

Ascorbate levels in the different mutant lines

The results indicate that five mutant lines (Fig 2) showed significantly higher levels of ascorbate compared to the wild type control (WT). The maximal ascorbate content (4.70 ± 0.3 Mol/g FW) was detected in the N666896 mutant. This amount was significantly higher by more than two fold compared with the WT (1.9 ± 0.13 Mol/g FW). Moreover, the ascorbate levels in the other three mutants (N652869, N338446 and N683883) was also significantly different compared to the WT. The amount of ascorbate in the latter mutants was significantly higher by more than two fold (4.27 ± 0.2 , 4.37 ± 0.1 and 4.48 ± 0.2 Mol/g FW respectively) compared with the WT. Despite giving a slightly higher level of ascorbate, the N668172 mutant showed a significantly higher content of ascorbate (2.93 ± 0.14 Mol/g FW) than the WT.

However, the results indicate that in four mutant lines, N598611, N678627, N671573, and N679576, no significant differences in ascorbate contents were detected between them (1.8 ± 0.1 , 2.15 ± 0.1 , 2.31 ± 0.2 , and 2.80 ± 0.2 Mol/g FW respectively) and the WT control (1.9 ± 0.13 Mol/g FW). Although the mutant N668172 showed a significant difference compared to the WT, no significant differences were observed between the latter mutant and both N671573 and N679576 mutants.

Protein-protein interactions with relevant ascorbate-function proteins

An analysis was conducted to determine whether the activities of respective dioxygenase proteins directly affect enzymes involved in ascorbate biosynthesis. These enzymes included: CYT1 or VTC1, a GDP-mannose pyrophosphorylase/mannose-1-pyrophosphatase [17];




Fig 2. The content of ascorbate in WT and nine mutant lines. The values are expressed as means \pm SE ($n = 9$), ($P < 0.05$). Means followed by the same letters within a column do not differ significantly from each other according to Duncan's multiple range tests. FW denotes fresh weight.

<https://doi.org/10.1371/journal.pone.0242833.g002>

VTC2, a GDP-L-galactose phosphorylase 1 [18]; VTC3, a protein kinase and PP2C-like domain-containing protein [19]; VTC4, a L-galactose-1-phosphate phosphatase [20], and VTC5, a GDP-L-galactose phosphorylase [21]. The results of protein-protein interactions presented in Fig 3 show that indirect protein links are predicted between respective proteins and VTC1-5 in *A. thaliana*.

In this figure several interactions were identified; they included:- (I) Proline dehydrogenase 1 (EARLY RESPONSIVE TO DEHYDRATION 5: ERD5) that helps the recovery of plants from drought stress and is involved in the biosynthesis of proline [22,23], (II) PYRROLINE-5-CARBOXYLATE REDUCTASE (P5CR), an enzyme linked to the stress-induced accumulation of proline [24], (III) MYO-INOSITOL OXYGENASE 2 (MIOX2), an enzyme involved in ascorbate biosynthesis in plants. VTC1 (CYT1) was shown to interact with dioxygenases by only one textmining line ('automated text-mining is applied to uncover statistical and/or semantic links between proteins, based on Medline abstracts and a large collection of full-text articles') [25]. However, neither VTC2 nor VTC5 had any direct links with dioxygenases. On the other hand, VTC4 was intermediate between both VTC2 and VTC5 with dioxygenase by two steps through both MIOX2 and ERD5. These combinations were based on the textmining and database lines. Moreover, in this protein-protein interaction study there was no obvious association between VTC3 and respective proteins.

While various protein-protein interactions were predicted between underlying proteins and VTC, MIOX2, ERD5 and P5CR, no gene co-expression was predicted between latter proteins and selected dioxygenases (Fig 4).

Discussion

Ascorbate levels

Based on their ascorbate levels, mutants could be divided into two groups. The first group comprise five mutant lines that had significant increases in the level of ascorbate in comparison with WT. This increase may be linked to the down-regulation of respective genes in mutants leading to the activation of other genes that encode enzymes involved in the biosynthetic pathways of ascorbate [11,26,27].

The second group comprise four mutants that all showed an increased amount of ascorbate compared to the WT, although these increases were not significant. In summary, these results




Fig 3. A prediction of protein-protein links between respective dioxygenases and other ascorbate-related proteins in *A. thaliana* (<http://string-db.org/>).

<https://doi.org/10.1371/journal.pone.0242833.g003>




Fig 4. A prediction of co-expression links between respective dioxygenase genes and other ascorbate-related genes in *A. thaliana* and other organisms (<http://string-db.org/>).

<https://doi.org/10.1371/journal.pone.0242833.g004>

may be linked to the suppression of respective genes in these mutants leading to the activation of other genes that encode enzymes involved in the biosynthetic pathways of ascorbate [26,27].

The ascorbate synthesis route in plants is principally involves the L-galactose pathway, which is also known as the Wheeler–Smirnoff pathway [28]. The enzyme GDP-L galactose phosphorylase (GGP, also known as VTC4) has an important role and is responsible for the sixth enzymatic step of this pathway. Numerous studies have documented that the overexpression of the GGP gene consistently leads to a two–six fold increases in ascorbate levels in a wide range of species including *A. thaliana* and [3,29]. Given the above premises, a link among 2 (OG)-dioxygenase genes and the GGP gene expression is possible, whereby metabolic reprogramming occurs in the mutants studied. The present results indicate that suppression of certain 2(OG)-dioxygenase genes may possibly up-regulate expression of the GGP gene and subsequently increase the ascorbate level in plants.

Although ascorbate has an active role as a co-factor of 2(OG)-dioxygenase family enzymes, it is also a natural substrate of ascorbate peroxidases in plants [30]. These latter enzymes are important players in regulating levels of cellular reactive oxygen species (ROS) under stress conditions [31] and in addition, 2(OG)-dioxygenase are thought to act as direct sensors of oxygen and regulators of hypoxia in both plants, and animals. Specifically, ROS metabolism consists of a complex network, which interacts closely with hormonal signalling systems, and allows plants a subtle developmental regulation against biotic and abiotic stress responses [32,33]. In light of the results from the present study, it is expected that selected mutants found to contain a high amount of ascorbate may be more tolerant to abiotic stress including ozone (O_3), high salinity and drought stress [2,6,11]. Despite the expectation of that selected dioxygenases are involved in the metabolic and biosynthetic pathways of ascorbate, to date there is no experimental evidence to confirm this idea yet. However, to further investigate this concept and to inform future studies, we studied protein-protein interactions between specific dioxygenase enzymes and ascorbate-function proteins.

Protein-protein interactions with relevant ascorbate-function proteins

Analysis of Protein-protein interactions predicted several interactions with gene products known to be involved in biosynthetic and metabolic processes of ascorbate. The prediction of protein interactions between VTC, MIOX and the selected 2(OG)-dioxygenases may indicate the latter enzymes are involved in regulatory mechanisms during embryonic development,

germination, fruit ripening and in the regulation of auxins, which comprise an important class of plant hormone. These regulatory mechanisms are known to be involved in ascorbate biosynthesis [34–37]. Possible protein interaction between P5CR, VTC and underlying 2(OG)-dioxygenases, may be a signal of the active role of such dioxygenases in protecting plants against drought and salinity. These expectations derive from previous studies demonstrating the role of P5CR in the biosynthetic pathways of proline, a widely distributed osmolyte that protect plants from stress conditions [38,39]. ERD5 is another protein predicted to be intermediated between VTC and underlying proteins. These interactions indicate the oxidative effect of dioxygenase proteins in the epigenetic changes of plants; these changes may serve as adaptive mechanisms to environmentally induced abiotic stress. This expectation derives from findings that reveal the involvement of ERD5 in the adaptation of *A. thaliana* accessions to cold, drought and salt tolerance [40]. It has also been shown that one of the mechanisms underlying the early adaptation of plants to salt stress is associated with an epigenetic mark [23].

In conclusion, evidence presented here strongly suggest that products of certain dioxygenase genes are expected to be actively involved in the metabolic and biosynthetic pathways of ascorbate [41]. This involvement may be of importance in future efforts to increase ascorbate amounts in plants for human nutrition [1–3,6,11,42], and to protect crop species against unfavourable conditions including drought, cold, light, O₃ and salinity [7,43].

Supporting information

S1 Fig. Showing area measurements of ascorbate level (at approximately 6 min.) in WT and nine mutants.

(ZIP)

S1 Table. List of primer sequences for detecting DNA insertion in mutant lines of *Arabidopsis*.

(DOCX)

Acknowledgments

I would like to express my appreciation to the Ministry of higher education/KRG, Human Capacity Development Program (HCDP) staff and staff members of the School of Biological Sciences, University of Reading for their tremendous support, help and great encouragement given to me throughout the research period.

Author Contributions

Data curation: Asaad M. Mahmood.

Formal analysis: Asaad M. Mahmood.

Investigation: Asaad M. Mahmood.

Methodology: Asaad M. Mahmood.

Supervision: Jim M. Dunwell.

Writing – original draft: Asaad M. Mahmood.

Writing – review & editing: Asaad M. Mahmood.

References

1. Hernandez A, Papadakos PJ, Torres A, González D, Vives M, Ferrando C, et al. Two known therapies could be useful as adjuvant therapy in critical patients infected by COVID-19. *Rev Esp Anesthesiol Reanim*. 2020; 67(5):245–252. <https://doi.org/10.1016/j.redar.2020.03.004> PMID: 32303365
2. Smirnoff N. Vitamin C booster. *Nat Biotechnol*. 2003; 21(2):134–136. <https://doi.org/10.1038/nbt0203-134> PMID: 12560838
3. Macknight RC, Laing WA, Bulley SM, Broad RC, Johnson AA, Hellens RP. Increasing ascorbate levels in crops to enhance human nutrition and plant abiotic stress tolerance. *Curr Opin Biotechnol*. 2017; 44:153–160. <https://doi.org/10.1016/j.copbio.2017.01.011> PMID: 28231513
4. Bulley S, Laing W. The regulation of ascorbate biosynthesis. *Curr Opin Plant Biol*. 2016; 33:15–22. <https://doi.org/10.1016/j.pbi.2016.04.010> PMID: 27179323
5. Laing WA, Martínez-Sánchez M, Wright MA, Bulley SM, Brewster D, Dare AP, et al. An upstream open reading frame is essential for feedback regulation of ascorbate biosynthesis in *Arabidopsis*. *Plant Cell*. 2015; 27(3):772–786. <https://doi.org/10.1105/tpc.114.133777> PMID: 25724639
6. Barth C, Gouzd ZA, Steele HP, Imperio RM. A mutation in GDP-mannose pyrophosphorylase causes conditional hypersensitivity to ammonium, resulting in *Arabidopsis* root growth inhibition, altered ammonium metabolism, and hormone homeostasis. *J Exp Bot*. 2010; 61(2):379–394. <https://doi.org/10.1093/jxb/erp310> PMID: 20007685
7. Veljović-Jovanović S, Vidović M, Morina F. Ascorbate as a key player in plant abiotic stress response and tolerance. *Ascorbic acid in plant growth, development and stress tolerance*: Springer; 2017. p. 47–109.
8. Smirnoff N. L-ascorbic acid biosynthesis. *Vitam Horm*. 2001; 61:241–266. [https://doi.org/10.1016/s0083-6729\(01\)61008-2](https://doi.org/10.1016/s0083-6729(01)61008-2) PMID: 11153268
9. McDonough MA, Loenarz C, Chowdhury R, Clifton IJ, Schofield CJ. Structural studies on human 2-oxoglutarate dependent oxygenases. *Curr Opin Struct Biol*. 2010; 20(6):659–672. <https://doi.org/10.1016/j.sbi.2010.08.006> PMID: 20888218
10. Gao Y, Han Z, Li Q, Wu Y, Shi X, Ai Z, et al. Vitamin C induces a pluripotent state in mouse embryonic stem cells by modulating micro RNA expression. *FEBS J*. 2015; 282(4):685–699. <https://doi.org/10.1111/febs.13173> PMID: 25491368
11. Wang J, Yu Y, Zhang Z, Quan R, Zhang H, Ma L, et al. *Arabidopsis* CSN5B interacts with VTC1 and modulates ascorbic acid synthesis. *Plant Cell*. 2013; 25(2):625–636. <https://doi.org/10.1105/tpc.112.106880> PMID: 23424245
12. Martínez-Andújar C, Martínez-Pérez A, Ferrández-Ayela A, Albacete A, Martínez-Melgarejo PA, Dodd IC, et al. Impact of overexpression of 9-cis-epoxycarotenoid dioxygenase on growth and gene expression under salinity stress. *Plant Sci*. 2020; 295:110268. <https://doi.org/10.1016/j.plantsci.2019.110268> PMID: 32534608
13. Mahmood AM, Dunwell JM. Evidence for novel epigenetic marks within plants. *Aims Genet*. 2019; 6(4):70–87. <https://doi.org/10.3934/genet.2019.4.70> PMID: 31922011
14. Smirnoff N. Ascorbic acid metabolism and functions: A comparison of plants and mammals. *Free Radic Biol Med*. 2018; 122:116–129. <https://doi.org/10.1016/j.freeradbiomed.2018.03.033> PMID: 29567393
15. Mahmood AM, Dunwell JM. Nuclear localization signals, genetic characterisation and morphological study of wild type and 14 *Arabidopsis* mutant lines. *Journal of University of Garmian*. 4(ICBS Conference):676–692.
16. Szklarczyk D, Gable AL, Lyon D, Junge A, Wyder S, Huerta-Cepas J, et al. STRING v11: protein–protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic Acids Res*. 2019; 47(D1):D607–D613. <https://doi.org/10.1093/nar/gky1131> PMID: 30476243
17. Conklin PL, Norris SR, Wheeler GL, Williams EH, Smirnoff N, Last RL. Genetic evidence for the role of GDP-mannose in plant ascorbic acid (vitamin C) biosynthesis. *Proc Natl Acad Sci U S A*. 1999; 96(7):4198–4203. <https://doi.org/10.1073/pnas.96.7.4198> PMID: 10097187
18. Gao Y, Badejo AA, Shibata H, Sawa Y, Maruta T, Shigeoka S, et al. Expression analysis of the *VTC2* and *VTC5* genes encoding GDP-L-galactose phosphorylase, an enzyme involved in ascorbate biosynthesis, in *Arabidopsis thaliana*. *Biosci Biotechnol Biochem*. 2011; 110320:1–6. <https://doi.org/10.1271/bbb.110320> PMID: 21897033
19. Conklin PL, DePaolo D, Wintle B, Schatz C, Buckenmeyer G. Identification of *Arabidopsis* VTC3 as a putative and unique dual function protein kinase:: protein phosphatase involved in the regulation of the ascorbic acid pool in plants. *J Exp Bot*. 2013; 64(10):2793–2804. <https://doi.org/10.1093/jxb/ert140> PMID: 23749562

20. Torabinejad J, Donahue JL, Gunesequera BN, Allen-Daniels MJ, Gillaspay GE. VTC4 is a bifunctional enzyme that affects myoinositol and ascorbate biosynthesis in plants. *Plant Physiol.* 2009; 150(2):951–961. <https://doi.org/10.1104/pp.108.135129> PMID: 19339506
21. Linster CL, Adler LN, Webb K, Christensen KC, Brenner C, Clarke SG. A second GDP-L-galactose phosphorylase in *Arabidopsis* en route to vitamin C covalent intermediate and substrate requirements for the conserved reaction. *J Biol Chem.* 2008; 283(27):18483–18492. <https://doi.org/10.1074/jbc.M802594200> PMID: 18463094
22. Cecchini NM, Monteoliva MI, Alvarez ME. Proline dehydrogenase contributes to pathogen defense in *Arabidopsis*. *Plant Physiol.* 2011; 155(4):1947–1959. <https://doi.org/10.1104/pp.110.167163> PMID: 21311034
23. Borges AA, Jiménez-Arias D, Expósito-Rodríguez M, Sandalio LM, Pérez JA. Priming crops against biotic and abiotic stresses: MSB as a tool for studying mechanisms. *Front in Plant Sci.* 2014; 5:642. <https://doi.org/10.3389/fpls.2014.00642> PMID: 25429297
24. Giberti S, Funck D, Forlani G. Δ 1-pyrroline-5-carboxylate reductase from *Arabidopsis thaliana*: stimulation or inhibition by chloride ions and feedback regulation by proline depend on whether NADPH or NADH acts as co-substrate. *New Phytol.* 2014; 202(3):911–919. <https://doi.org/10.1111/nph.12701> PMID: 24467670
25. Szklarczyk D, Franceschini A, Wyder S, Forslund K, Heller D, Huerta-Cepas J, et al. STRING v10: protein–protein interaction networks, integrated over the tree of life. *Nucleic Acids Res.* 2015; 43(D1): D447–D452. <https://doi.org/10.1093/nar/gku1003> PMID: 25352553
26. Ye N, Zhu G, Liu Y, Zhang A, Li Y, Liu R, et al. Ascorbic acid and reactive oxygen species are involved in the inhibition of seed germination by abscisic acid in rice seeds. *J Exp Bot.* 2012; 63(5):1809–1822. <https://doi.org/10.1093/jxb/err336> PMID: 22200664
27. Ye N, Zhang J. Antagonism between abscisic acid and gibberellins is partially mediated by ascorbic acid during seed germination in rice. *Plant Signal Behav.* 2012; 7(5):563–565. <https://doi.org/10.4161/psb.19919> PMID: 22516812
28. Wheeler G, Jones M, Smirnov N. The biosynthetic pathway of vitamin C in higher plants. *Nature.* 1998; 393(6683):365–369. <https://doi.org/10.1038/30728> PMID: 9620799
29. Broad RC, Bonneau JP, Hellens RP, Johnson AA. Manipulation of ascorbate biosynthetic, recycling, and regulatory pathways for improved abiotic stress tolerance in plants. *Int J Mol Sci.* 2020; 21(5):1790.
30. Xia X-J, Zhou Y-H, Shi K, Zhou J, Foyer CH, Yu J-Q. Interplay between reactive oxygen species and hormones in the control of plant development and stress tolerance. *J Exp Bot.* 2015; 66(10):2839–2856. <https://doi.org/10.1093/jxb/erv089> PMID: 25788732
31. Van Aken O, Van Breusegem F. Licensed to kill: mitochondria, chloroplasts, and cell death. *Trends Plant Sci.* 2015; 20(11):754–766. <https://doi.org/10.1016/j.tplants.2015.08.002> PMID: 26442680
32. Suzuki N, Miller G, Sejima H, Harper J, Mittler R. Enhanced seed production under prolonged heat stress conditions in *Arabidopsis thaliana* plants deficient in cytosolic ascorbate peroxidase 2. *J Exp Bot.* 2013; 64(1):2532–2563. <https://doi.org/10.1093/jxb/ers335> PMID: 23183257
33. Mittler R, Blumwald E. Genetic engineering for modern agriculture: challenges and perspectives. *Annu Rev Plant Biol.* 2010; 61:443–462. <https://doi.org/10.1146/annurev-arplant-042809-112116> PMID: 20192746
34. Lorence A, Chevone BI, Mendes P, Nessler CL. Myo-inositol oxygenase offers a possible entry point into plant ascorbate biosynthesis. *Plant Physiol.* 2004; 134(3):1200–1205. <https://doi.org/10.1104/pp.103.033936> PMID: 14976233
35. Zhang W, Gruszewski HA, Chevone BI, Nessler CL. An *Arabidopsis* purple acid phosphatase with phytase activity increases foliar ascorbate. *Plant Physiol.* 2008; 146(2):431–440. <https://doi.org/10.1104/pp.107.109934> PMID: 18065557
36. Luo M, Liu X, Singh P, Cui Y, Zimmerli L, Wu K. Chromatin modifications and remodeling in plant abiotic stress responses. *Biochim Biophys Acta.* 2012; 1819(2):129–136. <https://doi.org/10.1016/j.bbagen.2011.06.008> PMID: 21708299
37. Cruz-Rus E, Amaya I, Sanchez-Sevilla JF, Botella MA, Valpuesta V. Regulation of L-ascorbic acid content in strawberry fruits. *J Exp Bot.* 2011; 62(12):4191–4201. <https://doi.org/10.1093/jxb/err122> PMID: 21561953
38. Bohnert HJ, Nelson DE, Jensen RG. Adaptations to environmental stresses. *Plant Cell.* 1995; 7(7):1099–1111. <https://doi.org/10.1105/tpc.7.7.1099> PMID: 12242400
39. Colmer T, Voesenek L. Flooding tolerance: suites of plant traits in variable environments. *Funct Plant Biol.* 2009; 36(8):665–681. <https://doi.org/10.1071/FP09144> PMID: 32688679
40. Lewandowska-Sabat AM, Winge P, Fjellheim S, Dørum G, Bones AM, Rognli OA. Genome wide transcriptional profiling of acclimation to photoperiod in high-latitude accessions of *Arabidopsis thaliana*. *Plant Sci.* 2012; 185:143–155. <https://doi.org/10.1016/j.plantsci.2011.10.009> PMID: 22325875

41. De Tullio MC. Is ascorbic acid a key signaling molecule integrating the activities of 2-oxoglutarate-dependent dioxygenases? Shifting the paradigm. *Environ Exp Bot*. 2020; 178:104173.
42. Macknight RC, Laing WA, Bulley SM, Broad RC, Johnson AA, Hellens RP. Increasing ascorbate levels in crops to enhance human nutrition and plant abiotic stress tolerance. *Current opinion in biotechnology*. 2017; 44:153–160. <https://doi.org/10.1016/j.copbio.2017.01.011> PMID: 28231513
43. Foyer CH, Kyndt T, Hancock RD. Vitamin C in plants: novel concepts, new perspectives, and outstanding issues. *Antioxid Redox Signal*. 2020; 32(7):463–485. <https://doi.org/10.1089/ars.2019.7819> PMID: 31701753