

Hydrogen sulfide inhibits the browning of fresh-cut apple by regulating the antioxidant, energy and lipid metabolism

Article

Accepted Version

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Chen, C., Jiang, A., Liu, C., Wagstaff, C. ORCID: <https://orcid.org/0000-0001-9400-8641>, Zhao, Q., Zhang, Y. and Hu, W. (2021) Hydrogen sulfide inhibits the browning of fresh-cut apple by regulating the antioxidant, energy and lipid metabolism. *Postharvest Biology and Technology*, 175. 111487. ISSN 0925-5214 doi: 10.1016/j.postharvbio.2021.111487 Available at <https://centaur.reading.ac.uk/95960/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1016/j.postharvbio.2021.111487>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

1 **Hydrogen sulfide inhibits the browning of fresh-cut apple by**
2 **regulating the antioxidant, energy and lipid metabolism**

3
4 **Chen Chen^a, Aili Jiang^a, Chenghui Liu^a, Carol Wagstaff^b, Qiqi Zhao^a,**
5 **Yanhui Zhang^a, Wenzhong Hu^{a*}**

6
7 ^a*Key Laboratory of Biotechnology and Bioresources Utilization, Ministry of Education, College of*
8 *Life Science, Dalian Minzu University, Dalian, 116600, China*

9 ^b*School of Chemistry Food & Pharmacy, Harry Nursten Building, University of Reading,*
10 *Whiteknights, Reading, Berkshire, RG6 6DZ, UK*

11
12
13
14
15
16 *Corresponding author

17 Wenzhong Hu

18 Mailing address: College of Life Science, Dalian Minzu University, No. 18, Liaohe

19 West Road, Jinzhou New District, 116600, Dalian, P. R. China

20 Tel.: +86 411 87656213 E-mail: hwz@dlmu.edu.cn

Abstract: Surface browning is the primary limiting factor for the shelf-life of fresh-cut apple. Hydrogen sulfide (H₂S) treatment is known to effectively inhibit the browning, however, little is known about the underlying molecular mechanism. In the present paper RNA-Seq technology was used to analyse the transcript expression profiles of control and H₂S treated fresh-cut apple immediately after treatment (C0 and S0) and 6 d of storage (C6 and S6) at 4 °C. The results identified 3782 and 1164 differentially expressed unigenes (DEGs) in S0 vs. C0 and S6 vs. C6, respectively. Expression of most DEGs related to antioxidant systems and energy metabolism was up-regulated after H₂S treatment, whilst expression of genes encoding polyphenol oxidase, peroxidase, lipid-degrading enzymes, such as lipoxygenase and phospholipase D, was repressed. Further quantitative real-time PCR testing validated the reliability of our RNA-Seq results. We therefore propose that H₂S treatment inhibited the surface browning of fresh-cut apple by regulating antioxidant, energy and lipid metabolism to maintain the membrane integrity of the plant tissue.

Keywords: Fresh-cut apple; hydrogen sulfide, browning, transcriptome

1. Introduction

Fresh-cut apple, which contain phenolic components that have an antioxidant capacity in the fruit, have recently emerged as popular snacks in food service establishments, school lunch programs, and for family consumption (Guan and Fan, 2010). However, they are more perishable than intact produce because apple fruit undergo enzymatic browning during minimal processing. Surface browning is the major limiting factor for the shelf-life of fresh-cut apple, and its appearance strongly

and negatively affects the consumer's purchase decision (Shrestha et al., 2020; Toivonen and Brummell, 2008).

Enzymatic browning of fresh-cut fruit and vegetable is a complex and highly regulated process. It is generally considered that polyphenol oxidase (PPO) catalyses the oxidation of phenolic compounds to quinones which then condense to form brown polymers (Milani and Hamed, 2004). In a typical plant cell PPO is associated with the plastid and phenolic substrates are located in the vacuole. Cellular and intracellular disruption by cutting allows the substrates to mix and hence react to form browning polymers (Landrigan et al., 1996). However, PPO activity and phenolic content are not the only factors affecting browning reaction. Li et al. (2017) indicated that the lipid membrane of cells may be an important factor in browning of fresh-cut pear as a higher lipoxygenase (LOX) activity, lower unsaturated fatty acid ratio and severe cell membrane damage accompanied a stronger degree of browning. Cysteine protease inhibitors inhibited the browning of fresh-cut potato through reducing the accumulation of free amino acids (Dong et al., 2020), implying that protein degradation is an integral component of the browning mechanism. After the browning of fresh-cut lotus tubers, the ROS metabolism homeostasis was damaged and energy efficiency decreased as the antioxidant enzymes (TPX and SOD) and energy metabolism related protein (H⁺-ATPase, PDC) were down-regulated (Jiang et al., 2012). Recent studies have used high-throughput transcriptome sequencing techniques to characterize the expression profile of the browning response in different types of fruit and vegetable after cutting. Several enzymatic browning-related differentially expressed genes are consistently implicated, including *PPO*, *phenylalanine ammonia lyase (PAL)*, *peroxidase (POD)*, *catalase (CAT)*, *superoxide dismutase (SOD)* (Docimo et al., 2016; Zhang et al., 2019; Zhu et al., 2017). However,

the transcriptomic events that occur during the browning process of fresh-cut apple is still unknown.

Hydrogen sulfide (H₂S) is the third endogenous signaling molecule after carbon monoxide and nitric oxide, which is traditionally known as a toxic gas. In the past decade, studies showed that it can be generated in many types of mammalian cells and might be beneficial to human health via multiple mechanisms (Giovinazzo et al., 2021; Wang, 2002). In plant, H₂S has been revealed not only to regulate normal physiological processes (Hancock and Whiteman, 2014; Jin and Pei, 2015), but also to prevent the postharvest senescence of fruit and vegetable (Hu et al., 2012; Hu et al., 2014a). Recent research has demonstrated H₂S treatment could inhibit surface browning in many types of fresh-cut fruit and vegetable by regulating the phenolic metabolism or antioxidant defense system (Sun et al., 2015; Zheng et al., 2016). In these paper, the application concentration of H₂S were quite low. Furthermore, the essential oils in organosulphur rich fruits and vegetables, such as onion, garlic, shallot, leek, can be metabolized to generate H₂S in biological conditions and modulate cell signaling (Liang et al., 2015). Thus we propose that trace H₂S gas used in storage and preservation of fruits and vegetables could be safe. However, the underlying molecular mechanism of browning inhibition by H₂S is still unknown. In the present study, transcriptome analysis was performed using a high-throughput sequencing platform (Illumina HiSeqTM2000) to give new insights into the mechanism of browning inhibition by H₂S treatment in fresh-cut apple.

2. Materials and Methods

2.1 Sample preparation and treatments

“Fuji” apple (*Malus domestica* cv. Red Fuji) were freshly harvested from a commercial farm in Dalian City, P. R. China in October 2017. Apples were selected based on uniform color, size, hardness, and the absence of any visible physical defects and fungal infections. After harvest, the fruit were immediately transported to the laboratory at Dalian Minzu University and stored at 4 °C prior to the experiments. In these apples, weight (312.9 ± 37.4 g), equatorial (9.1 ± 0.5 cm), longitudinal (8.1 ± 0.3 cm) diameters were monitored. These fruits were rinsed gently by hand, using tap water, and were dried naturally. After that, apples were peeled, cored, then cut into 2 cm-thick cubes with a sharp stainless-steel knife. The cubes of each apple were divided into two groups, one was used for the H₂S treatment and the other one was the control. Sodium hydrosulfide (NaHS, Sigma) solutions were used as the H₂S donor and 200 mL of $0.7 \text{ mmol} \cdot \text{L}^{-1}$ NaHS was prepared and placed into the bottom of sealed containers (desiccators) which themselves had a total volume 3 L. This solution could release H₂S gas into the container headspace rapidly and reached the peak concentration within 30 min, which was then maintained at a constant concentration of $0.30 \times 10^{-10} \text{ mol} \cdot \text{L}^{-1}$ H₂S gas for the duration of sample treatment (Hu et al., 2012). Fresh-cut apple were placed on a grid above the NaHS solution and thereby exposed to H₂S treatment for 24 h. Water was used instead of NaHS solution in the control samples. Three independent containers were set up to generate H₂S and likewise there were three independent control samples. After treatment, samples were placed in a different plastic box ($15 \times 21 \times 2.5$ cm) for each replicate and each box was wrapped with polyethylene cling film to prevent further gas exchange. The time

when the control and H₂S treatment ended was defined as day 0 and from this time point on, all samples were stored for up to 12 d at 4 ± 1 °C.

2.2 Measurement of color

The color of fresh-cut apple was measured every two days using repeated measures of eight fresh cut apple cubes. Color data were captured using a Minolta Chroma Meter Model CR-300 (Minolta, Tokyo, Japan), using the CIELAB color parameters, L^* (lightness), a^* (green chromaticity) and b^* (yellow chromaticity). Browning index (BI) was calculated as follows (Palou et al., 1999):

$$BI = [100(x-0.31)]/0.172$$

where $x = (a^*+1.75L^*)/(5.645L^*+a^*-3.012b^*)$

2.3 Enzyme activities

All enzyme extracts were prepared by homogenizing 5 g of fresh-cut apple in a homogenizer (T-25, IKA, Germany), on ice, using the following extraction media: 20 mL of 0.2 mol L⁻¹ phosphate buffer (pH 6.4) containing 0.1 g polyvinylpyrrolidone (PVPP) for PPO, POD, SOD, CAT, glutathione reductase (GR) and 20 mL of 0.1 mol L⁻¹ phosphate buffer (pH 7.5) containing 1 mmol L⁻¹ ethylenediaminetetraacetic acid (EDTA) and 3 mmol L⁻¹ ascorbic acid for ascorbate peroxidase (APX). For phenylalanine ammonia lyase (PAL): the extraction solution was 20 mL of 0.1 mol L⁻¹ sodium borate buffer (pH 8.7) containing 0.037 % EDTA, 0.137 % β-mercaptoethanol and 3 % PVPP. Extracts were then centrifuged at 12 000 g, for 30 min, at 4 °C, and supernatants were collected for further analysis.

PPO and POD activities were examined using the method reported by Chen et al. (2016a). One unit of PPO and POD activities was defined as a decrease of OD value in absorbance per minute.

PAL activity was measured by using a method reported by Yin et al. (2012). One unit of PAL activity was defined as a decrease of $0.01 \times$ OD value in absorbance per minute.

SOD activity was measured using p-nitro-blue tetrazolium chloride (NBT) according to Ren et al. (2012). One unit of SOD activity was defined as the amount of enzyme that caused 50 % inhibition of NBT.

CAT activity was determined according to Ren et al. (2012), by monitoring the disappearance of H_2O_2 by recording the decrease in absorbance at 240 nm. One unit of CAT activity was defined as a decrease of $0.01 \times$ OD value in absorbance per minute.

APX activity was measured by determining the amount of oxidized ascorbate using the method of Ren et al. (2012). One unit of APX activity was defined as a decrease in the OD value in absorbance per minute.

GR activity was examined based on the oxidation of nicotinamide-adenine dinucleotide phosphate (NADPH) and the change of absorbance at 340 nm was monitored (Zhang et al., 2017). One unit of GR activity was defined as the amount of enzyme that oxidized 1 μ mol NADPH per min.

All the enzymatic activities were expressed as $U\ kg^{-1}\ FW$.

2.4 H_2O_2 content

Three grams of fresh-cut apple were put into 5 mL of cold acetone (100 %) and

then homogenized. Subsequently, the mixture was centrifuged (Allegra X-30R, Beckman Coulter, USA) for 20 min (10000 g, 4 °C). The supernatant was used to analyze H₂O₂ content by the titanium peroxide method (Patterson et al., 1984). The result was expressed as mmol kg⁻¹. There were three independent biological replicates for the control and the H₂S treatment.

2.5 Ascorbic acid content

Ascorbic acid (AsA) content was determined according to AOAC (1990) by using titration with 2,6-dichlorophenolindophenol. The result was calculated by comparison to a standard AsA curve and expressed in mg of AsA per kg fresh weight. There were three independent biological replicates for the control and the H₂S treatment.

2.6 Phenolic content

Phenolic compounds were extracted into 75 % ethanol and the content of individual phenolics was determined by using an Agilent Technologies (Waldbronn, Germany) 1100 Series HPLC system according to Chen et al. (2016b). Ten µL of each sample was filtered and then separated using a Hypersil BDS C18 column (250 mm × 4.6 mm, 5 µm; Thermo, Bellefonte, PA, USA). The phenolics were eluted by using two mobile phases (A: 2 % v/v acetic acid in water, B: 2 % v/v acetic acid in water and acetonitrile, 50:50, v/v) with a linear gradient from 10 to 55 % B in 50 min and monitored at 254 nm (for quercetin, eugenol, epicatechin), 280 nm (for catechin, tannic acid, vanillic acid) and 320 nm (for caffeic acid and chlorogenic acid). The

flow rate was 1 mL min⁻¹. The result was calculated by comparison to standard curves constructed from dilutions of pure standards for each phenolic compound and expressed as mg per kg of sample.

2.7 Transcriptome Analysis

2.7.1 mRNA library construction and sequencing

Total RNA was extracted from each biological replicate of the control and H₂S treatment using Trizol reagent (Invitrogen, CA, USA) following the manufacturer's procedure. The total RNA quantity and purity were analysed using a Bioanalyzer 2100 and RNA 6000 Nano LabChip Kit (Agilent, CA, USA) with RIN number >7.0. Approximately 10 µg of total RNA representing a specific adipose type was subjected to isolation of Poly (A) mRNA with poly-T oligo attached magnetic beads (Invitrogen). Following purification, the mRNA was fragmented into small pieces using divalent cations under elevated temperature. Then the cleaved RNA fragments were reverse-transcribed to create the final cDNA library in accordance with the protocol for the mRNA Seq sample preparation kit (Illumina, San Diego, USA), the average insert size for the paired-end libraries was 300 bp (±50 bp). We then performed the paired-end sequencing on an IlluminaHiseq4000 at the LC Sciences (Hangzhou, Zhejiang, China), following the vendor's recommended protocol.

2.7.2 RNA-seq reads mapping

The clean reads were mapped onto the apple genome GDDH13 Version 1.1 (<https://iris.angers.inra.fr/gddh13/the-apple-genome-downloads.html>) using HISAT (Johns Hopkins University Center for Computational Biology, Baltimore, MD, USA).

<https://daehwankimlab.github.io/hisat2/,version:hisat2-2.0.4>), which initially removed a portion of the reads based on quality information accompanying each read and then mapped the reads to the reference genome (Kim et al., 2015). HISAT allows multiple alignments pre-read (up to 20 by default) and a maximum of two mismatches when mapping the reads to the reference. HISAT build a database of potential splice junctions and confirms these by comparing the previously unmapped reads against the database of putative junctions.

2.7.3 Transcript abundance estimation and differentially expressed testing

The mapped reads of each sample were assembled using StringTie (http://ccb.jhu.edu/software/stringtie/,version:stringtie-1.3.4d.Linux_x86_64, Pertea et al., 2015) with default parameters (command line: `~stringtie -p 4 -G genome.gtf -o output.gtf -l sample input.bam`). Then, all transcriptomes from samples were merged to reconstruct a comprehensive transcriptome using perl scripts. After the final transcriptome was generated, StringTie and Ballgown (<http://www.bioconductor.org/packages/release/bioc/html/ballgown.html>) was used to estimate the expression levels of all transcripts (Frazee et al., 2015; Pertea et al., 2015). StringTie was used to perform expression level for mRNAs by calculating FPKM. The differentially expressed mRNAs and genes were selected with $|\log_2FC| \geq 1$ and with statistical significance ($P < 0.05$) by R package-Ballgown

2.8 Quantitative Real-Time PCR (qRT-PCR)

To validate the RNA-Seq data, the expression of 10 selected DEGs were quantified by using qRT-PCR following the manufacturer's protocols (One Step

SYBR[®] PrimeScript[™] RT-PCR Kit, TaKaRa, Japan). Total RNA was extracted from three independent biological replicates of both the control and H₂S treated apple material and purified by using TaKaRa MiniBEST Plant RNA Extraction Kit (TaKaRa, Japan). The sequences of gene specific qRT-PCR primers are listed in Supplementary Table S1. *Md18sRNA* was used as a housekeeping gene. Calculation of the relative quantification was performed by the comparative 2^{- $\Delta\Delta$ CT} method (Livak and Schmittgen, 2001). Three biological replicates were used for qRT-PCR analysis.

3. Results and discussion

3.1 Effect of H₂S treatment on the browning of fresh-cut apple

Surface browning is one of the major limiting factors for the shelf life of fresh-cut apple and strongly and negatively affects the consumers' decision whether or not to purchase the product. As shown in Fig. 1A, the initial surface color of H₂S-treated fresh-cut apple was a clean, pale yellow. The control samples had already started to discolor from the time spent in the incubator post-cutting. As the storage time progressed, the color of control samples turned to a pale brown. H₂S treatment effectively inhibited the surface browning of fresh-cut apple over the 12 d of storage, with discoloration remaining less than the browning index (BI) of the control at day 0 throughout the experiment. The BI refers to the intensity of the brown color and has been considered an important indicator of the browning degree. A continual increase in BI of fresh-cut apple was observed during storage (Fig. 1B). H₂S treatment clearly suppressed the increase in browning degree over the entire storage period.

3.2 Physiological responses of fresh-cut apple to H₂S treatment

To assess the physiological responses of fresh-cut apple to H₂S treatment, we monitored the activities of PPO, POD, PAL, SOD, CAT, APX and GR, the contents of H₂O₂, AsA and phenolics during storage at 4 °C (Fig. 2). H₂S treatment significantly increased the enzyme activities of SOD, CAT, GR, PAL and increased the contents of AsA and phenolics, whilst it inhibited the accumulation of H₂O₂ and PPO, POD and APX activities, as compared with the control ($P < 0.05$). It is interesting to note that the activities of five enzymes (PPO, PAL, CAT, SOD and APX) reached their maximum or minimum on the 6th day of storage. Therefore, fresh-cut apple samples that were taken immediately after H₂S treatment (day 0) and after storage at 4 °C for 6 d were used for transcriptome analysis to explore the molecular mechanism of browning inhibition.

3.3 Transcriptome gene expression analysis

In order to investigate the molecular mechanism initiated by H₂S treatment in browning inhibition of fresh-cut apple, RNA-Seq libraries were designed including control and H₂S treated samples at two stage (day 0 and day 6). Supplementary Table S2 and S3 provide a quality summary of the sequence data and the number of unigene and transcript identified in each sample. There were 21, 26, 6 and 12 uniquely expressed genes were found in C0, S0, C6 and S6, respectively (Table S4). We totally identified 3782 (1638 up-regulated and 2144 down-regulated) and 1164 DEGs (529 up-regulated and 635 down-regulated) in S0 vs. C0 and S6 vs. C6, respectively, only 406 genes in common (Fig. 3A, 3B). There are more DEGs in S0 vs. C0 than that in

S6 vs. C6, indicating that a majority of transcriptional responses were manifested soon after the treatments. The top up-regulated DEG in S0 vs. C0 and S6 vs. C6 were *zinc finger AN1 domain-containing stress-associated protein 12* and *glutathione S-transferase (GST)*, respectively. *Polygalacturonase inhibitor 2* was the top down-regulated DEG in both S0 vs. C0 and S6 vs. C6 (Table S5). Then we analyze the possible browning-related genes in the DEGs of two samples. In the control, there were 775 and 879 DEGs up- and down-regulated respectively at day 6 relative to day 0. While in the H₂S treatment, 1171 and 1144 DEGs up- and down-regulated were identified respectively after 6 d storage, only 312 were in common (Fig. 3C). This limited overlap in transcriptomic events indicating that H₂S treatment altered the gene expression patterns of fresh-cut apple during storage and this change might be related to the browning inhibition.

GO term enrichment analyses were conducted for DEGs in S0 vs. C0, S6 vs. C6, C6 vs. C0 and S6 vs. S0, respectively, to evaluate the potential functions of these genes. The top 15 enriched GO terms ($P < 0.05$) of biological process, cellular component and molecular function are shown in Fig. 4A-D. When comparing the GO terms in S0 vs. C0 to S6 vs. C6, 18 out of 45 GO term were found in common. Within the biological process category, “oxidation-reduction process” was the most enriched term in both S0 vs. C0 and S6 vs. C6. Besides, several GO terms related to response to abiotic stresses, such as salt stress, cold, heat, water deprivation, were significantly enriched in both two comparable pairs. For cellular component, GO terms related to membrane, such as “plasma membrane”, “integral component of plasma membrane”, “vacuolar membrane”, dominated in both S0 vs. C0 and S6 vs. C6. In S0 vs. C0, DEGs associated with “transferase activity” was the most dominated GO term in molecular function, while that was “metal ion binding” in S6 vs. C6. There were five

common GO terms in S0 vs. C0 and S6 vs. C6, including genes coding proteins with “oxidoreductase”, “glutathione transferase activity”, “pyridoxal phosphate binding” etc.

Next, we compared the function of DEGs in C6 vs. C0 and S6 vs. S0, only 11 GO terms were found in common, indicating that H₂S treatment changed the metabolism of fresh-cut apple, which might be involved in the browning inhibition (Fig 4C, 4D). For biological process, the most dominated GO term was “oxidation-reduction process” in the control, while that was “protein ubiquitination” in the H₂S treatment. Six common GO terms were found in C6 vs. C0 and S6 vs. S0. Within the cellular component category, “cytoplasm”, “integral component of plasma membrane” and “nuclear speck” were significantly enriched in both control and H₂S treatment. “cytoplasm” was the most dominated GO terms in the control, which was the second enriched GO terms in the H₂S treatment. The largest group number of DEGs in the H₂S treatment were enriched in “nucleus”. Besides, several GO terms related to membrane, such as “plasma membrane”, “vacuolar membrane”, “endoplasmic reticulum membrane”, “Golgi membrane” and “endosome membrane” were only enriched in the H₂S treatment. The top three GO terms for molecular function in the control and H₂S treatment were “nucleic acid binding”, “oxidoreductase activity” “ubiquitin-protein transferase activity” and “protein serine/threonine kinase activity”, “nucleotide binding”, “transferase activity”, respectively. To further understand the changes induced by H₂S related to the browning process, the gene expression pattern of DEGs in the control and H₂S treatment within the common GO terms were analyzed (Fig. S1). H₂S treatment increased the number of up-regulated DEGs in “glucose import” and “glucose transmembrane transporter activity”, while increased the down-regulated DEGs in

“integral component of plasma membrane”, “lipid metabolic process”, “negative regulation of transcription”, “protein ubiquitination” and “response to cold”.

Therefore, these DEGs expression pattern changes together with the involvement of “membrane”, “nucleus” induced by H₂S might be related to the browning inhibition.

In order to characterize the pathways that related to the browning inhibition by H₂S treatment, we performed pathway enrichment analysis on DEGs based on the KEGG database with $P < 0.05$ as the threshold. We found that 25/102, 11/98, 11/125 and 6/123 pathways demonstrated significant changes in S0 vs.C0, S6 vs. C6, C6 vs. C0 and S6 vs. S0, respectively (Table S6). There were seven pathways that were all enriched in the H₂S treatment versus the control at both stages of shelf life, including two within carbohydrate metabolism and two representing lipid metabolism. The pathways that were specifically enriched in S0 vs.C0 included five of carbohydrate metabolism and six of amino acid metabolism. However, three pathways representing biosynthesis of secondary metabolites (phenylpropanoid biosynthesis, flavonoid biosynthesis and anthocyanin biosynthesis) were only found in S6 vs. C6. Two pathways “Circadian rhythm-plant” and “Biotin metabolism” were found in both control and H₂S treatment during storage. “Ubiquitin mediated proteolysis” had the largest number of DEGs in S6 vs. S0.

3.4 Differentially expressed genes related to phenol peroxidases and phenolics

Surface browning of fresh-cut products results from the oxidation of phenolic substances to quinones catalyzed by phenol oxidases (such as PPO and/or POD), which subsequently condense to form browning polymers (Degl'Innocenti et al., 2005; Saltveit, 2000). Suppression of *PPO* genes was an effective approach to reduce the

browning of potato tuber (Chi et al., 2014; Coetzer et al., 2001). In the same way, numerous studies have shown that the inhibition of surface browning of fresh-cut apple was accompanied by a decrease in PPO activity (Hemachandran et al., 2017; Saba and Sogvar, 2016). In the present study, one gene encoding PPO was up-regulated (*MSTRG.17066*, a 1.72 log₂FC significant increase) in the control at day 6 relative to day 0, but no significant change in *PPO* expression was observed in H₂S treatment during storage (Fig. 5). H₂S treatment showed significant reductions in gene expression of *PPO* relative to the control at day 0 (*MSTRG.17063*, a 2.2 log₂FC significant reduction) and day 6 (*MSTRG.9035*, a 1.9 log₂FC significant reduction). This result fits with the observed pattern of PPO activity of fresh-cut apple (Fig. 2). The PPO activity in the control samples at day 6 was higher than that at day 0 and H₂S treated samples showed consistently lower PPO activity than the control.

The POD enzyme, which uses H₂O₂ as a catalyst for the oxidation of phenolic compounds (Reyes et al., 2007), could also enhance browning reactions in the presence of ongoing PPO-mediated browning (Richard-Forget and Gauillard, 1997). In the present study, two *POD* genes (*MSTRG.11943*, *MSTRG.22268*) were induced, while one *POD* (*MSTRG.27890*) was repressed by H₂S treatment. *MSTRG.22268* was up-regulated and *MSTRG.27890* was down-regulated in the control at day 6 relative to day 0, but they did not differentially expressed in the H₂S treatment during storage. *MSTRG.11943* was down-regulated in the H₂S treatment but was not differentially expressed in the control during storage. Three genes encoding POD (*MSTRG.28170*, *MSTRG.6438*, *MSTRG.898*) were repressed in S6 vs. C6. Only *MSTRG.6438* was found differentially expressed in both control (a 2.11 log₂FC significant increase) and H₂S treatment (a 1.78 log₂FC significant increase) during storage (Fig. 5). Therefore, H₂S treatment was effective at repressing induction of *MSTRG.6438* that would

normally be expressed at a later stage post-cutting, which might be related to the browning inhibition. It is unclear why fresh-cut apple has different copies of *POD* genes and showed different expression patterns in H_2S treatment relative to the control at two stages. We speculated that the cutting process induced the production of ROS, but H_2S treatment stopped the induction of the *POD* that would normally use H_2O_2 as a catalyst for oxidation of phenolic compounds. Six-day storage led to accumulation of ROS in control tissues, providing a higher concentration of H_2O_2 and enabling significant levels of *POD*-mediated polyphenol browning. However, H_2S treatment was not able to completely inhibit either the expression of *POD* genes or the enzymic activity of *POD*. There is also a debate on the correlation between *POD* activity and browning reactions of fresh-cut fruit and vegetable in previous reports (Liu et al., 2019; Sun et al., 2015) and leads to the conclusion that the function of the *POD* gene family needs to be further studied.

The biosynthesis of phenolics, which are the substrates of enzymatic browning reactions, involves a complex network of routes based on the shikimate and phenylpropanoid pathways. Genes encoding chorismate mutase (CM), chorismate synthase (CS), PAL and shikimate O-hydroxycinnamoyltransferase (HCT), which involved in the shikimic acid pathway and subsequent conversion of chorismate to phenylalanine, and phenylpropanoid pathways were all up-regulated in S0 vs. C0 or S6 vs. C0, while *bifunctional 3-dehydroquinase dehydratase/shikimate dehydrogenase* (*3DD*, *SD*), *4-coumarate-CoA ligase* (*4CL*), were down-regulated. The implication of these findings is that H_2S treatment actually induces the synthesis of phenolic compounds and this is further supported by the biochemical assays which demonstrated the significantly higher accumulation of several phenolic compounds (chlorogenic acid, vanillic acid, catechin, tannic acid, quercetin), in H_2S treated

samples compared to the control during the storage. Chlorogenic acid and catechin were reported to be good substrates for PPO in apple (Amaki et al., 2011). Therefore, we concluded that H₂S mediated delay of browning of fresh-cut apple is not through reduction of the phenolic substrates.

3.5 Differentially expressed genes related to lipid metabolism

Although the eventual browning symptoms occur because of the interactions between PPO and/or POD enzymes and phenolic substrates, the enzymes and substrates are ordinarily separated from each other by intracellular membranes. Docimo et al. (2016) has reported that there was a lack of correlation between flesh browning and PPO activity or phenolic content in cut eggplant. Similarly, Cantos et al. (2002) has found that PPO activity and phenolic compounds are not rate-limiting in browning development of fresh-cut potatoes. Li et al. (2017) has reported that the membrane may be an important factor in browning of fresh-cut pear and the browning was caused by the cell membrane degradation which damaged the cell compartmentalization. Therefore, the progress of the browning reaction was thought to be related to the processes that affect the membrane integrity. Besides the loss of cell integrity due to processing, the breakdown of membranes within cells of fruit tissues during storage also causes the loss of cellular compartmentalization, and subsequently increases the degree of browning (Toivonen and Brummell, 2008, Li et al., 2017). Since lipids are essential constituents of the cell membrane, any alteration of membrane lipid composition such as the decrease of unsaturated fatty acids, the increase of saturated fatty acids, or degradation of membrane phospholipids, may change the biophysical and/or biochemical membrane properties and damage the

integrity of membrane (Saquet et al., 2003; Zhang et al., 2018)). In the present study, of the 13 DEGs related to the fatty acid metabolism in the control samples, three were up-regulated at day 6 compared to day 0, whereas ten were down-regulated over the same storage period. With the addition of H₂S treatment, 12 genes were differentially expressed, with ten up-regulated at day 6 compared to day 0, and two down-regulated over the same storage period. Among the DEGs in the control and H₂S treatment during storage, only two genes were in common, which were up-regulated in both samples. Comparing the DEGs in H₂S treatment to the control found that most genes (19 out of 21) were down-regulated at day 0, but three out of four genes were up-regulated at day 6 (Table S7). Glycerophospholipid metabolism also changed after H₂S treatment, 11 (six genes were up-regulated and five genes were down-regulated) and 20 DEGs (eight genes were up-regulated and 12 genes were down-regulated) were found in the control and H₂S treatment at day 6 relative to day 0, respectively. There were 37 and 10 genes differentially expressed in S0 vs. C0 and S6 vs. C6, respectively (Table S8).

LOX and PLD are two important lipids-degrading enzymes, which induce the degradation of unsaturated fatty acids and phospholipids, respectively. Mellidou et al. (2014) revealed that the expression of genes encoding LOX and PLD increased when browning developed during the storage of apple. In the present study, we did not find the same result as the literature shown, one *LOX* and one *PLD* were down-regulated in the control at day 6 relative to day 0, which was not differentially expressed in the H₂S treatment during storage. But comparing the H₂S treatment to the control, three genes encoding PLD and two genes encoding LOX were down-regulated at day 0 and two of these *PLDs* and one *LOX* maintained low expression by day 6 of storage (Fig. 5). The down-regulation of these genes might increase the content of unsaturated fatty

acids and phospholipids in H₂S treated material, thus retarding the damage of cellular membrane structure that would otherwise occur in the control. H₂S treatment reduced LOX activity and lipid peroxidation has also been observed in other fresh-cut fruit (Gao et al., 2013; Zheng et al., 2016). It can thus be suggested that H₂S treatment could suppress the activation of membrane lipids-degrading enzymes, possibly at a level of transcriptional regulation by inhibiting the expression of genes *LOX* and *PLD*, which in turn might retard the degradation of cell membrane lipids and maintain the integrity of cell membrane structure.

3.6 Differentially expressed genes related to antioxidant systems

Mechanical injury by cutting processing increased the production of reactive oxygen species (ROS), which induce membrane lipid peroxidation and promote browning reactions. Antioxidant enzymes and antioxidants inside the fruit tissues are necessary for ROS detoxification (Møller, 2001). Enhancing enzymatic and non-enzymatic antioxidant capacities to alleviate the oxidative damage is one way to inhibit the browning of fresh-cut fruit and vegetable during storage (Hu et al., 2014b; Zheng et al., 2016). H₂S treatment induced the expression of genes encoding GST, as 16 and 6 *GSTs* were found up-regulated in H₂S treatment relative to the control at day 0 and day 6, respectively. Six and two *GSTs* were up-regulated and down-regulated in the control at day 6 relative to day 0, respectively. Seven of these genes remained unchanged in the H₂S treated samples over the same storage period, but five detected *GSTs* in the H₂S treatment were all down-regulated. These results indicated that H₂S treatment activated the GSTs expression at an earlier stage, but most of them have less longevity (Fig. 5). Besides, one gene encoding CAT and one gene encoding SOD all

showed significantly higher expression, while two *APXs* showed significantly lower expression in tissue from the H_2S treatment compared to control at day 0. Physiological results supports this, the activities of SOD and CAT increased while APX activity decreased after H_2S treatment (Fig. 2). After 6 d of storage, only one APX-encoding gene was up-regulated, whilst one gene encoding APX and one encoding SOD were down-regulated in H_2S treatment relative to the control. There were three and two *SODs* down-regulated in the control and H_2S treatment, respectively. Therefore, it is reasonable to see that the number of up-regulated DEGs involved in antioxidant systems was greater in S0 vs. C0 compared to those in S6 vs. C6 (Fig. 5), implying that the antioxidant capacities of H_2S treated sample at day 0 were higher than at day 6.

AsA is an antioxidant that is able to interact with ROS directly or together with antioxidant enzymes. AsA plays an important role in the development of browning. Exogenous application of AsA effectively inhibited the browning of fresh-cut fruit (Yan et al., 2017) and browning does not occur unless the endogenous AsA concentration falls below a certain threshold value (Franck et al., 2007). In the plant cell, AsA can be continuously oxidized through enzymatic (APX and AO) or non-enzymatic reactions and can be reduced back to AsA by recycling reductases (DHAR, MDHAR) (Cocetta et al., 2014). The AsA recycling pathway has been evidenced as important for the re-establishment of AsA in horticultural crops (Mellidou and Kanellis, 2007). Although *APX* in fresh-cut apple was down-regulated in S0 vs. C0, the expression of *AO* and *MDHAR* was up-regulated in S0 vs. C0, while they were unchanged after 6 d of storage. In the control sample, one *AO* was up-regulated at day 6 relative to day 0, but three *AOs* were down-regulated and two *MDHARs* were up-regulated in the H_2S treatment during storage (Fig. 5). This result

suggested that AsA recycling pathway was triggered by H₂S, but it did not work efficiently after storage. The transcriptome results corresponded well to the content of AsA. We observed a decrease of AsA in fresh-cut apple during storage, regardless of treatment, but H₂S treatment significantly inhibited the decline. Overall, the up-regulation of antioxidant enzymes and AsA recycling-related gene expression aided ROS scavenging, as we observed that H₂S significantly alleviated the accumulation of H₂O₂ and maintained it at a lower level compared with the control throughout storage. The ability of H₂S treatment to increase the antioxidant capacity and reduce the accumulation of ROS has also been reported in various fresh-cut fruit and vegetable (Sun et al., 2015; Zheng et al., 2016). Therefore, we concluded that H₂S treatment delayed browning of fresh-cut apple might be associated with its effect on enhancing the antioxidant defense capacity, which reduced the oxidative damage from ROS.

3.7 Differentially expressed genes related to energy metabolism

An increasing body of evidence has demonstrated that the browning of postharvest horticultural crops is often accompanied by lower levels of ATP content and energy charge (Franck et al., 2007; Lin et al., 2018). ATP, as an "energy flux", plays an important role in lipid synthesis and cell membrane restoration. Once ATP production rate is lower than a certain threshold, free fatty acids will be released from phospholipids and the cell membrane structure will be damaged (Rawyler et al., 1999), which allows PPO to mix with phenolics and react to produce browning compounds. Furthermore, ATP also plays an important role in AsA cycling. ATP deficiency can lead to ROS accumulation, which resulted in membrane lipid peroxidation and the

membrane structural disruption. Adequate ATP is beneficial to inhibit flesh and peel browning of fruit (Saquet et al., 2003). In postharvest fruit, ATP is mainly generated by oxidative phosphorylation. In our work, 62.5% of DEGs related to oxidative phosphorylation were down-regulated in the control at day 6 relative to day 0. However, in H₂S treatment, that percent of down-regulated DEGs was reduced to 31.8% (Table S9), indicating that H₂S treatment retained oxidative phosphorylation capacity, which might be related to promoting browning inhibition. ATPase and NAD are key enzymes for oxidative phosphorylation (Wang et al., 2018). Jiang et al. (2012) has reported ATPase was down-regulated after fresh-cut lotus tissue browning. Transcript levels of the genes encoding the *PhATPase* γ and *PhATPase* α subunits decreased during *Phalaenopsis* explant browning (Xu et al., 2015). Seven genes encoding ATPase (include calcium-transporting ATPase, copper-transporting ATPase etc.) and three genes encoding NAD were induced after H₂S treatment (S0 vs. C0). With the prolonged storage time, the energy production capacity was gradually reduced. However, the expression of two *ATPase* and one *NAD* gene remained up-regulated in H₂S treated tissue relative to control after six days of storage (Fig. 5). Therefore, the up-regulation of these genes encoding energy metabolism enzymes involved in oxidative phosphorylation by H₂S treatment might increase the synthesis of ATP and thus provide enough energy for maintaining the cell membrane structure of fresh-cut apple.

The respiratory pathways, such as glycolysis, tricarboxylic acid (TCA) cycle, mitochondrial electron transport chain (ETC) also supply energy to maintain normal metabolism in postharvest fruit and vegetables. Most DEGs related to the respiratory pathway were down-regulated in the control fresh-cut apple during storage, while in the H₂S treated sample, the number of up-regulated DEGs in glycolysis and TCA

increased (Table S10 and Table S11). H₂S treatment therefore appears to promote the retention of metabolic processes linked to energy generation in stored fresh-cut apple. At day 0, H₂S treatment induced the expression of genes encoding sucrose synthase (*SUS*) while it repressed the expression of genes encoding fructokinase as compared to the control (Fig. 5). It is suggested that *SUS* was used to cleave sucrose to provide hexose-phosphates instead of using hexokinases, which is an ATP consuming process. This alteration could save energy by using only one molecule of PPi instead of two molecules of ATP (Boeckx et al., 2019). Three genes encoding 6-phosphofructokinase, one of the rate-limiting enzymes in glycolysis, were up-regulated by H₂S treatment. Aldehyde dehydrogenase genes, which encode a critical enzyme for anaerobic respiration, were repressed by H₂S treatment. Thus, we propose that H₂S regulated glycolysis provides substrate for ATP production, rather than being consumed by anaerobic respiration. In the TCA cycle most of the DEGs (13/15) were down-regulated in S0 vs. C0, indicating that TCA pathway of fresh-cut apple was repressed by H₂S treatment (Table S11). Furthermore, genes encoding mitochondrial respiratory chain complex I, II and III were induced in S0 vs. C0 or S6 vs. C6 (Fig. 5), which might be expected to result in the elevated synthesis of energy-generating molecules. It is worth noting that excessive respiration would also consume metabolic substances and lead to aging and senescence. Therefore, maintaining proper respiration intensity and simultaneously supplying sufficient energy play important roles in the inhibition of browning. It can be assumed that H₂S-inhibited browning of fresh-cut apple could be related to the reduction of respiratory intensity via conserving higher levels of energy in the form of ATP. This was supported by the previous reports, which confirmed that exogenous H₂S treatment significantly reduced respiration rates whilst concomitantly inducing higher

energy production (Hu et al., 2012).

3.8 Validation of RNA-Seq by qRT-PCR

A total of 10 genes with differing expression patterns were selected to validate the RNA-Seq data by qRT-PCR evaluation using specific primers (Table S1). As shown in Fig. 6, correlation analysis of the gene expression ratios showed a good correlation ($R^2 = 0.92886$) between qRT-PCR and RNASeq, indicating the high reliability of the RNA-Seq data obtained in our study.

4. Conclusions

Surface browning, induced by mechanical damage, is a complex and highly regulated process. H_2S treatment effectively inhibited surface browning and maintained the initial color of fresh-cut apple during the storage. Transcriptome analysis revealed that the browning inhibition by H_2S was mainly related to the regulation of lipid metabolism, antioxidant system, and energy metabolism. The results are helpful to understand the browning inhibition mechanism conferred by H_2S on fresh-cut apple, which is beneficial for the technological innovation of browning prevention and amelioration for fresh-cut fruit.

Funding: The work was financially supported by National Key R&D Program of China (2016YFD0400903) and the National Natural Science Foundation of China (31601517, 31801598).

Notes:

The authors declare no competing financial interest.

References:

Amaki, K., Saito, E., Taniguchi, K., Joshita, K., Murata, M., 2011. Role of chlorogenic acid quinone and interaction of chlorogenic acid quinone and catechins in the enzymatic browning of apple. *Biosci. Biotechnol. Biochem.* 75, 100444-1–4. DOI:10.1271/bbb.100444

AOAC, 1990. Vitamin C (ascorbic acid) in vitamin preparations and juices. Official methods of analysis, 15th edn. AOAC, Arlington.

Boeckx, J., Pols, S., Hertog, M.L., Nicolai, B.M., 2019. Regulation of the central carbon metabolism in apple fruit exposed to postharvest low-oxygen stress. *Front. Plant Sci.* 10, 1384. DOI:10.3389/fpls.2019.01384

Cantos, E., Tudela, J.A., Gil, M.I., Espín, J.C., 2002. Phenolic compounds and related enzymes are not rate-limiting in browning development of fresh-cut potatoes. *J. Agric. Food Chem.* 50, 3015–3023. DOI:10.1021/jf0116350

Chen, C., Hu, W., He, Y., Jiang, A., Zhang, R., 2016a. Effect of citric acid combined with UV-C on the quality of fresh-cut apple. *Postharvest Biol. Technol.* 111, 126–131. DOI:10.1016/j.postharvbio.2015.08.005

Chen, C., Hu, W., Zhang, R., Jiang, A., Zou, Y., 2016b. Levels of phenolic compounds, antioxidant capacity, and microbial counts of fresh-cut onions after treatment with a combination of nisin and citric acid. *Hortic. Environ. Biotechnol.* 57, 266–273. DOI:10.1007/s13580-016-0032-x

Chi, M., Bhagwat, B., Lane, W.D., Tang, G., Su, Y., Sun, R., Oomah, B.D., Wiersma, P.A., Xiang, Y., 2014. Reduced polyphenol oxidase gene expression and

644 enzymatic browning in potato (*Solanum tuberosum* L.) with artificial microRNAs.
645 BMC Plant Biol. 14, 62. DOI:10.1186/1471-2229-14-62

646 Cocetta, G., Baldassarre, V., Spinardi, A., Ferrante, A., 2014. Effect of cutting on
647 ascorbic acid oxidation and recycling in fresh-cut baby spinach (*Spinacia oleracea* L.)
648 leaves. Postharvest Biol. Technol. 88, 8–16. DOI:10.1016/j.postharvbio.2013.09.001

649 Coetzer, C., Corsini, D., Love, S., Pavek, J., Tumer, N., 2001. Control of
650 enzymatic browning in potato (*Solanum tuberosum* L.) by sense and antisense RNA
651 from tomato polyphenol oxidase. J. Agric. Food Chem. 49, 652–657.
652 DOI:10.1021/jf001217f

653 Degl'Innocenti, E., Guidi, L., Pardossi, A., Tognoni, F., 2005. Biochemical study
654 of leaf browning in minimally processed leaves of lettuce (*Lactuca sativa* L. var.
655 Acephala). J. Agric. Food Chem. 53, 9980–9984. DOI:10.1021/jf050927o

656 Docimo, T., Francese, G., De Palma, M., Mennella, D., Toppino, L., Lo Scalzo,
657 R., Mennella, G., Tucci, M., 2016. Insights in the fruit flesh browning mechanisms in
658 *Solanum melongena* genetic lines with opposite postcut behavior. J. Agric. Food
659 Chem. 64, 4675–4685. DOI:10.1021/acs.jafc.6b00662

660 Dong, T., Cao, Y., Jiang, C.-Z., Li, G., Liu, P., Liu, S., Wang, Q., 2020. Cysteine
661 protease inhibitors reduce enzymatic browning of potato by lowering the
662 accumulation of free amino acids. J. Agric. Food Chem. 68, 2467–2476.
663 DOI:10.1021/acs.jafc.9b07541

664 Franck, C., Lammertyn, J., Ho, Q.T., Verboven, P., Verlinden, B., Nicolai, B.M.,
665 2007. Browning disorders in pear fruit. Postharvest Biol. Technol. 43, 1–13.
666 DOI:10.1016/j.postharvbio.2006.08.008

Frazer, A.C., Pertea, G., Jaffe, A.E., Langmead, B., Salzberg, S.L., Leek, J.T.,
2015. Ballgown bridges the gap between transcriptome assembly and expression
analysis. *Nat. Biotechnol.* 33, 243–246. DOI:10.1038/nbt.3172

Gao, S.-P., Hu, K.-D., Hu, L.-Y., Li, Y.-H., Han, Y., Wang, H.-L., Lv, K., Liu,
Y.-S., Zhang, H., 2013. Hydrogen sulfide delays postharvest senescence and plays an
antioxidative role in fresh-cut kiwifruit. *HortScience* 48, 1385–1392.
DOI:10.21273/HORTSCI.48.11.1385

Giovinazzo et al., 2021. Hydrogen sulfide is neuroprotective in Alzheimer's
disease by sulphydrating GSK3 β and inhibiting Tau hyperphosphorylation. *PNAS*
118(4), e2017225118. DOI:10.1073/pnas.2017225118

Guan, W., Fan, X., 2010. Combination of sodium chlorite and calcium
propionate reduces enzymatic browning and microbial population of fresh-cut
“Granny Smith” apple. *J. Food Sci.* 75, M72–M77.
DOI:10.1111/j.1750-3841.2009.01470.x

Hancock, J.T., Whiteman, M., 2014. Hydrogen sulfide and cell signaling: team
player or referee? *Plant Physiol. Biochem.* 78, 37–42.
DOI:10.1016/j.plaphy.2014.02.012

Hemachandran, H., Anantharaman, A., Mohan, S., Mohan, G., Kumar, D.T., Dey,
D., Kumar, D., Dey, P., Choudhury, A., Doss, C.G.P., 2017. Unraveling the inhibition
mechanism of cyanidin-3-sophoroside on polyphenol oxidase and its effect on
enzymatic browning of apple. *Food Chem.* 227, 102–110.
DOI:10.1016/j.foodchem.2017.01.041

Hu, Hu, S.-L., Wu, J., Li, Y.-H., Zheng, J.-L., Wei, Z.-J., Liu, J., Wang, H.-L.,
Liu, Y.-S., Zhang, H., 2012. Hydrogen sulfide prolongs postharvest shelf life of

691 strawberry and plays an antioxidative role in fruits. J. Agric. Food Chem. 60,
692 8684–8693. DOI:10.1021/jf300728h

693 Hu, Wang, Q., Hu, L.-Y., Gao, S.-P., Wu, J., Li, Y.-H., Zheng, J.-L., Han, Y., Liu,
694 Y.-S., Zhang, H., 2014b. Hydrogen sulfide prolongs postharvest storage of fresh-cut
695 pears (*Pyrus pyrifolia*) by alleviation of oxidative damage and inhibition of fungal
696 growth. PloS one 9, e85524. DOI:10.1371/journal.pone.0085524

697 Hu, H., Shen, W., Li, P., 2014a. Effects of hydrogen sulphide on quality and
698 antioxidant capacity of mulberry fruit. Int. J. Food Sci. Tech. 49, 399–409.
699 DOI:10.1111/ijfs.12313

700 Jiang, J., Jiang, L., Zhang, L., Luo, H., Opiyo, A.M., Yu, Z., 2012. Changes of
701 protein profile in fresh-cut lotus tuber before and after browning. J. Agric. Food Chem.
702 60, 3955–3965. DOI:10.1021/jf205303y

703 Jin, Z., Pei, Y., 2015. Physiological implications of hydrogen sulfide in plants:
704 pleasant exploration behind its unpleasant odour. Oxid Med Cell Longev. 397502.
705 DOI:10.1155/2015/397502

706 Kim, D., Langmead, B., Salzberg, S.L., 2015. HISAT: a fast spliced aligner with
707 low memory requirements. Nat. Methods 12, 357–360. DOI:10.1038/nmeth.3317

708 Landrigan, M., Morris, S., Eamus, D., McGlasson, W., 1996. Postharvest water
709 relationships and tissue browning of rambutan fruit. Sci. Horticul. 66, 201-208.
710 DOI:10.1016/S0304-4238(96)00915-6

711 Li, Z., Zhang, Y., Ge, H., 2017. The membrane may be an important factor in
712 browning of fresh-cut pear. Food Chem. 230, 265–270. DOI:
713 10.1016/j.foodchem.2017.03.044

714 Liang, D., Wang, C., Tocmo, R., Wu, H., Deng, L.-W., Huang, D., 2015.
715 Hydrogen sulphide (H₂S) releasing capacity of essential oils isolated from

716 organosulphur rich fruits and vegetables. J. Funct. Foods 14, 634–640.
 717 DOI:10.1016/j.jff.2015.02.007

718 Lin, Y., Lin, Y., Lin, H., Chen, Y., Wang, H., Shi, J., 2018. Application of propyl
 719 gallate alleviates pericarp browning in harvested longan fruit by modulating
 720 metabolisms of respiration and energy. Food Chem. 240, 863–869.
 721 DOI:10.1016/j.foodchem.2017.07.118

722 Liu, X., Yang, Q., Lu, Y., Li, Y., Li, T., Zhou, B., Qiao, L., 2019. Effect of
 723 purslane (*Portulaca oleracea* L.) extract on anti-browning of fresh-cut potato slices
 724 during storage. Food Chem. 283, 445–453. DOI:10.1016/j.foodchem.2019.01.058

725 Livak, K.J., Schmittgen, T.D., 2001. Analysis of relative gene expression data
 726 using real-time quantitative PCR and the 2⁻ΔΔCT method. Methods 25, 402–408.
 727 DOI:10.1006/meth.2001.1262

728 Mellidou, I., Buts, K., Hatoum, D., Ho, Q.T., Johnston, J.W., Watkins, C.B.,
 729 Schaffer, R.J., Gapper, N.E., Giovannoni, J.J., Rudell, D.R., 2014. Transcriptomic
 730 events associated with internal browning of apple during postharvest storage. BMC
 731 Plant Biol. 14, 328. DOI:10.1186/s12870-014-0328-x

732 Mellidou, I., Kanellis, A.K., 2017. Genetic control of ascorbic acid biosynthesis
 733 and recycling in horticultural crops. Front. Chem. 5, 50.
 734 DOI:10.3389/fchem.2017.00050

735 Milani, J., Hamed, M., 2004. Susceptibility of five apple cultivars to enzymatic
 736 browning, V International Postharvest Symposium 682, 2221–2226.
 737 DOI:10.17660/ActaHortic.2005.682.303

738 Møller, I.M., 2001. Plant mitochondria and oxidative stress: electron transport,
 739 NADPH turnover, and metabolism of reactive oxygen species. Annu. Rev. Plant Biol.
 740 52, 561–591. DOI:10.1146/annurev.arplant.52.1.561

Palou, E., López-Malo, A., Barbosa-Cánovas, G., Welti-Chanes, J., Swanson, B.,
 1999. Polyphenoloxidase activity and color of blanched and high hydrostatic pressure
 treated banana puree. J. Food Sci. 64, 42–45.
 DOI:10.1111/j.1365-2621.1999.tb09857.x

Patterson, B.D., MacRae, E.A., Ferguson, I.B., 1984. Estimation of hydrogen
 peroxide in plant extracts using titanium (IV). Anal. Biochem. 139, 487–492.
 DOI:10.1016/0003-2697(84)90039-3

Pertea, M., Pertea, G.M., Antonescu, C.M., Chang, T.-C., Mendell, J.T., Salzberg,
 S.L., 2015. StringTie enables improved reconstruction of a transcriptome from
 RNA-seq reads. Nat. Biotechnol. 33, 290–295. DOI:10.1038/nbt.3122

Rawyler, A., Pavelic, D., Gianinazzi, C., Oberson, J., Braendle, R., 1999.
 Membrane lipid integrity relies on a threshold of ATP production rate in potato cell
 cultures submitted to anoxia. Plant Physiol. 120, 293–300. DOI:10.1104/pp.120.1.293

Ren, Y., Wang, Y., Bi, Y., Ge, Y., Wang, Y., Fan, C., Li, D., Deng, H., 2012.
 Postharvest BTH treatment induced disease resistance and enhanced reactive oxygen
 species metabolism in muskmelon (*Cucumis melo* L.) fruit. Eur. Food Res. Technol.
 234, 963–971. DOI:10.1007/s00217-012-1715-x

Reyes, L.F., Villarreal, J.E., Cisneros-Zevallos, L., 2007. The increase in
 antioxidant capacity after wounding depends on the type of fruit or vegetable tissue.
 Food Chem. 101, 1254–1262. DOI:10.1016/j.foodchem.2006.03.032

Richard-Forget, F.C., Gauillard, F.A., 1997. Oxidation of chlorogenic acid,
 catechins, and 4-methylcatechol in model solutions by combinations of pear (*Pyrus*
communis cv. Williams) polyphenol oxidase and peroxidase: a possible involvement
 of peroxidase in enzymatic browning. J. Agric. Food Chem. 45, 2472–2476.
 10.1021/jf970042f

Saba, M.K., Sogvar, O.B., 2016. Combination of carboxymethyl cellulose-based coatings with calcium and ascorbic acid impacts in browning and quality of fresh-cut apple. *LWT-Food Science and Technology* 66, 165–171. DOI: 10.1016/j.lwt.2015.10.022

Saltveit, M.E., 2000. Wound induced changes in phenolic metabolism and tissue browning are altered by heat shock. *Postharvest Biol. Technol.* 21, 61–69. DOI:10.1016/S0925-5214(00)00165-4

Saquet, A.A., Streif, J., Bangerth, F., 2003. Energy metabolism and membrane lipid alterations in relation to brown heart development in ‘Conference’ pears during delayed controlled atmosphere storage. *Postharvest Biol. Technol.* 30, 123–132. DOI:10.1016/S0925-5214(03)00099-1

Shrestha, L., Kulig, B., Moschetti, R., Massantini, R., Pawelzik, E., Hensel, O., Sturm, B., 2020. Optimisation of Physical and Chemical Treatments to Control Browning Development and Enzymatic Activity on Fresh-cut Apple Slices. *Foods* 9, 76. DOI:10.3390/foods9010076

Sun, Y., Zhang, W., Zeng, T., Nie, Q., Zhang, F., Zhu, L., 2015. Hydrogen sulfide inhibits enzymatic browning of fresh-cut lotus root slices by regulating phenolic metabolism. *Food Chem.* 177, 376–381. DOI:10.1016/j.foodchem.2015.01.065

Toivonen, P.M.A., Brummell, D.A., 2008. Biochemical bases of appearance and texture changes in fresh-cut fruit and vegetables. *Postharvest Biol. Technol.* 48, 1–14. DOI:10.1016/j.postharvbio.2007.09.004

Wang, 2002. Two’s company, three’s a crowd: can H₂S be the third endogenous gaseous transmitter? *FASEB Journal* 16, 1792–1798. DOI:org/10.1096/fj.02-0211hyp

Wang, J., Lv, M., Li, G., Jiang, Y., Fu, W., Zhang, L., Ji, S., 2018. Effect of intermittent warming on alleviation of peel browning of ‘Nanguo’ pears by regulation

energy and lipid metabolisms after cold storage. *Postharvest Biol. Technol.* 142, 99–106. DOI:10.1016/j.postharvbio.2017.12.007

Xu, C., Zeng, B., Huang, J., Huang, W., Liu, Y., 2015. Genome-wide transcriptome and expression profile analysis of *Phalaenopsis* during explant browning. *PloS one* 10, e0123356. DOI:10.1371/journal.pone.0123356

Yan, S., Luo, Y., Zhou, B., Ingram, D.T., 2017. Dual effectiveness of ascorbic acid and ethanol combined treatment to inhibit browning and inactivate pathogens on fresh-cut apple. *LWT-Food Sci. Technol.* 80, 311–320. DOI:10.1016/j.lwt.2017.02.021

Yin, Y., Bi, Y., Li, Y., Wang, Y., Wang, D., 2012. Use of thiamine for controlling *Alternaria alternata* postharvest rot in Asian pear (*Pyrus bretschneideri* Rehd. cv. Zaosu). *Int. J. Food Sci. Tech.* 47, 2190–2197. DOI:10.1111/j.1365-2621.2012.03088.x

Zhang, J., Yuan, L., Liu, W., Lin, Q., Wang, Z., Guan, W., 2017. Effects of UV-C on antioxidant capacity, antioxidant enzyme activity and colour of fresh-cut red cabbage during storage. *Int. J. Food Sci. Tech.* 52, 626–634. DOI:10.1111/ijfs.13315

Zhang, S., Lin, Y., Lin, H., Lin, Y., Chen, Y., Wang, H., Shi, J., Lin, Y., 2018. *Lasiodiplodia theobromae* (Pat.) Griff. & Maubl. -induced disease development and pericarp browning of harvested longan fruit in association with membrane lipids metabolism. *Food Chem.* 244, 93–101. DOI:10.1016/j.foodchem.2017.10.020

Zhang, S., Tian, L., Zhang, Y., Zhao, H., Zhao, J., Guo, J., Zhu, G.-P., 2019. *De novo* transcriptome assembly of the fresh-cut white husk of *Juglans cathayensis* Dode: insights for enzymatic browning mechanism of fresh-cut husk of walnut. *Sci. Horticult.* 257, 108654. 10.1016/j.scienta.2019.108654

815 Zheng, J.-L., Hu, L.-Y., Hu, K.-D., Wu, J., Yang, F., Zhang, H., 2016. Hydrogen
816 sulfide alleviates senescence of fresh-cut apple by regulating antioxidant defense
817 system and senescence-related gene expression. HortScience 51, 152–158.
818 DOI:10.21273/HORTSCI.51.2.152

819 Zhu, H., Liu, J., Wen, Q., Chen, M., Wang, B., Zhang, Q., Xue, Z., 2017. De
820 novo sequencing and analysis of the transcriptome during the browning of fresh-cut
821 *Luffa cylindrica* 'Fusi-3' fruits. PloS one 12. DOI:10.1371/journal.pone.0187117

Figures

Fig. 1 Effects of H₂S treatment on the (A) appearance and (B) browning index of fresh-cut apple during storage at 4 °C.

Fig. 2 Physiological changes of fresh-cut apple after H₂S treatment during storage at 4 °C. (A) PPO, (B) POD, (C) PAL, (D) SOD, (E) CAT, (F) APX and (G) GR activities, (H) H₂O₂ content, (I) AsA content, (J) phenolic content.

Fig. 3 Differentially Expressed Genes (DEGs) between samples. (A) Numbers of up- and down-regulated DEGs in S0 vs. C0, S6 vs. C6, C6 vs. C0 and S6 vs. S0. Venn diagrams showing common DEGs in S0 vs. C0 and S6 vs. C6 (B), C6 vs. C0 and S6 vs. S0 (C).

Fig. 4 The top 15 enriched GO terms ($P < 0.05$) of biological process, cellular component and molecular function in S0 vs. C0 (A), S6 vs. C6 (B), C6 vs. C0 (C) and S6 vs. S0 (D).

Fig. 5 Proposed model for the browning inhibition of fresh-cut apple by H₂S.

Fig. 6 Correlation analysis between RNA-seq and qRT-PCR data.

837 **Supplementary material**

838 Fig. S1 Gene expression pattern of the DEGs of the control and H₂S treatment at day
839 6 relative to day 0 in the common GO terms.

840 Table S1 Primers used in qRT-PCR for the validation of RNA-Seq data.

841 Table S2 Quality summary of sequencing data.

842 Table S3 The number of unigene and transcript identified in each sample.

843 Table S4 Uniquely expressed genes in each samples.

844 Table S5 Top 10 up- and down-regulated genes after H₂S treatment.

845 Table S6 KEGG pathway classification of the DEGs in S0 vs. C0, S6 vs. C6, C6 vs.
846 C0 and S6 vs. S0

847 Table S7 DEGs involved in fatty acid metabolism.

848 Table S8 DEGs involved in Glycerophospholipid metabolism.

849 Table S9 DEGs involved in oxidative phosphorylation.

850 Table S10 DEGs involved in glycolysis.

851 Table S11 DEGs involved in TCA.





Fig. 1 Effects of H₂S treatment on the (A) appearance and (B) browning index of fresh-cut apple during storage at 4 °C. The error bars represent standard deviation (SD, n = 3). *: statistically significant differences (P < 0.05).



5

6 Fig. 2 Physiological changes of fresh-cut apple after H_2S treatment during storage at
7 4 °C. (A) PPO, (B) POD, (C) PAL, (D) SOD, (E) CAT, (F) APX and (G) GR activities,
8 (H) H_2O_2 content, (I) AsA content, (J) phenolic content. The error bars represent
9 standard deviation (SD, n = 3). *: statistically significant differences ($P < 0.05$).







Fig. 3 Differentially Expressed Genes (DEGs) between samples. (A) Numbers of up- and down-regulated DEGs in S0 vs. C0, S6 vs. C6, C6 vs. C0 and S6 vs. S0. Venn diagrams showing common DEGs in (B) S0 vs. C0 and S6 vs. C6, (C) C6 vs. C0 and S6 vs. S0. DEGs were filtered with a cut-off of $|\log_2FC| \geq 1$ and $P < 0.05$.




16

17 Fig. 4 The top 15 enriched GO terms ($P < 0.05$) of biological process, cellular
 18 component and molecular function in (A) S0 vs. C0, (B) S6 vs. C6, (C) C6 vs. C0 and
 19 (D) S6 vs. S0.



23 means gene was not differentially expressed ($P \geq 0.05$ or $-1 < \log_2FC < 1$). 3DD,SD: bifunctional 3-dehydroquinase dehydratase/shikimate
 24 dehydrogenase, 4CL: 4-coumarate-CoA ligase, AA: aspartate aminotransferase, ACD: aconitate hydratase, ADT,PDT: arogenate
 25 dehydratase/prephenate dehydratase, AO: ascorbate oxidase, APX: ascorbate peroxidase, AsA: ascorbic acid, CAT: catalase, CM: chorismate
 26 mutase, CS: chorismate synthase, FK: fructokinase, FUH: fumarate hydratase, G3PD: glyceraldehyde-3-phosphate dehydrogenase, GST:
 27 glutathione S-transferase, HCT: shikimate O-hydroxycinnamoyltransferase, ICDH: isocitrate dehydrogenase, LOX: lipoxidase, LPD:
 28 phospholipase D, MDH: malate dehydrogenase, MDHAR: monodehydroascorbate reductase, NAD: NADH dehydrogenase, PAL: phenylalanine
 29 ammonia-lyase, PDH: pyruvate dehydrogenase, PFK: 6-phosphofructokinase, PGK: phosphoglycerate kinase, PK: pyruvate kinase, POD:
 30 peroxidase, PPH: phosphopyruvate hydratase, PPO: polyphenol oxidase, SOD: superoxide dismutase, SDH: succinate dehydrogenase, SUS:
 31 sucrose synthase, SK: shikimate kinase.



32

33 Fig. 6 Correlation analysis between RNA-seq and qRT-PCR data. Ten DEGs from the
 34 RNA-seq assay were used for qRT-PCR assay. The \log_2FC obtained by qRT-PCR
 35 (X-axis) was plotted against \log_2FC by RNA-seq (Y-axis).