

*An uncertainty-focused database
approach to extract spatiotemporal trends
from qualitative and discontinuous lake-
status histories*

Article

Accepted Version

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

de Cort, G., Chevallier, M., Burrough, S., Chen, C. and Harrison, S. P. ORCID: <https://orcid.org/0000-0001-5687-1903> (2021) An uncertainty-focused database approach to extract spatiotemporal trends from qualitative and discontinuous lake-status histories. *Quaternary Science Reviews*, 258. 106870. ISSN 0277-3791 doi: 10.1016/j.quascirev.2021.106870 Available at <https://centaur.reading.ac.uk/96508/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1016/j.quascirev.2021.106870>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

An uncertainty-focused database approach to extract spatiotemporal trends from qualitative and discontinuous lake-status histories

^{1,2,3*}De Cort, Gijs; ^{4,5}Chevalier, Manuel; ⁶Burrough, Sallie L.; ^{7,8,9}Chen, Christine Y.; ¹⁰Harrison, Sandy P.

¹Limnology Unit, Department of Biology, Ghent University, Belgium

²Division of Ocean and Climate Physics, Lamont-Doherty Earth Observatory, Columbia University, USA

³Department of Earth Sciences, Royal Museum for Central Africa, Belgium

⁴Institute of Earth Surface Dynamics, University of Lausanne, Switzerland

⁵Institute of Geosciences, Sect. Meteorology, Rheinische Friedrich-Wilhelms-Universität Bonn, Auf dem Hügel 20, 53121 Bonn, Germany

⁶School of Geography and the Environment, University of Oxford, UK

⁷Massachusetts Institute of Technology-Woods Hole Oceanographic Institution Joint Program in Oceanography, Cambridge, MA, USA

⁸Department of Earth, Atmospheric and Planetary Sciences, Massachusetts Institute of Technology, Cambridge, MA, USA

⁹Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, CA, USA

¹⁰School of Archaeology, Geography and Environmental Science, University of Reading, UK

*Corresponding author:

E-mail: gijs.decort@ugent.be

Current address:

Limnology Unit, Department of Biology, Ghent University

K.L. Ledeganckstraat 35, 9000 Ghent, Belgium

Abstract

Changes in lake status are often interpreted as palaeoclimate indicators due to their dependence on precipitation and evaporation. The Global Lake Status Database (GLSDB) has since long provided a standardised synopsis of qualitative lake status over the last 30,000 ^{14}C years. Potential sources of uncertainty however are not recorded in the GLSDB. Here we present an updated and improved relational-database framework that incorporates uncertainty in both chronology and the interpretation of palaeoenvironmental data. The database uses peer-reviewed palaeolimnological studies to produce a consensus on qualitative lake-status histories, whose chronologies are revised and standardized through the recalibration of radiocarbon dates and the application of Bayesian age-depth modelling for stratigraphic archives. Quantitative information on absolute water-level elevation is preserved if available from geomorphological sources. We also propose a new probabilistic analytical framework that accounts for these uncertainties to reconstruct synoptic, integrated environmental signals. The process is based on a Monte Carlo algorithm that iteratively samples individual lake-status histories within the limits of their uncertainties to produce many possible scenarios. We then use Recursively-Subtracted Empirical Orthogonal Function analysis to extract dominant patterns of lake-status variability from these scenarios.

As a proof of concept, we apply this framework to 67 sites in eastern and southern Africa whose lake-status histories cover part of the late Pleistocene and/or Holocene. We show that, despite the sometimes large temporal and interpretation uncertainties, and the inclusion of highly discontinuous lake-status time series, identifying the major known millennial-scale climatic phases during the last 20,000 years is possible. Our framework was also able to identify an antiphased response between the lake basins in eastern and interior southern Africa to these changes. We propose that our new database and methodology framework serves as a template for efficient lake-status data synthesis, encourages the incorporation of lake-status data in more palaeoclimate syntheses, and expands the possibilities for the use of such data in the evaluation of climate models.

Keywords (*max. 10*)

Lake status; lake level; palaeolimnology; palaeoclimatology; relational database; data uncertainty; Monte Carlo algorithm; Recursively-Subtracted Empirical Orthogonal Function; Africa; Quaternary

Highlights (*3-5*)

- We present a new database format to synthesize lake-status histories, conserving data uncertainty in both chronology and lake status.

- A combination of Monte Carlo iterative sampling and recursively-subtracted empirical orthogonal function analysis is employed to extract dominant patterns from uncertain lake-status histories.
- A proof-of-concept study highlights the dominant millennial-scale palaeoclimatic trends of eastern and southern Africa of the last 20,000 years.

1. Introduction

Understanding hydroclimate dynamics on timescales longer than a few decades depends largely on natural environmental archives that have recorded past climate changes. Changes in lake volume/area provide important information about past hydroclimates because they reflect changes in the balance of precipitation and evaporation over the lake and its catchment (Cheddadi et al., 1997; Harrison et al., 2002). The first efforts to reconstruct late-Quaternary palaeo-hydroclimate dynamics from lake data started in the late 1970's (Street and Grove, 1979) and led to the creation of several regional compilations (e.g., Harrison, 1989; Fang, 1991; Harrison, 1993; Harrison et al., 1993; Tarasov et al., 1994; Yu and Harrison, 1996; Harrison et al., 1996; Jolly et al., 1998). These regional databases were subsequently homogenised and integrated into the first Global Lake Status Database (GLSDB; Qin et al., 1998) that contained histories of 'lake status' (a term that covers all parameters of lake size, i.e. depth, level, area and volume) across the world spanning some or all of the last 30,000 radiocarbon years (~34,000 cal yr BP). Initially, the GLSDB records were categorised into three qualitative lake-status classes – 'low', 'intermediate' and 'high' – at 1000 radiocarbon-year intervals. Later, this structure was adapted to provide continuous records and allow an assessment of dating quality and flexible definitions of status class. The GLSDB has been used to assess palaeoenvironmental changes since the Last Glacial Maximum (LGM) and to evaluate climate-model hindcasting performance for key time windows in the past (e.g., Qin et al., 1998; Kohfeld and Harrison, 2000; Coe and Harrison, 2002; Tierney et al., 2011; Bartlein et al., 2017).

Unfortunately, the GLSDB was last updated in the late 1990s and does not include the many new palaeolimnological studies published since then. An update of the chronologies of the records stored in the database is also necessary because all the chronologies were reported as uncalibrated radiocarbon dates, and because many other geochronological methods have since been applied to palaeolimnological archives (e.g., Kutterolf et al., 2016; Roberts et al., 2018; Chen et al., 2020; Hatfield et al., 2020). In parallel, awareness has grown regarding the necessity to account for uncertainties when interpreting paleoenvironmental records and their age models (e.g., Telford et al., 2004; Heegaard et al., 2005; Blaauw, 2010; Blaauw and Christen, 2011; Parnell et al., 2011; Breitenbach 2012). In its original form, the GLSDB cannot account for such uncertainties. One particular consequence of such limited consideration of uncertainties is that all the records are treated as equivalent. Such an assumption is obviously problematic as some records are more informative than others, due to their better dating and/or to their more direct/simpler interpretations.

This paper presents a new relational database structure that efficiently stores lake-status histories, including both chronological and status uncertainties, and demonstrates how such a uncertainty-

focused database can enable more complex reconstructions of past environmental change. As a proof-of-concept, we present a new compilation of 67 late-Quaternary lake-status histories from eastern and southern Africa (ESA), many of which are discrete and discontinuous. We employ a new probabilistic approach based on a Monte Carlo (MC) algorithm and empirical orthogonal functions (EOFs) to identify the dominant hydroclimate features across ESA during the last 20,000 years. Finally, based on the results of this concept study, we explore and discuss the relative strengths and weaknesses of our framework.

2. Lake status as an indicator of past climate

The balance between precipitation and evaporation (P-E) exerts a universal control on lacustrine systems by regulating lake volume, which itself depends on lake depth and surface area (Mason et al., 1994; Cheddadi et al., 1996; Harrison et al., 2002). Responses of lake volume to climatic change are most striking in endorheic, or ‘closed’ lakes, i.e. lakes without an (above-ground) outflow where most or all water leaving the lake is lost through evaporation. Closed lakes are found most often in semi-arid environments, where catchment evapotranspiration exceeds water input, thus constraining lake-level equilibria below overflow level (Meybeck, 1995). While the response of exorheic, or ‘open’ lakes, to climatic change is often less dramatic, these also undergo lake-level changes if changes in the hydrologic budget are not compensated by inflow/outflow adjustments (Cheddadi et al., 1996). On longer time scales, many lakes can be classified as transitional, shifting between open and closed states as changing climate controls their water balance (Kalff, 2001).

Past lake status is derived from two main lines of evidence. The first source is geomorphic features formed by marginal processes that reflect displacements of the past shore or near-shore environment (‘palaeoshorelines’). Palaeoshorelines above the present water level can be dated using radiometric or luminescence dating techniques to estimate the timing at which they were last active. Combined with elevation data and knowledge of basin morphometry – and if necessary corrected for post-depositional deformation that might occur through tectonic vertical displacement (e.g., Garcin et al., 2009) or isostatic adjustments of the lithosphere (e.g., Chen and Maloof, 2017) – they allow for a direct, quantitative reconstruction of water level and lake hydrology at their time of formation. Palaeoshorelines represent discrete, rather than continuous, recordings of lake-level history as they are often only deposited and preserved under specific hydrological conditions such as water-level stasis. As such, they only provide snapshots of past hydrological states. Palaeoshoreline records tend to dominate the palaeolimnological record in semi-arid to arid regions, where the climate regime necessary to maintain steady state lake-full conditions deviates most strongly from present-day climate conditions (Bowler, 1986; Burrough and Thomas, 2009).

The second source of evidence is derived from the lakebed, where the progressive deposition of sediments results in sequences representative of the character of the overlying water column through time. Such stratigraphies can be retrieved as sediment cores or as outcropping sections of sedimentary landforms above the modern water surface. Lakebed archives can be (but are not necessarily) uninterrupted deposits and thus provide a continuous time series of past lake status. Absolute reconstructions of water depth can be made from lake-bed records using sequences of cores to trace the limit of lacustrine deposition through time (e.g., Digerfeldt, 1965; Schneider Tobolski, 1985; Winkler et al., 1986; Almqvist-Jacobsen, 1995; Shuman et al., 2009) but the procedure is time-consuming. Generally lake-bed records only provide indirect information of relative changes in lake status approximated through sensors of water depth such as sedimentological characteristics or floral and faunal community composition. For these reasons, lake-level reconstructions derived from lakebed archives tend to be continuous but qualitative in nature.

Reconstructing lake status and associated climatic change from sedimentology and geomorphology can be challenging for several reasons (e.g., Gasse, 2000; Holmes and Hoelzmann, 2017):

1. Like all natural palaeoenvironmental archives, the chronological control on lake records is dependent on the dating method applied. Apart from lakes containing annually varved sediments, dating of lacustrine deposits relies mostly on radiometric techniques with varying precision.
2. Lake-status archives can be incomplete and low stands are often underrepresented. In sedimentological records, low water levels or dry stands can inhibit deposition and even remove previously deposited sediments, thus creating stratigraphic hiatuses or unconformities. Similarly, palaeoshorelines above a lakebed only reflect instances where water level was higher than today, and prolonged high stands can erode previously formed shoreline structures thereby eliminating evidence for older episodes of lower water level (Burrough and Thomas, 2009).
3. The information about lake-status changes tends to decrease further back in time, which can be partly explained by practical fieldwork constraints (i.e., coring older deposits is technically challenging and costly) but also reflects difficulties of obtaining records after lake desiccation because of compaction, pedogenesis and/or mineralisation of the lakebed (e.g., Stager et al., 2002; Bessems et al., 2008; De Cort et al., 2018). Lake coring is therefore often restricted to the time window following the most recent dry stand. The result is that lake-status data is often skewed towards recent periods of higher water level, which have a higher probability of being recorded than periods of greater age and/or lower lake status.

4. The sensitivity of lake level to climate is site-specific, as factors such as physical and ecological catchment characteristics and groundwater inflow or outflow can affect the amplitude and timing of lake response to variability in P-E (Vassiljev et al. 1998). Different archives also show different levels of sensitivity to lake-status changes, which may differ between indicators but also be non-stationary through both space and time for the same indicator (Juggins, 2013).
5. Lake-status changes can also be driven by non-climatic factors, such as gradual basin infilling, basin deformation (e.g. by tectonism or volcanism) or sea-level change. These processes can mask or modify the climate signal preserved in sedimentary or geomorphic archives.

Lakes vary widely in their climate sensitivity. Most lakes act as low-pass filters that mainly respond to lower-frequency climate variability (Mason et al., 1994; Liu and Schwartz, 2014). Catchment size and morphometry, as well as relative groundwater flux, are major factors in determining the sensitivity of lake level to climate change (Vassiljev et al. 1998; Burrough and Thomas, 2009; Olaka et al., 2010). The time needed for a lake to return to equilibrium creates a lagged response which usually varies from months to a few years (Yi and Zhang, 2015) but can take multiple millennia in extremely large catchments (Singarayer et al., 2019).

3. The new relational lake-status database

This section describes the principles behind the construction of the new database, as applied to the ESA sites used in our case study (see section 5).

3.1 Data selection

Studies containing data on past lake-status change were included in the database if they fulfilled the following criteria:

1. Data and publications were peer-reviewed;
2. Chronological data was available to anchor past lake variability in time (e.g., radiometric dating, reliable historical accounts or instrumental records);
3. Evidence indicates that lake-level changes were primarily driven by climate and were interpreted as such in the original literature. Sites or sections of individual records where changes in lake level were demonstrably non-climatic in origin (e.g., direct influence of sea-level changes, volcanism, basin infilling or anthropogenic activities) were excluded to avoid mixing environmental signals;
4. The palaeoenvironmental evidence reflected changes in lake level, depth, volume and/or area rather than other hydrological variables such as precipitation, runoff, or moisture source.

Interpretations from the original literature were followed in this matter, unless competing evidence (e.g., from contradictory studies) demanded otherwise.

3.2 Compilation and revision of chronological information

A wide range of approaches for age-model construction were used in the original literature, so we built standardised chronologies for all records. Radiocarbon dates were recalibrated using IntCal13 for Northern Hemisphere sites (Reimer et al., 2013) and SHCal13 for Southern Hemisphere sites (Hogg et al., 2013). Age-depth models for sediment profiles were reconstructed using the Bayesian age-depth modelling software Bacon in R (Blaauw and Christen, 2011; R Core Team, 2020). Core-top ages were included in the age model when intact recovery of the surface sediment was reported, in which case the core top was assumed to correspond to the time of sampling. The specifics of all the age-depth models, such as the exclusion of anomalous outlying dates, the excision of event deposits, the inclusion of sediment hiatuses, and the settings of the Bacon run were adjusted on an *ad-hoc* basis for each individual sequence to ensure the most reliable results and with the objective of staying as close as possible to interpretations in the original literature.

The discrete nature of palaeoshorelines means that calibrated ages obtained from them were used directly as a shoreline age, except in the case where multiple shoreline dates overlapped within their uncertainty ranges to such a degree that they could be interpreted as representing a single continuous time interval for a specific lake status.

New chronologies were not created for sites where i) the original chronological data was (partly) unavailable but the site was considered fundamental for regional reconstruction, or ii) where the published age model was already created using Bacon with the appropriate calibration curve. Original ¹⁴C measurements and associated information, as well as data from other geochronological techniques, were retained in the database to enable future updates.

3.3 Construction of consensus lake-status histories

Many lines of evidence only provide indirect evidence of past lake status. Thus, assumptions on the relationship between such proxies and lake level are often required to reconstruct lake status. Our standard approach was to combine all available sources of information on the assumption that if several lines of evidence point to a change in water depth or area, this consensus is real (Harrison et al., 1991; Harrison and Digerfeldt, 1993). For every site, the information on past changes in lake status was combined with the revised chronologies to produce sequences of lake-status episodes. Each episode is defined by i) a best-estimate of the start and end age, each with their associated 2- σ credible intervals, taken from the reconstructed age-depth models of sediment cores (Fig. 1a) or

individual/clustered ages from discontinuous archives (Fig. 1b); and ii) a relative qualitative lake-status class represented by a positive integer, in which a higher status (higher number) signifies a higher relative lake level, volume or surface area (1 being the lowest status documented and not necessarily meaning drystand; Fig. 1). The weighted mean of the probability distribution produced by the Bacon model, or in the case of palaeoshorelines, of the single defining chronological point, was taken as the best estimate of the 'true' age.

There is no upper limit to the number of different lake-status classes assigned to a lake, as this depends on the available lines of evidence and their sensitivity to lake-status change, the resolution of the available information and the total time span of the archive. The ranking is specific to each lake and status classes cannot be directly compared between lakes. In general, the original authors' interpretation was followed unless there were contrasting conclusions published in the literature. When the status at a particular time relative to other periods in a lake's history was ambiguous, lower and upper limits on the possible lake status class were defined. This occurred when status was based on ambiguous data interpretation, or when several archives were used that do not overlap sufficiently in time or type of evidence so that comparison of status in different periods of a lake's history was not straightforward.

Assignment of lake-status classes to lakebed sediments and palaeoshorelines was challenging in different ways. The density of palaeoshoreline dates was often insufficient to create continuous reconstructions. Additional assumptions were often required concerning the duration of events represented by features dated by only one or a few ages. The same was true for other discontinuous records such as non-shoreline geomorphological features or archaeological artefacts. With such 'snapshot' information, event duration had to be estimated conservatively, informed by the nature of the deposit and the sensitivity of the basin to hydrological changes. For example, raised beaches may be assigned shorter durations (on the order of years) than thick carbonate deposits that accumulated in the littoral zone over multiple decades or centuries. Similarly, some lake basins are well buffered against short-term water-level fluctuations whereas others exhibit significant variability on (sub-)annual time scales, depending on basin characteristics such as morphometry, hydrography and vegetation.

3.4 Database structure and content

Data on site characteristics, chronology and lake-status history was assembled into 9 tables, which were combined into a relational database and exported as a SQLite3 file (Fig. 2, SI).

Lakes – This table contains a unique site identifier, site name, coordinates and basic information on present hydrology and morphometry.

Contributors – This table includes the name and ORCID of the researcher who contributed the site history to the database, and the date (month/year) when the data were included in the database.

Alternative Names – This table links the site identifier with alternative site names used in the literature.

Refs – This table contains all the references used to compile the database.

Dating – This table contains the original individual dates as well as the outcome of their reanalysis as part of the newly generated chronologies.

Coding Basis – This table contains the site-specific definitions of all status classes.

Coding Source – This table lists which lake-status indicators were available in the consulted literature, and which of these were used in the consensus lake-status reconstruction.

Lake Size – This table provides information on absolute water depth or lake-level elevation as recorded by geomorphological features. Best estimates of the mean, lower and upper still-water bound of each shoreline, and corresponding estimates and uncertainties of lake surface area and lake volume, are given if available from the original literature.

Coding – This table contains the lake-status histories. For each site, a sequence of episodes is defined. Each of these episodes is defined by a best-estimate start and end age, together with their associated 2σ uncertainties, and by a best estimate, a minimum and a maximum value for lake status. The Coding table also links lake-status episodes to corresponding quantitative information, if available in the Lake Size table.

A number of additional tables is used to store lists of accepted values for specific fields or to link the different data tables together (Fig. 2, SI).

4. Analytical approach

Integrating temporal and lake-status uncertainties allows for the extraction of robust spatiotemporal palaeoclimatic trends from data (e.g., Breitenbach et al. 2012; Anchukaitis and Tierney, 2013; Chevalier and Chase 2015). Here, we designed a specific Monte-Carlo (MC) framework to iteratively sample the lake-status histories, taking account of their uncertainties in chronology and lake status, to produce a large number of possible historical scenarios (section 4.1). These randomised iterations

were subsequently used to extract the principal spatiotemporal modes of variability (section 4.2). A flowchart of this analytical approach is shown in Fig. 3.

4.1 Monte-Carlo sampling of time- and status-uncertain lake histories

Our MC algorithm iteratively sampled the Coding table to produce a large number of variants of each lake's status history; in our application (see section 5), we used 10,000 iterations. In each iteration, uncertainty ranges of start and end ages of subsequent status episodes were sampled unidirectionally through time to conserve the chronological sequence of defined episodes. The outcome of the sampling of a given time point (i.e. the start or end of a certain status episode) was thus constrained by the uncertainty defined for that point and by the result of the priorly sampled point. To avoid biasing the sampled records either towards younger or towards older ages – which can happen especially for sequences of relatively short episodes with highly overlapping temporal uncertainty ranges – the sampling was randomly performed by either moving forward or backward in time. This means that for any lake, roughly half of the iterations sampled the coding table in an old-to-young sequence and the other half the iterations sampled the coding table starting at the youngest and ending at the oldest episode. Similar to the sampling of the start and end dates of each episode, the status class was also sampled from its uncertainty range but without being constrained by the priorly sampled point, since we assumed that status class is independent between episodes.

Although the distribution of estimated ages produced by radiocarbon calibration or Bayesian age-depth modelling are generally not symmetric or even unimodal, only the model weighted means and upper and lower 95% confidence intervals are used to define the sequence of episodes that make up a lake's history in the database. Similarly, probability distributions of status classes are also not normally distributed and may be highly asymmetrical (see Fig. 1c). To cope with the asymmetrical nature of age uncertainties, the MC algorithm samples start and end ages of status episodes following a probability density function defined by a split-normal distribution, also known as a two-piece normal distribution. The mode of the split-normal distribution is equal to the best estimate of that age (Fig. 1c) and the left and right standard deviations (1σ) of the distribution are derived from the lower and upper limit of the 2σ range as defined in the coding table. Uncertain status classes were set to follow a triangular probability density function, peaking at best-estimate values and linearly decreasing towards 0 at values of minimum-1 and maximum+1, which precludes obtaining values outside of the defined minimum-to-maximum range (Fig. 1c). The split-normal (for ages) and triangular (for status class) distributions are scaled to ensure that their cumulated probabilities sum to 1.

4.2 Recursively-Subtracted Empirical Orthogonal Function analysis

To extract regional palaeoclimatic trends, MC-sampled lake-history variants were generated for a set of lakes of interest. These histories were then analysed using Empirical Orthogonal Functions (EOFs), which decompose data of high spatiotemporal dimensionality into its dominant spatiotemporal features of variability. Recursively-Subtracted Empirical Orthogonal Function (RSEOF) analysis is a specific case of EOF analysis that is able to deal with data sets with high proportions of missing values (Taylor et al., 2013), as can be the case with lake-status data. All MC-produced lake histories were interpolated to a common time scale with regular spacing. In our proof-of-concept study using African lakes (see Section 5), a time step of 5 years was employed. Time series were centred and scaled to a mean of 0 and a standard deviation of 1 prior to the analysis. A reference RSEOF was also calculated for the scenario where all ages and statuses of included sites were assumed at their best-estimate value. The result was an ensemble of 10,000 (MC-generated) + 1 (best-estimate) sets of principal components (PCs), each summarizing the dominant spatiotemporal modes of variability in their corresponding set of lake-status histories. By construction, the sign of each eigenvector is arbitrary (Legendre & Legendre, 2012). To homogenise the sign of corresponding eigenvectors across all RSEOF iterations, each PC was multiplied by -1 if this resulted in a reduction of its distance to the corresponding PC of the set of best-estimate lake-status histories, *i.e.* the sign and direction of each eigenvector was aligned with the sign and direction of the eigenvector of the reference RSEOF. For each RSEOF iteration, unicity of EOF modes was assessed using North's Rule of Thumb, which determines which eigenvectors are distinct from their nearest neighbour (North, 1982). RSEOF analyses were carried out using the R package *sinkr* (Taylor, 2017; R Core Team, 2020).

5. Case study: lake-status history in eastern and southern Africa

5.1 Regional setting

Our study area – eastern and southern Africa (ESA) – is delimited by the Sahel to the north and by the Congo Basin and the Angolan highlands to the west (Fig. 4) and encompasses different climatic zones (Gasse et al., 2008; Burrough and Thomas, 2013). Tropical East Africa and Madagascar mainly receive monsoonal rainfall from the Indian Ocean associated with the yearly migration of the tropical rain belt, although Atlantic moisture can also significantly contribute in western areas (Nicholson, 2017; 2018). In southern Africa, the strength of the monsoon decreases with distance from the Indian Ocean, grading from the subtropical grasslands and savannas in the east into the arid Kalahari over the southern African plateau and finally into the hyper-arid Namib Desert along the west coast. This pattern is additionally influenced by the cold Benguela current in the Atlantic, further depriving the

Namib of rain. The southern and southwestern coasts of South Africa are dominated by a Mediterranean climate, where winter rains prevail (Tyson and Preston-White 2000).

Lake sediments are the dominant source of palaeoenvironmental and palaeoclimatic reconstructions for East Africa (Verschuren, 2003). Tectonic activity and faulting associated with the East-African Rift System (EARS) have produced a string of basins which are arranged in two distinct rift branches (Chorowicz, 2005). The western branch, or Albertine Rift, holds several large and deep freshwater lakes (Tiercelin and Lezzar, 2002). Lakes of the eastern branch, or Gregory Rift, are typically smaller, shallower and more saline (Schagerl and Renaut, 2016). Additionally, volcanic activity associated with the EARS created a number of explosion craters, many of which hold small but sometimes relatively deep lakes. High densities of these crater lakes are found for example in the Rungwe Volcanic Province of southern Tanzania (Delalande et al., 2008) and the maar crater lake districts of southwestern Uganda (Melack, 1978). Locally concentrated small lakes of glacial origin are found in the alpine zones of the highest peaks and mountain ranges of East Africa (e.g., Mahaney, 2004; Eggermont and Verschuren, 2007; Eggermont et al., 2007; Tiercelin et al., 2008).

Because of southern Africa's dry conditions; a deep covering of sand related to its cratonic origins (Haddon and McCarthy, 2005); and the restricted influence of rifting, the southern African interior has less potential for deep lakes than East Africa. Today, most lakes and wetlands are found along the coast of South Africa, where they are often under significant marine influence (Hill, 1975; Whitfield et al., 2017). Nevertheless, the subcontinental interior holds many endorheic basins of various sizes, most of which are perennially dry or hold seasonally filled playas today, but where geomorphological features provide evidence for long-lived standing water in the past.

5.2 The ESA lake-status database

We identified a total of 67 lakes in ESA that met the criteria defined in Sect. 3.1 (Fig. 4, Table 1). Sites from the original GLSDB were only incorporated here if they met these criteria, and their GLSDB histories were critically re-evaluated in the process. Data for these sites was sourced from 244 publications (see Supplementary Material). Their catchments were delineated using the HydroSHEDS database (Lehner et al., 2008) or, for the smallest drainage basins, by manual delineation using Google satellite imagery and Advanced Spaceborne Thermal Emission and Reflection Radiometer (ASTER) Global Digital Elevation Model (GDEM) v2 data (ASTER GDEM is a product of METI and NASA). The ESA database is available online, along with supplementary documentation for each individual site (see Data availability).

The current nature of the lakes in these basins reflects regional climate. Most basins are topographically closed today (54/67). Seasonally or perennially dry basins are disproportionately common in the interior and western portions of southern Africa, while 10 of the 13 currently overflowing lakes are located within 12° of the Equator. Slightly more than half the sites (39/67) occur below 1000 m a.s.l., while 4/67 of sites lie above 2000 m asl. Most lakes have a surface area smaller than 100 km² (50/67) while 9/67 are currently larger than 1000 km². Catchment area ranges from < 1 km² for the smallest crater lakes to 1,205,959 km² for the Makgadikgadi basin (Fig. 5a). The most common sources of evidence for lake status (Table 1) are stratigraphy (used in 36/67 of all sites) and sedimentological composition of deposits (34/67), diatoms (32/67), non-shoreline geomorphology (27/67) and palaeoshorelines (21/67). Historical and/or instrumental gauge data was used for 12/67 and 8/67 of lakes, respectively. The number of defined status classes per site ranged from 2 to 15.

The chronological data is summarised in Fig 5. The database contains 1856 published chronological points, of which 72% are radiocarbon dates. A significant portion of radiocarbon dates (29%) had not been calibrated in the original publication. Optically stimulated luminescence (14% of the total), U/Th (1%) and thermoluminescence (1%) dates were almost exclusively obtained from non-core geomorphic features. Only 1460 chronological points (79% of the total) were used to establish the updated chronologies. Excluded dates either were demonstrated or suspected to be erroneous, or were from archives that were not used in the final consensus lake-status history. Chronologies were reconstructed for all except 2 sites where chronological data were unavailable and 4 sites where the published age model was already created using Bacon with appropriate radiocarbon calibration. Only 3 sites have lake-status information at 50,000 cal yr BP. This number increases to 16 at 20,000 cal yr BP, 27 at 10,000 cal yr BP, and 36 at 1000 cal yr BP. The average resolution of revised chronologies over the last 20,000 years, here defined as the number of used dates divided by the total duration of the record, is 4.1 (min 0.03, max 38.4) dates per 1000 years. For sites that go back to at least 20,000 cal yr BP, average dating resolution over the period 20,000 cal yr BP to 2020 CE is 1.3 (min 0.0, max 8.3, n = 26) dates per 1000 years. Similarly, over the period 1000 cal yr BP to CE 2020 it is 5.3 (min 0.0, max 57, n = 58) dates per 1000 years.

The number of informative sites decreases with age (Figs. 5b, 6, cf. Section 2). The varying availability of palaeolimnological records through time in ESA is partly the result of the lake-level changes these records document. For example, the steep decrease in the number of sites with lake-status information before ca. 2000 cal yr BP (Fig. 7) can be traced at least in part to the many relatively small and shallow lakes in East Africa that desiccate relatively regularly and therefore do not contain archives older than a couple of millennia (Verschuren, 2003). This bias in archive development by local climate is inherent to palaeolimnology in arid to semi-arid climate zones.

Although long histories are available from catchments of all sizes, there is a positive correlation between catchment size and the age of the oldest available information on lake status (Fig. 5a). Large catchments are generally associated with a higher variety in depositional environments, producing a diversity in palaeolimnological evidence that is more likely to survive for extended periods of time. Larger basins may also get more attention from the scientific community, resulting in a higher likelihood of deep-drilling campaigns or intensive geomorphological surveys. The largest catchments in the database are of such dimensions that their water levels are no longer driven solely by local climate but follow an integrated response to distinct climate patterns within several climate zones (e.g., Burrough et al., 2009). On average more status classes were defined for long histories from large catchments (Fig. 5a). This might reflect a higher sensitivity to moisture-balance change that comes with large, climatologically and ecologically diverse catchments and higher catchment to lake area ratios (Street, 1980; Olaka et al., 2010), but it is also likely that longer records contain evidence of higher-amplitude climate variability justifying more status classes. Shorter records typically have more status classes defined per unit of time (not shown) and the number of defined status episodes increases closer to the present (Fig. 5b), reflecting the different resolutions typically employed when studying archives of different length.

5.3 Application of joint MC-RSEOF analysis

We applied our joint MC-RSEOF framework (Fig. 3) to extract the dominant histories of the studied lakes across ESA during the past 20,000 years. Very few sites completely cover this time window: some sites have continuous histories but do not extend back to 20,000 years, others' histories are fragmentary across that time window. The RSEOF method allows for incomplete datasets, but the meaning of the PCs is directly related to the amount of information available in the system. Temporal coverage determines how complete a certain lake-status history is over the studied period, while the number of status classes reflects the sensitivity of a lake to undergo significant change and to record that change. The number of included lakes is also likely to have an influence on the PCs.

To determine to what extent the outcomes of our analyses are affected by temporal coverage, sensitivity to environmental change, data resolution and the number of the included lake-status records, we investigated different combinations of these site-selection criteria. Temporal coverage of the 20,000-0 cal yr BP window was set to range between 10 and 100% and the minimum required number of status episodes covering (part of) that window was set to 2 or 4, which resulted in the number of sites meeting these criteria, and thus being retained for subsequent RSEOF analyses, varying between 7 and 45 (Fig. 8a).

5.4 Millennial-scale lake-status variability in ESA over the last 20,000 years

The number of non-overlapping (i.e., distinct from neighbouring) EOFs and the amount of variance explained by the first EOF modes varied as different criteria for site inclusion were employed (Fig. 8a). Within this range of experiments, best-estimate scenarios result in 0 to 2 distinct (non-overlapping with their nearest neighbour) EOFs, which increases up to 4 in a minority of MC-produced scenarios. Distinct EOFs emerge when the coverage threshold is intermediate, as low temporal coverage produces too sparsely populated covariance matrices and results in gamma amplification due to lack of information (Taylor et al., 2013), while high temporal coverage excludes too many sites from the analysis. This may also explain why the amount of variance explained by EOF1 does not change significantly when the time-coverage criterion is changed between 33 and 100 %, i.e. the advantage of using more complete records is balanced by the reduction in their number. The time series of EOF1 is relatively consistent across tested subsets, despite the EOFs being based on a variable number of lake sites (Fig. 8b). This supports the notion that EOF1 reflects lake-status behaviour that is relatively consistent across the study region.

Fig. 9 highlights the RSEOF results for the subset of sites that cover at least 33% of 20,000-0 cal BP and for which at least 4 status episodes were defined during this time window. The scree plot (Fig. 9b) shows eigenvalues levelling from EOF3 onward, and North's rule of thumb identifies only the first 2 EOFs as non-overlapping. The first EOF explains $40.3 \pm 3.8\%$ of the variability in lake status across the MC ensemble of 10,000 simulations, or 43.6% in the best-estimates scenario. When interpreted as a trajectory of lake status, sites that score positively on this EOF tend to have low lake status at 20,000 cal yr BP, culminating in a minimum around c. 17,500-16,500 cal yr BP (Fig. 9c, d). Afterwards, lake status increases towards the start of the Holocene, although with a temporary reversal of this trend between c. 13,000-12,000 cal yr BP. A prolonged maximum in lake status is registered between c. 11,000-5,000 cal yr BP. After c. 4,000 cal yr BP, levels are intermediate between those of the start of the record and those of the early Holocene. All the sites that correlate positively to EOF1 are located in the EARS and Madagascar (Fig. 9a, c). Sites negatively correlated to EOF1 (i.e. showing opposite trends in lake status) are confined to interior and western southern Africa, with the exception of Lake Masoko in the southern Rift. The EOF biplot also shows the distinction between interior southern Africa and East Africa along the first axis, while homogeneous distribution among negatively scoring subregions reveals little to no spatial pattern within the EARS (Fig. 9a). The consistency of EOF1 across experiments with differential site selection (Fig. 8b), as well as the relatively narrow 95-% uncertainty envelope that is similar to EOF1 of the best-estimate lake histories (Fig. 9d), suggest that the dominant millennial-scale lake-status patterns in ESA over the last 20,000 years are widely represented and robust against the combined influence of chronological and status uncertainties. The first EOF shows

that lake status variability over the last 20,000 years is dominated by a coherent pattern of regionally opposed behaviour between the EARS and interior southern Africa.

The late-Glacial portion of EOF1 is remarkably similar to the climatic trend of the deglaciation in Africa according to transient palaeoclimate model simulations (Otto-Bliesner et al., 2014) and important palaeoclimate records from Africa (e.g., deMenocal 2000; Weldeab et al., 2007; Tierney and deMenocal 2013). Heinrich Stadial 1, which has been described as an extreme climatic event for the Afro-Asian monsoon region (Stager et al., 2011), is associated with minimum lake status across the EARS and high lake status in interior southern Africa between ca. 18,000-16,000 cal yr BP. From ca. 15,000 cal yr BP, lake-status EOF1 shows the increase of monsoonal precipitation over East Africa as a possible response to rising greenhouse-gas concentrations (Otto-Bliesner et al., 2014). This evolution is temporarily interrupted during the Younger Dryas (YD), a global event that is also captured in our EOF1 (Rasmussen et al., 2006; Carlson, 2013). With their almost uniformly positive loading on EOF1, lakes within the EARS had a high status during the early to mid-Holocene, concurrent with the widespread African Humid Period (Jolly et al., 1998; deMenocal et al., 2000; Shanahan et al., 2015). In contrast to northern Africa, where a similar early- to mid-Holocene humidity maximum was induced by the precessional expansion of the West-African monsoon (Kutzbach and Otto-Bliesner, 1982; Otto-Bliesner et al., 2014), the causes for tropical East Africa's similar moisture-balance behaviour during this episode remain unclear (Tierney et al., 2011; Liu et al., 2017; Reid et al., 2019; Wang et al., 2019).

The number of informative sites through time reflects the trends reconstructed by EOF1 (Fig. 7). The maximum of informative Kalahari sites is observed between 18,500-15,500 cal yr BP, as well as prolonged recording in the Namib between 17,000-11,000 cal yr BP, meaning that shorelines formed across the region during these times due to increased lake status. Afterwards, Holocene palaeoshorelines are less common across the interior of southern Africa, except for a short-lived increase in Kalahari sites around 6000-5500 cal yr BP. The aridification of southern-African drylands, mostly reflected in our database through absence of lake-status evidence, has been reconstructed from other data sources (Goudie and Thomas, 1986; Telfer and Thomas, 2006; Thomas, 2016, Chase et al. 2019). In contrast, the number of records from the EARS increases as lakes recovered from widespread lowstands or desiccation during HS1 (Stager et al., 2011). High numbers of sites reflect maximum positive moisture balance during the early Holocene, before a slight decrease during the mid Holocene and a final rise towards maximum numbers from ca. 4000 cal yr BP to the present.

More caution is necessary when interpreting higher-order EOFs of uncertain time series (e.g., Anchukaitis and Tierney, 2013). While significantly different from its nearest neighbours, EOF2 has large uncertainties and a low explanatory power (Fig. 9f), explaining $19.8 \pm 1.9\%$ of the total variance

across all MC iterations, and 18.7% in the case when only best-estimate lake-status histories are used. The main difference between EOF2 and EOF1 time series is the Holocene section, where EOF2 values for the early to mid-Holocene are similar to those of the preceding millennia, and the last 4,000 years show the most negative values of the entire period (Fig. 9f). The uncertainty range of EOF2 is considerably wider than the uncertainty range of EOF1. While sites south of 10°S load positively on EOF2, there is no evident spatial pattern for the EARS (Fig. 9a, e). We were not able to identify a mechanism for this trend with such a scatter of positive and negative loadings across the EARS. However, its general trend and consistent correlation to sites across the Kalahari could support the general evolution towards drier conditions over the last 20,000 years.

The trends and events identified in ESA have been described in earlier studies based either on individual records or from syntheses of selected palaeoclimate records (e.g., Chase et al., 2010; Stager et al., 2011; Stone, 2014; Singarayer and Burrough, 2015). While this paper does not aim at exploring the causal climatological processes behind them, the coherence of our results with well-established climatic events as evident in independent records demonstrates that robust palaeoclimate trends can be identified from qualitative and discontinuous lake-status data when age-model and interpretation uncertainties are taken into account.

6. Discussion and recommendations

Any collection of palaeoenvironmental data is confronted with what can be called a ‘synthesis dilemma’: the choice between targeting a low number of high-quality records at the expense of spatial and/or temporal coverage, or including as much data as possible, thereby increasing the internal noise of the data compilation due to lower average data quality. For the former, the curation of records often requires subjective interpretations of what is considered ‘high-quality’, which is not without its own limitations. The lake-status synthesis method presented in this paper adopts a more conservative approach and includes as many lakes as possible. Lake-status histories are often fragmentary and of variable temporal resolution, making them challenging to incorporate in palaeoenvironmental syntheses that often focus on selected archives or records of higher continuity and/or resolution (e.g., Tierney et al., 2013; McKay and Kaufman, 2014; Steiger et al., 2018; Atsawawaranunt et al., 2019). Our ESA case study nevertheless demonstrates that robust palaeoclimate patterns can still be extracted when the proxy and dating uncertainties are taken into account, even from collections of lake-status histories containing highly discontinuous and uncertain time series. Most interestingly, our final product derived from our MC-RSEOF approach has relatively narrow uncertainties – compared to the uncertainties of the composing records (see Fig. 1), thus highlighting its capacity to extract reliable signals, even from noisy data.

Uncertainty in relative lake status can have different causes, such as poorly documented sedimentation processes, uncertain translation of biological indicators to environmental change, and uncertainties comparing environmental indicators, especially when different indicators are used for different parts of a record. Consideration of uncertainty in the interpretation of a record is rarely taken into account in palaeoclimate-data syntheses, but can be important (e.g., Lohmann et al., 2013). The definition of minimum and maximum possible lake status, which together with the best estimate describe a triangular probability density function, is a practical solution to such challenges.

The incorporation of temporal uncertainty is crucial to assess past environmental variability (McKay and Kaufman, 2014). For example, Comas-Bru et al. (2019) explored the influence of chronological uncertainty in matching speleothem isotope data to climate model simulations over time windows of variable length. According to Anchukaitis and Tierney (2013), the instability introduced by temporal uncertainty is one of the main limits to recovering higher-order modes of variability from MC-sampled uncertain time series. When using our algorithm to study a fixed-boundary time window, chronological uncertainty not only affects the temporal data distribution but also the number of sites providing those data. In practice, the number of sites meeting a certain temporal-coverage criterion varies considerably between the 10,000 iterations of our last-20k experiments (Fig. 8a). While EOF1 exhibits clear similarities with the well-known succession of events that characterise the last 20,000 years, the meaning of EOF2 was less clear, and may support the results of Anchukaitis and Tierney (2013). Because our MC-RSEOF approach simultaneously integrates chronological and lake-status uncertainty, the uncertainty of its product is influenced by both factors. Since the lake-status classification system of the database is qualitative, all derived products should be treated as qualitative as well. However, quantitative information is available for a subset of the lakes of our database (Table 1) for more specific studies on climate-driven water budget.

In addition to the proposed methodology, our database format also offers major improvements in terms of uniformity and precision of lake-status chronologies over the GLSDB. The preservation of relevant information in a relational database format enables exploring and selecting sites based on sensitivity or temporal coverage, or other characteristics such as catchment size or type of chronological data. While we primarily focused on the past 20,000 years, different site selection based on density or other aspects of dating control could be advisable when studying shorter time scales or time windows. Similarly, caution is advised when studying poorly represented time windows or regions, where inclusion or exclusion of individual sites may significantly alter the resulting EOFs.

Our database format offers major improvements in terms of uniformity and precision of lake-status chronologies over the GLSDB. Almost one third of all reported radiocarbon dates were not calibrated

upon original publication (Fig. 6). Our updated chronologies commonly resulted in significant changes in comparison to their original counterparts. An important factor in radiocarbon-based chronologies is the dated material and, correspondingly, old-carbon effects. While our data compilation conserves information on reported carbon-reservoir effects, these effects are often not investigated in the original studies, which may cast additional uncertainty even on relatively high-resolution chronologies.

Our new database approach offers a new perspective on the use of lake-status data for comparative studies between reconstructions based on natural archives and palaeoclimate model simulations. Data-model comparisons have become an integral part of evaluating model performance (Kohfeld and Harrison, 2000; Harrison et al., 2014) but their effectiveness requires standardised quantification of data uncertainty (Harrison et al., 2016). Our new database significantly improves the incorporation of such uncertainties in assessing model simulations using lake-status data and provides significant improvements over previous lake-status syntheses, such as the widely used GLSDB.

The utility of our analytical framework combining MC sampling of uncertainties and RSEOF identification of dominant modes of variability extends beyond lake-status data. Indeed, it can be applied to any palaeoenvironmental data set that comes in the form of time series with chronological, analytical and/or interpretational uncertainties. This opens new possibilities for more inclusive syntheses, employing a standardised methodology across all archive types, data sources and palaeoenvironmental variables.

However, to enable the development of such large-scale studies, adequate reporting of data is crucial. Most of the challenges involved in assembling the ESA lake-status database were related to incomplete reporting of data. While we fully acknowledge that reporting data for purposes beyond the original goal of a paper can be complex and time-consuming, the following points emphasise key elements to consider in order to foster the best use of palaeolimnological data by independent groups:

- Raw data (e.g., lake-level curves, stratigraphic profiles, time series of key indicators) were not always accessible. Where the information was not available from the data producers, we had to digitise the data from figures, which at best led to a loss of data quality, and possibly to a distortion of some signals.
- Stratigraphic data series were often only expressed against the calculated age and not versus original core or section depth. The inclusion of depth information, either in the original publication or in archived data sets, is essential for future updates of the chronologies (e.g., when the radiocarbon calibration curves are updated).

- In some cases chronological data was only presented graphically. Dating information presented in tabular form allow for both a quick assessment of the raw measurements and their errors, and contribute to a better reuse of the data (e.g., Millard, 2014; Courtney Mustaphi et al., 2019).
- The details of the construction of the published chronology were sometimes incomplete. More details on the radiocarbon calibration curve used and/or the method of interpolation between multiple age points would contribute to a more accurate reproduction of the data.
- We encourage a clear definition and discussion of which environmental variables are reconstructed based on the available evidence.

7. Conclusion

Lake-status histories are important sources of information on past climate change, but their synthesis and translation to palaeoclimate change is challenging. We present a new relational lake-status database that combines updated chronologies with the incorporation of uncertainties. We have shown that the combination of MC sampling to integrate the full range of data uncertainty and RSEOF to extract the dominant modes of variability allows the identification of dominant historical trends, and an assessment of their robustness. The application of this approach to ESA identified the dominant millennial-scale palaeoclimate features of the last 20,000 years, consistent with the current understanding of environmental change in the region, including a tendency towards anti-phased lake-status behaviour between the EARS and interior and eastern southern Africa. These results highlight the potential contributions lake-status data can make to large-scale palaeoclimate syntheses, despite their often discontinuous and qualitative nature. The data and methods presented here serve as a proof-of-concept study for future large-scale syntheses of lake status in other regions of the world.

Data availability

The following materials are available on the open data repository Zenodo (<https://doi.org/10.5281/zenodo.4494804>).

- The ESA lake-status database, in both SQLite and spreadsheet format.
- Supplementary description and constraints of all database fields.
- Supplementary documentation for all sites included in the database. This includes relevant background information of the site, presentation of original literature from which lake-status information was sourced, and discussion of consensus lake-status history.
- Catchment shapefiles of all included sites.
- R code for MC-RSEOF analysis on our database format.

Author contributions

GDC: Project administration, Methodology, Software, Formal analysis, Investigation, Data curation, Writing – original draft, Writing – review and editing, Visualisation

MC: Project administration, Methodology, Software, Investigation, Data curation, Writing – review and editing, Visualisation

SLB: Project administration, Methodology, Investigation, Writing – review and editing

CYC: Project administration, Methodology, Writing – review and editing

SPH: Supervision, Conceptualisation, Methodology, Resources, Writing – review and editing

Acknowledgements

GDC acknowledges support from the Research Foundation Flanders (FWO), the Belgian-American Educational Foundation (BAEF) and Belgian Science Policy Brain-be project BR/121/A2 PAMEXEA (Patterns and mechanisms of climate extremes in East Africa). MC has been funded by the Swiss National Science Foundation (SNF) through the HORNET project (200021_169598). SLB acknowledges support from the Trapnell Fund, Environmental Change Institute at the University of Oxford. CYC acknowledges financial support from the National Science Foundation Graduate Research Fellowship and Department of Earth, Atmospheric, and Planetary Sciences at MIT. SPH acknowledges funding from the ERC-funded project GC2.0 (Global Change 2.0: Unlocking the past for a clearer future, grant number 694481) and from JPI-Belmont Forum project entitled Palaeoclimate Constraints on Monsoon Evolution and Dynamics (PaCMEDy). We gratefully acknowledge the support of INQUA funding under the project PALCOM 1609P (Palaeolakes of the Arid Southern Hemisphere). We sincerely thank Bryan N. Shuman, David McGee, Sarah Metcalfe, Maarten Blaauw, Mengna Liao, Bronwyn Dixon, Nicholas Primmer, Luciana Figueiredo Prado and David T. Liefert for constructive discussions on synthesizing lake-status histories.

Figure captions

Fig. 1 Examples of lake-status histories over the last 25,000 years as incorporated in the database. **a)** The palaeoshoreline-based reconstruction of Etosha, Namibia. **b)** The (mostly) sediment-core based record of Lake Naivasha, Kenya. Horizontal error bars indicate the outer limits of 2- σ age uncertainty of each episodes' start and end as determined by reworked chronologies. Relative status class (y-axis) is qualitative, but episodes in orange are linked to quantitative information in the form of absolute water-level elevations from shorelines or other geomorphological evidence. In contrast, grey boxes depict episodes where such a link does not exist. Uncertainty in relative lake status is depicted by lighter-colored areas outlined by a dash-dotted line. **c)** The MC sampling process illustrated on part of Lake Naivasha's lake-status history. Start and end of subsequent lake-status episodes are sampled from split-normal distributions defined by the 95% credible intervals provided in the database, which themselves are derived from Bayesian age-depth modelling or, in the case of palaeoshorelines, from individual geochronological data points. For each episode, status class is sampled from a triangular distribution defined by best-estimate and minimum and maximum possible status classes defined in the database. **Figure size: 1-column width.**

Fig. 2 Schematic overview of the structure of the SQLite database. In addition to the 9 main tables (blue), the database contains additional tables that are either used to define lists of accepted values for specific fields (black) or to link the different data tables together (orange). The black tables help to homogenise the information that is entered in the database (e.g., basin type) and facilitate querying the data. The orange tables account for the multiple relationships that can exist between tables. For example, the LakeRefs table connects the Lakes and Refs tables because there is no one-to-one connection between the two tables: one reference can be associated with several lakes and one lake can be associated with several references. For a detailed overview of these tables, see SI. **Figure size: 1.5-column width.**

Fig. 3 Flowchart of analytical approach employed in this study. **Figure size: 1.5-column width.**

Fig. 4 Location of sites in our ESA database against **a)** elevation and **b)** aridity index, which is defined as the ratio of mean annual precipitation over mean annual evapotranspiration (MAP/MAE; Trabucco en Zomer, 2009). Map a shows whether or not the sites were incorporated in the GLSDB, while map b shows the current hydrology of the sites, with symbol colour differentiating between (seasonally) dry, closed, and open basins. Black lines delineate catchment boundaries. Site numbers refer to Table 1. **Figure size: 2-column width.**

Fig. 5 a) Lake sites in the database plotted according to catchment size, best-estimate age of the earliest status-class information, best-estimate temporal coverage, and number of defined status classes. **b)** Number of lake-status episodes of each site's best-estimate history falling (partly) within 500-year bins over the last 25,000 years. **Figure size: 1-column width.**

Fig. 6 a) Spatial distribution of the amount and type of chronological data points used in reworked chronologies. **b)** Site-specific temporal distribution of chronological data points used in reworked chronologies (coloured circles depict best-estimate age) and lake-status information (covered parts of history are denoted by a full black line). Site numbers refer to Table 1. **c)** Database-wide counts of chronological data points used in reworked chronologies, in 1,000-year bins according to type. **Figure size: 2-column width.**

Fig. 7 Number of sites with lake-status information through time over the last 20,000 years, for the entire database and per subregion discussed in the text, across 10,000 MC-generated historical scenarios. Full lines depict the best-estimate scenario for all sites, dark and light shading depicts the 1- σ and 2- σ MC uncertainty envelope. **Figure size: 1-column width.**

Fig. 8 a) Overview of MC-RSEOF experiments over 20,000-0 cal yr BP on MC-generated and best-estimate lake-status histories. Results are plotted according to the site-selection criteria that were used to retain for RSEOF analysis only those sites that met minima of temporal coverage (varying between experiments between 10 and 100 %) and lake-status episodes (2 or 4). The top row of plots shows the frequency of the number of incorporated sites across each experiment's 10,000 MC iterations. The bin also containing the number of sites in the scenario where all uncertainties are set to their best estimate, is shown by the darker shade of purple. The middle row shows the frequency of resulting non-overlapping EOFs from each experiment's 10,000 RSEOF runs, with the darker shade again depicting the value that was also obtained for the best-estimate run. The bottom row of plots shows the distribution of the variance explained by the first two EOFs across each experiment's 10,000 RSEOF runs. Values for the best-estimate scenario are depicted by a darker-shaded point. **b)** Time series of EOF1 from RSEOF analyses over 20,000-0 cal yr BP on MC-generated ensemble of lake-status histories. Site-selection criteria of minimum temporal coverage and defined status episodes vary between 10-100% and 2-4, respectively. Black line depicts the median across iterations, dark grey and light grey shading depicts the 1- σ and 2- σ MC uncertainty envelope. **Figure size: entire page in landscape format.**

Fig. 9 Results of RSEOF analysis over 20,000-0 cal yr BP. Selection criteria for sites were a minimum of 4 defined status episodes within this time window and a temporal coverage of at least 33%. Best-estimate histories as well as MC-sampled histories (10,000 iterations) are considered. **a)** Biplot of EOF1

and 2, depicting loadings of best-estimate scenario (thick-bordered circles) and means of MC-generated scenarios with 1- σ error bars (thin-bordered circles). Point size is proportional to the record's temporal coverage. Only sites for which the best-estimate history met the selection criteria are depicted. **b)** Lambda scores of EOFs for MC-generated (vertical columns and error bars show MC mean $\pm 1\sigma$) and best-estimate (points) histories. The percentage of MC iterations in which a particular EOF is represented is shown above each column. **c)** EOF1 loadings of best-estimate (upper half-circles) and MC-sampled histories (lower half-circles). Half-circle size is proportional to the record's temporal coverage. Only sites for which the best-estimate history met the selection criteria are depicted. **d)** Time series of EOF1. The thick black line represents EOF1 of the best-estimate scenario for each site, the thin line represents the median EOF1 across MC iterations. Dark grey and light grey shading depicts the 1- σ and 2- σ MC uncertainty envelope. **e)** Same as c, but for EOF2. **f)** Same as d, but for EOF2. **Figure size: entire page in landscape format.**

Tables

Table 1 Summary of sites incorporated in our ESA lake-status database and sources of evidence for constructing their lake-status history. For full references, see Supplementary Material.

References

- Almquist-Jacobson, H., 1995. Lake-level fluctuations at Ljustjärnen, central Sweden and their implications for the Holocene climate of Scandinavia. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 118, 269–290. [https://doi.org/10.1016/0031-0182\(95\)00002-2](https://doi.org/10.1016/0031-0182(95)00002-2)
- Anchukaitis, K.J., Tierney, J.E., 2013. Identifying coherent spatiotemporal modes in time-uncertain proxy paleoclimate records. *Clim. Dyn.* 41, 1291–1306. <https://doi.org/10.1007/s00382-012-1483-0>
- Atsawawaranunt, K., Comas-Bru, L., Amirnezhad Mozhdehi, S., Deininger, M., Harrison, S.P., Baker, A., Boyd, M., Kaushal, N., Ahmad, S.M., Ait Brahimi, Y., Arienzo, M., Bajo, P., Braun, K., Burstyn, Y., Chawchai, S., Duan, W., Hatvani, I.G., Hu, J., Kern, Z., Labuhn, I., Lachniet, M., Lechleiter, F.A., Lorrey, A., Pérez-Mejías, C., Pickering, R., Scroxton, N., 2018. The SISAL database: a global resource to document oxygen and carbon isotope records from speleothems. *Earth Syst. Sci. Data Discuss.* 1–64. <https://doi.org/10.5194/essd-2018-17>
- Bartlein, P.J., Harrison, S.P., Izumi, K., 2017. Underlying causes of Eurasian midcontinental aridity in simulations of mid-Holocene climate. *Geophys. Res. Lett.* 44, 9020–9028. <https://doi.org/10.1002/2017GL074476>
- Bessemers, I., Verschuren, D., Russell, J.M., Hus, J., Mees, F., Cumming, B.F., 2008. Palaeolimnological evidence for widespread late 18th century drought across equatorial East Africa 259, 107–120. <https://doi.org/10.1016/j.palaeo.2007.10.002>
- Blaauw, M., 2010. Methods and code for ‘classical’ age-modelling of radiocarbon sequences. *Quat. Geochronol.* 5, 512–518. <https://doi.org/10.1016/j.quageo.2010.01.002>
- Blaauw, M., Christen, J.A., 2011. Flexible paleoclimate age-depth models using an autoregressive gamma process. *Bayesian Anal.* 6, 457–474. <https://doi.org/10.1214/11-BA618>
- Bowler, J.M., 1986. Spatial variability and hydrologic evolution of Australian lake basins: Analogue for pleistocene hydrologic change and evaporite formation. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 54, 21–41. [https://doi.org/10.1016/0031-0182\(86\)90116-1](https://doi.org/10.1016/0031-0182(86)90116-1)
- Breitenbach, S.F.M., Rehfeld, K., Goswami, B., Baldini, J.U.L., Ridley, H.E., Kennett, D.J., Prufer, K.M., Aquino, V. V., Asmerom, Y., Polyak, V.J., Cheng, H., Kurths, J., Marwan, N., 2012. Constructing Proxy Records from Age models (COPRA). *Clim. Past* 8, 1765–1779. <https://doi.org/10.5194/cp-8-1765-2012>

715 Burrough, S.L., Thomas, D.S.G., 2013. Central southern Africa at the time of the African Humid Period:
 716 A new analysis of Holocene palaeoenvironmental and palaeoclimate data. *Quat. Sci. Rev.* 80, 29–
 717 46. <https://doi.org/10.1016/j.quascirev.2013.08.001>

718 Burrough, S.L., Thomas, D.S.G., Bailey, R.M., 2009. Mega-Lake in the Kalahari: A Late Pleistocene
 719 record of the Palaeolake Makgadikgadi system. *Quat. Sci. Rev.* 28, 1392–1411.
 720 <https://doi.org/10.1016/j.quascirev.2009.02.007>

721 Burrough, S.L., Thomas, D.S.G., 2009. Geomorphological contributions to palaeolimnology on the
 722 African continent. *Geomorphology* 103, 285–298.
 723 <https://doi.org/10.1016/j.geomorph.2008.07.015>

724 Carlson, A.E., 2013. The Younger Dryas climate event, in: *The Encyclopedia of Quaternary Science*, Vol.
 725 3. Elsevier, Amsterdam, pp. 126–134.

726 Chase, B.M., Meadows, M.E., Carr, A.S., Reimer, P.J., 2010. Evidence for progressive Holocene
 727 aridification in southern Africa recorded in Namibian hyrax middens: Implications for African
 728 Monsoon dynamics and the “African Humid Period.” *Quat. Res.* 74, 36–45.
 729 <https://doi.org/10.1016/j.yqres.2010.04.006>

730 Chase, B.M., Niedermeyer, E.M., Boom, A., Carr, A.S., Chevalier, M., He, F., Meadows, M.E., Ogle, N.,
 731 Reimer, P.J., 2019. Orbital controls on Namib Desert hydroclimate over the past 50,000 years.
 732 *Geology* 47, 867–871. <https://doi.org/10.1130/G46334.1>

733 Cheddadi, R., Yu, G., Guiot, J., Harrison, S.P., Prentice, I.C., 1996. The climate of Europe 6000 years
 734 ago. *Clim. Dyn.* 13, 1–9. <https://doi.org/10.1007/s003820050148>

735 Chen, C.Y., Maloof, A.C., 2017. Revisiting the deformed high shoreline of Lake Bonneville. *Quat. Sci.*
 736 *Rev.* 159, 169–189. <https://doi.org/10.1016/j.quascirev.2016.12.019>

737 Chen, C.Y., McGee, D., Woods, A., Pérez, L., Hatfield, R.G., Edwards, R.L., Cheng, H., Valero-Garcés, B.
 738 L., Lehmann, S.B., Stoner, J.S., Schwalb, A., Tal, I., Seltzer, G.O., Tapia, P.M., Abbott, M.A., Rodbell,
 739 D.T., 2020. U-Th dating of lake sediments: Lessons from the 700 ka sediment record of Lake Junín,
 740 Peru. *Quaternary Science Reviews*. 244: 106422. <https://doi.org/10.1016/j.quascirev.2020.106422>

741 Chevalier, M., Chase, B.M., 2015. Southeast African records reveal a coherent shift from high- to low-
 742 latitude forcing mechanisms along the east African margin across last glacial-interglacial transition.
 743 *Quat. Sci. Rev.* 125, 117–130. <https://doi.org/10.1016/j.quascirev.2015.07.009>

744 Chorowicz, J., 2005. The East African rift system. *J. African Earth Sci.* 43, 379–410.
 745 <https://doi.org/10.1016/j.jafrearsci.2005.07.019>

746 Coe, T., Harrison, P., 2002. The water balance of northern Africa during the mid-Holocene : an
 747 evaluation of the 6 ka BP PMIP simulations 155–166. <https://doi.org/10.1007/s00382-001-0219-3>

748 Comas-Bru, L., Harrison, S.P., Werner, M., Rehfeld, K., Scroxton, N., Veiga-Pires, C., 2019. Evaluating
 749 model outputs using integrated global speleothem records of climate change since the last glacial.
 750 *Clim. Past* 15, 1557–1579. <https://doi.org/10.5194/cp-15-1557-2019>

751 Courtney Mustaphi, C.J., Brahney, J., Aquino-López, M.A., Goring, S., Orton, K., Noronha, A.,
 752 Czaplewski, J., Asena, Q., Paton, S., Panga Brushworth, J., 2019. Guidelines for reporting and
 753 archiving 210Pb sediment chronologies to improve fidelity and extend data lifecycle. *Quat.*
 754 *Geochronol.* 52, 77–87. <https://doi.org/10.1016/j.quageo.2019.04.003>

755 De Cort, G., Verschuren, D., Ryken, E., Wolff, C., Renaut, R.W., Creutz, M., Van der Meeren, T., Haug,
 756 G., Olago, D.O., Mees, F., 2018. Multi-basin depositional framework for moisture-balance
 757 reconstruction during the last 1300 years at Lake Bogoria, central Kenya Rift Valley. *Sedimentology*
 758 38, 1667–1696. <https://doi.org/10.1111/sed.12442>

759 Delalande, M., 2008. Hydrologie et géochimie isotopique du Lac Masoko et de lacs volcaniques de la
 760 province active du Rungwe (Sud-Ouest Tanzanie). Université Paris Sud.

761 deMenocal, P., Ortiz, J., Guilderson, T., Adkins, J., Sarnthein, M., Baker, L., Yarusinsky, M., 2000. Abrupt
 762 onset and termination of the African Humid Period: *Quat. Sci. Rev.* 19, 347–361.
 763 [https://doi.org/10.1016/S0277-3791\(99\)00081-5](https://doi.org/10.1016/S0277-3791(99)00081-5)

764 Digerfeldt, G., 1965. Vielången och Farlången. En utvecklingshistorisk insjöundersrkning. *Skånes Natur*
 765 52, 162–183.

766 Eggermont, H., Russell, J.M., Schettler, G., Van Damme, K., Bessems, I., Verschuren, D., 2007. Physical
 767 and chemical limnology of alpine lakes and pools in the Rwenzori Mountains (Uganda–DR Congo).
 768 *Hydrobiologia* 592, 151–173. <https://doi.org/10.1007/s10750-007-0741-3>

769 Eggermont, H., Verschuren, D., 2007. Taxonomy and diversity of Afroalpine Chironomidae (Insecta:
 770 Diptera) on Mount Kenya and the Rwenzori Mountains, East Africa. *J. Biogeogr.* 34, 69–89.
 771 <https://doi.org/10.1111/j.1365-2699.2006.01590.x>

772 Fang, J., 1991. Lake evolution during the past 30,000 years in China, and its implications for
 773 environmental change. *Quat. Res.* 36, 37–60. [https://doi.org/10.1016/0033-5894\(91\)90016-X](https://doi.org/10.1016/0033-5894(91)90016-X)

774 Garcin, Y., Junginger, A., Melnick, D., Olago, D.O., Strecker, M.R., Trauth, M.H., 2009. Late Pleistocene-
 775 Holocene rise and collapse of Lake Suguta, northern Kenya Rift. *Quat. Sci. Rev.* 28, 911–925.
 776 <https://doi.org/10.1016/j.quascirev.2008.12.006>

777 Gasse, F., 2000. Hydrological changes in the African tropics since the Last Glacial Maximum. *Quat. Sci.*
778 *Rev.* 19, 189–211. [https://doi.org/10.1016/S0277-3791\(99\)00061-X](https://doi.org/10.1016/S0277-3791(99)00061-X)

779 Gasse, F., Chalié, F., Vincens, A., Williams, M.A.J., Williamson, D., 2008. Climatic patterns in equatorial
780 and southern Africa from 30,000 to 10,000 years ago reconstructed from terrestrial and near-shore
781 proxy data. *Quat. Sci. Rev.* 27, 2316–2340. <https://doi.org/10.1016/j.quascirev.2008.08.027>

782 Goudie, A., Thomas, D.S.G., 1986. Lunette dunes in southern Africa. *J. Arid Environ.* 10, 1–12.
783 [https://doi.org/10.1016/S0140-1963\(18\)31260-6](https://doi.org/10.1016/S0140-1963(18)31260-6)

784 Haddon, I.G., McCarthy, T.S., 2005. The Mesozoic–Cenozoic interior sag basins of Central Africa: The
785 Late-Cretaceous–Cenozoic Kalahari and Okavango basins. *J. African Earth Sci.* 43, 316–333.
786 <https://doi.org/10.1016/j.jafrearsci.2005.07.008>

787 Harrison, S.P., 1989. Lake levels and climatic change in eastern North America. *Clim. Dyn.* 3, 157–167.
788 <https://doi.org/10.1007/BF01080366>

789 Harrison, S.P., 1993. Late Quaternary lake-level changes and climates of Australia. *Quat. Sci. Rev.* 12,
790 211–231. [https://doi.org/10.1016/0277-3791\(93\)90078-Z](https://doi.org/10.1016/0277-3791(93)90078-Z)

791 Harrison, S.P., Saarse, L., Digerfeldt, G., 1991. Holocene changes in lake levels as climate proxy data in
792 Europe. *Paleoklimaforschung* 6, 159–170.

793 Harrison, S.P., Digerfeldt, G., 1993. European lakes as palaeohydrological and palaeoclimatic
794 indicators. *Quat. Sci. Rev.* 12, 233–248. [https://doi.org/10.1016/0277-3791\(93\)90079-2](https://doi.org/10.1016/0277-3791(93)90079-2)

795 Harrison, S.P., Prentice, I.C., Guiot, J., 1993. Climatic controls on Holocene lake-level changes in
796 Europe. *Clim. Dyn.* 8, 189–200. <https://doi.org/10.1007/BF00207965>

797 Harrison, S.P., Yu, G., Tarasov, P.E., 1996. Late Quaternary Lake-Level Record from Northern Eurasia.
798 *Quat. Res.* 45, 138–159. <https://doi.org/10.1006/qres.1996.0016>

799 Harrison, S.P., Yu, G., Vassiljev, J., 2002. Climate changes during the Holocene recorded by lakes from
800 Europe, in: *Climate Development and History of the North Atlantic Realm*. Springer, Berlin,
801 Heidelberg, pp. 191–204.

802 Harrison, S.P., Bartlein, P.J., Brewer, S., Prentice, I.C., Boyd, M., Hessler, I., Holmgren, K., Izumi, K.,
803 Willis, K., 2014. Climate model benchmarking with glacial and mid-Holocene climates. *Clim. Dyn.*
804 43, 671–688. <https://doi.org/10.1007/s00382-013-1922-6>

805 Harrison, S.P., Bartlein, P.J., Prentice, I.C., 2016. What have we learnt from palaeoclimate simulations?
806 *J. Quat. Sci.* 31, 363–385. <https://doi.org/10.1002/jqs.2842>

807 Hatfield, R., Stoner, J., Solada, K.E., Morey, A.E., Woods, A., Chen, C.Y., McGee, D., Abbott, M.B.,
808 Rodbell, D.T., 2020. Paleomagnetic Constraint of the Brunhes Age Sedimentary Record from Lake
809 Junín, Peru. *Frontiers in Earth Science* 8. <https://doi.org/10.3389/feart.2020.00147>.

810 Heegaard, E., Birks, H.J.B., Telford, R.J., 2005. Relationships between calibrated ages and depth in
811 stratigraphical sequences: an estimation procedure by mixed-effect regression. *The Holocene* 15,
812 612–618. <https://doi.org/10.1191/0959683605hl836rr>

813 Hill, B.J., 1975. The origin of southern African coastal lakes. *Trans. R. Soc. South Africa* 41, 225–240.
814 <https://doi.org/10.1080/00359197509519440>

815 Hogg, A.G., Hua, Q., Blackwell, P.G., Niu, M., Buck, C.E., Guilderson, T.P., Heaton, T.J., Palmer, J.G.,
816 Reimer, P.J., Reimer, R.W., Turney, C.S.M., Zimmerman, S.R.H., 2013. SHCal13 Southern
817 Hemisphere Calibration, 0–50,000 Years cal BP. *Radiocarbon* 55, 1889–1903.
818 https://doi.org/10.2458/azu_js_rc.55.16783

819 Holmes, J., Hoelzmann, P., 2017. The Late Pleistocene-Holocene African Humid Period as Evident in
820 Lakes. Oxford University Press. <https://doi.org/10.1093/acrefore/9780190228620.013.531>

821 Jolly, D., Prentice, I.C., Bonnefille, R., Ballouche, A., Bengo, M., Brenac, P., Buchet, G., Burney, D., Cazet,
822 J., Cheddadi, R., Edorh, T., Elenga, H., Elmoutaki, S., Guiot, J., Laarif, F., Lamb, H., Lezine, A., Maley,
823 J., Mbenza, M., Peyron, O., Reille, M., Reynaud-Farrera, I., Riollet, G., Ritchie, J.C., Roche, E., Scott,
824 L., Ssemmanda, I., Straka, H., Umer, M., Van Campo, E., Vilimumbalo, S., Vincens, A., Waller, M.,
825 1998. Biome reconstruction from pollen and plant macrofossil data for Africa and the Arabian
826 peninsula at 0 and 6000 years. *J. Biogeogr.* 25, 1007–1027. <https://doi.org/10.1046/j.1365-2699.1998.00238.x>

827

828 Juggins, S., 2013. Quantitative reconstructions in palaeolimnology: new paradigm or sick science?
829 *Quat. Sci. Rev.* 64, 20–32. <https://doi.org/10.1016/j.quascirev.2012.12.014>

830 Kalff, J., 2001. Chapter 5: Hydrology and climate, in: *Limnology*. Prentice Hall, Upper Saddle River, New
831 Jersey, pp. 53–71.

832 Kohfeld, K.E., Harrison, S.P., 2000. How well can we simulate past climates? Evaluating the models
833 using global palaeoenvironmental datasets 19.

834 Kutterolf, S., Schindlbeck, J.C., Anselmetti, F.S., Ariztegui, D., Brenner, M., Curtis, J., Schmid, D., Hodell,
835 D., Mueller, A., Pérez, W., Schwalb, A., Frische, M., Wang, K.L., 2016. A 400-ka tephrochronological
836 framework for Central America from Lake Petén Itzá (Guatemala) sediments. *Quaternary Science*
837 *Reviews* 150, 200–220. <https://doi.org/10.1016/j.quascirev.2016.08.023>.

838 Kutzbach, J.E., Otto-Bliesner, B.L., 1982. The Sensitivity of the African-Asian Monsoonal Climate to
839 Orbital Parameter Changes for 9000 Years B.P. in a Low-Resolution General Circulation Model. *J.*
840 *Atmos. Sci.* 39, 1177–1188. [https://doi.org/10.1175/1520-0469\(1982\)039<1177:TSOTAA>2.0.CO;2](https://doi.org/10.1175/1520-0469(1982)039<1177:TSOTAA>2.0.CO;2)

841 Legendre, P., Legendre, L.F.J., 2012. *Numerical Ecology Volume 24 Developments in Environmental*
842 *Modelling*. Elsevier.

843 Lehner, B., Verdin, K., Jarvis, A., 2008. New Global Hydrography Derived From Spaceborne Elevation
844 Data. *Eos, Trans. Am. Geophys. Union* 89, 93. <https://doi.org/10.1029/2008EO100001>

845 Liu, G., Schwartz, F.W., 2014. On modeling the paleohydrologic response of closed-basin lakes to
846 fluctuations in climate: Methods, applications, and implications. *Water Resour. Res.* 50, 2975–
847 2992. <https://doi.org/10.1002/2013WR014107>

848 Liu, X., Rendle-Bühning, R., Kuhlmann, H., Li, A., 2017. Two phases of the Holocene East African Humid
849 Period: Inferred from a high-resolution geochemical record off Tanzania. *Earth Planet. Sci. Lett.*
850 460, 123–134. <https://doi.org/10.1016/j.epsl.2016.12.016>

851 Lohmann, G., Pfeiffer, M., Laepple, T., Leduc, G., Kim, J.-H., 2013. A model–data comparison of the
852 Holocene global sea surface temperature evolution. *Clim. Past* 9, 1807–1839.
853 <https://doi.org/10.5194/cp-9-1807-2013>

854 Mahaney, W.C., 2004. Quaternary Glacial Chronology of Mount Kenya Massif, in: Ehlers, J., Gibbard,
855 P. (Eds.), *Quaternary Glaciations - Extent and Chronology, Part III*. Elsevier B.V., pp. 227–231.

856 Mason, I.M., Guzkowska, M.A.J., Rapley, C.G., Street-Perrott, F.A., 1994. The response of lake levels
857 and areas to climatic change. *Clim. Change* 27, 161–197. <https://doi.org/10.1007/BF01093590>

858 McKay, N.P., Kaufman, D.S., 2014. An extended Arctic proxy temperature database for the past 2,000
859 years. *Sci. Data* 1, 140026. <https://doi.org/10.1038/sdata.2014.26>

860 Melack, J.M., 1978. Morphometric, physical and chemical features of the volcanic crater lakes of
861 western Uganda. *Arch. fur Hydrobiol.* 84, 430–453.

862 Meybeck, M., 1995. Global distribution of lakes, in: Lerman, A., Imboden, D.M., Gat, J.R. (Eds.), *Physics*
863 *and Chemistry of Lakes*. Springer-Verlag Berlin Heidelberg, pp. 1–36.

864 Millard, A.R., 2014. Conventions for Reporting Radiocarbon Determinations. *Radiocarbon* 56, 555–
865 559. <https://doi.org/10.2458/56.17455>

866 Nicholson, S.E., 2018. The ITCZ and the seasonal cycle over equatorial Africa. *Bull. Am. Meteorol. Soc.*
867 99, 337–348. <https://doi.org/10.1175/BAMS-D-16-0287.1>

868 Nicholson, S.E., 2017. Climate and climatic variability of rainfall over eastern Africa. *Rev. Geophys.* 55,
869 590–635. <https://doi.org/10.1002/2016RG000544>

870 North, G.R., Bell, T.L., Cahalan, R.F., Moeng, F.J., 1982. Sampling Errors in the Estimation of Empirical
871 Orthogonal Functions. *Mon. Weather Rev.* 110, 699–706. [https://doi.org/10.1175/1520-0493\(1982\)110<0699:SEITEO>2.0.CO;2](https://doi.org/10.1175/1520-0493(1982)110<0699:SEITEO>2.0.CO;2)

873 Olaka, L.A., Odada, E.O., Trauth, M.H., Olago, D.O., 2010. The sensitivity of East African rift lakes to
874 climate fluctuations. *J. Paleolimnol.* 44, 629–644. <https://doi.org/10.1007/s10933-010-9442-4>

875 Otto-Bliesner, B.L., Russell, J.M., Clark, P.U., Liu, Z., Overpeck, J.T., Konecky, B., DeMenocal, P.,
876 Nicholson, S.E., He, F., Lu, Z., 2014. Coherent changes of southeastern equatorial and northern
877 African rainfall during the last deglaciation. *Science* (80-.). 346, 1223–1227.
878 <https://doi.org/10.1126/science.1259531>

879 Parnell, A.C., Buck, C.E., Doan, T.K., 2011. A review of statistical chronology models for high-resolution,
880 proxy-based Holocene palaeoenvironmental reconstruction. *Quat. Sci. Rev.* 30, 2948–2960.
881 <https://doi.org/10.1016/j.quascirev.2011.07.024>

882 Qin, B., Harrison, S.P., Kutzbach, J.E., 1998. Evaluation of modelled regional water balance using lake
883 status data. *Quat. Sci. Rev.* 17, 535–548. [https://doi.org/10.1016/S0277-3791\(98\)00011-0](https://doi.org/10.1016/S0277-3791(98)00011-0)

884 R Core Team 2020. R: A language and environment for statistical computing. R Foundation for
885 Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>

886 Rasmussen, S.O., Andersen, K.K., Svensson, A.M., Steffensen, J.P., Vinther, B.M., Clausen, H.B.,
887 Siggaard-Andersen, M.-L., Johnsen, S.J., Larsen, L.B., Dahl-Jensen, D., Bigler, M., Röthlisberger, R.,
888 Fischer, H., Goto-Azuma, K., Hansson, M.E., Ruth, U., 2006. A new Greenland ice core chronology
889 for the last glacial termination. *J. Geophys. Res.* 111, D06102.
890 <https://doi.org/10.1029/2005JD006079>

891 Reid, R.E.B., Jones, M., Brandt, S., Bunn, H., Marshall, F., 2019. Oxygen isotope analyses of ungulate
892 tooth enamel confirm low seasonality of rainfall contributed to the African Humid Period in
893 Somalia. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 534, 109272.
894 <https://doi.org/10.1016/j.palaeo.2019.109272>

895 Reimer, P.J., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G., Ramsey, C.B., Buck, C.E., Cheng, H.,
896 Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P., Haflidason, H., Hajdas, I., Hatté, C.,
897 Heaton, T.J., Hoffmann, D.L., Hogg, A.G., Hughen, K.A., Kaiser, K.F., Kromer, B., Manning, S.W., Niu,
898 M., Reimer, R.W., Richards, D.A., Scott, E.M., Southon, J.R., Staff, R.A., Turney, C.S.M., van der

899 Plicht, J., 2013. IntCal13 and Marine13 Radiocarbon Age Calibration Curves 0–50,000 Years cal BP.
900 Radiocarbon 55, 1869–1887. https://doi.org/10.2458/azu_js_rc.55.16947

901 Roberts, H.M., Bryant, C.L., Huws, D.G., Lamb, H.F., 2018. Generating long chronologies for lacustrine
902 sediments using luminescence dating: a 250,000 year record from Lake Tana, Ethiopia. *Quat. Sci.*
903 *Rev.* 202, 66–77. <https://doi.org/10.1016/j.quascirev.2018.10.037>.

904 Schagerl, M., Renaut, R.W., 2016. Dipping into the soda lakes of East Africa, in: Schagerl, M. (Ed.), *Soda*
905 *Lakes of East Africa*. Springer International Publishing Switzerland, pp. 3–24.

906 Schneider, R., Tobolski, K., 1985. Lago di Ganna - Late glacial and Holocene environments of a lake in
907 the southern Alps. *Diss. Bot.* 87, 229–271.

908 Shanahan, T.M., McKay, N.P., Hughen, K.A., Overpeck, J.T., Otto-Bliesner, B., Heil, C.W., King, J., Scholz,
909 C., Peck, J., 2015. The time-transgressive termination of the African Humid Period. *Nat. Geosci.* 8,
910 144–144. <https://doi.org/10.1038/ngeo2329>

911 Shuman, B., Henderson, A.K., Colman, S.M., Stone, J.R., Fritz, S.C., Stevens, L.R., Power, M.J., Whitlock,
912 C. 2009. Holocene lake-level trends in the Rocky Mountains, U.S.A. *Quaternary Science Reviews.*
913 28(19–20), 1861–1879. <https://doi.org/10.1016/j.quascirev.2009.03.003>

914 Singarayer, J.S., Burrough, S.L., 2015. Interhemispheric dynamics of the African rainbelt during the late
915 Quaternary. *Quat. Sci. Rev.* 124, 48–67. <https://doi.org/10.1016/j.quascirev.2015.06.021>

916 Singarayer, J.S., Williams, C., Koriche, S., Dadson, S., Burrough, S.L., 2019. Modelling late Quaternary
917 surface hydrology changes and climate feedbacks over African drylands, in: 20th Congress of the
918 International Union for Quaternary Research (INQUA). Dublin, Ireland.

919 Specht, T., Rosendahl, B., 1989. Architecture of the Lake Malawi Rift, East Africa. *J. African Earth Sci.*
920 8, 355–382.

921 Stager, J.C., Ryves, D.B., Chase, B.M., Pausata, F.S.R., 2011. Catastrophic Drought in the Afro-Asian
922 Monsoon Region During Heinrich Event 1. *Science* (80-.). 331, 1299–1302.
923 <https://doi.org/10.1126/science.1198322>

924 Stager, J.C., Mayewski, P.A., Meeker, L.D., 2002. Cooling cycles, Heinrich event 1, and the desiccation
925 of Lake Victoria. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 183, 169–178.
926 [https://doi.org/10.1016/S0031-0182\(01\)00468-0](https://doi.org/10.1016/S0031-0182(01)00468-0)

927 Steiger, N.J., Smerdon, J.E., Cook, E.R., Cook, B.I., 2018. A reconstruction of global hydroclimate and
 928 dynamical variables over the Common Era. *Sci. Data* 5:180086.
 929 <https://doi.org/10.1038/sdata.2018.86>

930 Stone, A.E.C., 2014. Last Glacial Maximum conditions in southern Africa. *Prog. Phys. Geogr. Earth*
 931 *Environ.* 38, 519–542. <https://doi.org/10.1177/0309133314528943>

932 Street, F.A., Grove, A.T., 1979. Global Maps of Lake-Level Fluctuations since 30,000 yr B.P. *Quat. Res.*
 933 12, 83–118. [https://doi.org/10.1016/0033-5894\(79\)90092-9](https://doi.org/10.1016/0033-5894(79)90092-9)

934 Street, F.A., 1980. The relative importance of climate and local hydrogeological factors in influencing
 935 lake-level fluctuations. *Palaeoecol. Africa* 12, 137–158.

936 Tarasov, P.E., Harrison, S.P., Saarse, L., Pushenko, M.Y., Andreev, A.A., Aleshinskaya, Z. V., Davydova,
 937 N.N., Dorofeyuk, N.I., Efremov, Y. V., Khomutova, V.I., Sevastyanov, D. V., Tamosaitis, J.,
 938 Uspenskaya, O.N., Yakushko, O.F., Tarasova, I. V., 1994. Lake status records from the Former Soviet
 939 Union and Mongolia, data base documentation. NOAA Paleoclimatology Publications Series Report
 940 2. Boulder.

941 Taylor, M., 2017. sinkr: Collection of functions with emphasis on multivariate data analysis. R package
 942 version 0.6.

943 Taylor, M.H., Losch, M., Wenzel, M., Schröter, J., 2013. On the sensitivity of field reconstruction and
 944 prediction using empirical orthogonal functions derived from Gappy data. *J. Clim.* 26, 9194–9205.
 945 <https://doi.org/10.1175/JCLI-D-13-00089.1>

946 Telfer, M.W., Thomas, D.S.G., 2006. Complex Holocene lunette dune development, South Africa:
 947 Implications for paleoclimate and models of pan development in arid regions. *Geology* 34, 853.
 948 <https://doi.org/10.1130/G22791.1>

949 Telford, R.J., Heegaard, E., Birks, H.J.B., 2004. Alle age-depth models are wrong: but how badly? *Quat.*
 950 *Sci. Rev.* 23, 1-5. <https://doi.org/10.1016/j.quascirev.2003.11.003>

951 Thomas, D., 2016. Desert dune environments, in: Knight, J., Grab, S.W. (Eds.), *Environmental Change*
 952 *in Southern Africa: Physical and Human Dimensions*. Cambridge University Press, Cambridge.

953 Tiercelin, J.-J., Gibert, E., Umer, M., Bonnefille, R., Disnar, J.-R., Lézine, A.-M., Hureau-Mazaudier, D.,
 954 Travi, Y., Keravis, D., Lamb, H.F., 2008. High-resolution sedimentary record of the last deglaciation
 955 from a high-altitude lake in Ethiopia. *Quat. Sci. Rev.* 27, 449–467.
 956 <https://doi.org/10.1016/j.quascirev.2007.11.002>

957 Tiercelin, J.-J., Lezzar, K.E., 2002. A 300 million years history of rift lakes in Central and East Africa: an
 958 updated broad review, in: Odada, E., Olago, D.O. (Eds.), *The East African Great Lakes: Limnology,*
 959 *Palaeolimnology and Biodiversity.* Kluwer Academic Publishers, pp. 3–60.

960 Tierney, J.E., Lewis, S.C., Cook, B.I., LeGrande, A.N., Schmidt, G.A., 2011. Model, proxy and isotopic
 961 perspectives on the East African Humid Period. *Earth Planet. Sci. Lett.* 307, 103–112.
 962 <https://doi.org/10.1016/j.epsl.2011.04.038>

963 Tierney, J.E., Smerdon, J.E., Anchukaitis, K.J., Seager, R., 2013. Multidecadal variability in East African
 964 hydroclimate controlled by the Indian Ocean. *Nature* 493, 389–392.
 965 <https://doi.org/10.1038/nature11785>

966 Trabucco, A., Zomer, R., 2009. Global aridity index (Global-Aridity) and global potential
 967 evapotranspiration (Global-PET) geospatial database [WWW Document]. CGIAR Consort. Spat. Inf.
 968 URL <http://www.csi.cgiar.org>

969 Tyson, P.D., Preston-Whyte, R., 2000. *The weather and climate of southern Africa.* Oxford University
 970 Press, Cape Town.

971 Vassiljev, J., Harrison, S.P., Guiot, J., 1998. Simulating the Holocene Lake-Level Record of Lake Bysjön,
 972 Southern Sweden. *Quat. Res.* 49, 62–71. <https://doi.org/10.1006/qres.1997.1942>

973 Verschuren, D., 2003. Lake-based climate reconstruction in Africa: progress and challenges.
 974 *Hydrobiologia* 500, 315–330. <https://doi.org/10.1023/A:1024686229778>

975 Wang, L., Brook, G.A., Burney, D.A., Voarintsoa, N.R.G., Liang, F., Cheng, H., Edwards, R.L., 2019. The
 976 African Humid Period, rapid climate change events, the timing of human colonization, and
 977 megafaunal extinctions in Madagascar during the Holocene: Evidence from a 2m Anjohibe Cave
 978 stalagmite. *Quat. Sci. Rev.* 210, 136–153. <https://doi.org/10.1016/j.quascirev.2019.02.004>


979 Weldeab, S., Lea, D.W., Schneider, R.R., Andersen, N., 2007. 155,000 years of West African Monsoon
 980 and ocean thermal evolution. *Science* 316, 1303–1307. <https://doi.org/10.1126/science.1140461>

981 Winkler, M.G., Swain, A.M., Kutzbach, J.E., 1986. Middle Holocene Dry Period in the Northern
 982 Midwestern United States: Lake Levels and Pollen Stratigraphy. *Quat. Res.* 25, 235–250.
 983 [https://doi.org/10.1016/0033-5894\(86\)90060-8](https://doi.org/10.1016/0033-5894(86)90060-8)


984 Whitfield, A.K., Weerts, S.P., Weyl, O.L.F., 2017. A review of the influence of biogeography, riverine
 985 linkages, and marine connectivity on fish assemblages in evolving lagoons and lakes of coastal
 986 southern Africa. *Ecol. Evol.* 7, 7382–7398. <https://doi.org/10.1002/ece3.3266>

- 987 Yi, G., Zhang, T., 2015. Delayed Response of Lake Area Change to Climate Change in Siling Co Lake,
988 Tibetan Plateau, from 2003 to 2013. *Int. J. Environ. Res. Public Health* 12, 13886–13900.
989 <https://doi.org/10.3390/ijerph121113886>
- 990 Yu, G., Harrison, S.P., 1996. An evaluation of the simulated water balance of Eurasia and northern
991 Africa at 6000 y BP using lake status data. *Clim. Dyn.* 12, 723–735.
992 <https://doi.org/10.1007/s003820050139>


A Etosha, Namibia




B Lake Naivasha, Kenya




C



 Primary key
 Foreign key


contributors	
	lakeID
	contributor
	ORCID
	date

alternativeNames	
	lakeID
	alternativeName


basinTypes	
	basinType





dataSources	
	dataSource



dateTypes	
	dateType




materialDated	
	materialDated

lakes	
	lakeID
	lakeName
	latitude
	longitude
	elevation
	basinType
	area
	volume
	catchment








refs	
	referenceID
	abbrev
	reference
	DOI



dating*	
	lakeID
	dateID
	dateType
	materialDated
	locationType
	featureName
	datedReference
	depth
	elevation
	thickness
	labnumber
	rc_age
	...
	...

codingBases	
	lakeID
	codingCategory
	evidence


codingSources	
	lakeID
	dataSource
	referenceID
	used




lakeRefs	
	lakeID
	referenceID

coding	
	lakeID
	levelID
	codingCategory
	minStatus
	maxStatus
	basinType
	shorelineID
	startDate
	oldestStartDate
	youngestStartDate
	endDate
	oldestEndDate
	youngestEndDate

lakeSizes	
	lakeID
	shorelineID
	shorelineFeature
	stillWaterLevel
	stillWaterLevelLower
	stillWaterLevelHigher
	area
	areaUncertainty
	volume
	volumeUncertainty
	catchmentArea
	catchmentAreaUncer.

codingLakeSizes	
	lakeID
	levelID
	shorelineID

lakeSizesRefs	
	lakeID
	shorelineID
	referenceID

datingRefs	
	lakeID
	dateID
	referenceID

I

Probabilistic lake-status histories with defined uncertainties

Lake 1

Lake 2

...

Lake x

II

MC scenario 1

Lake 1

Lake 2

...

Lake x

MC scenario 2

Lake 1

Lake 2

...

Lake x

MC scenario 3

Lake 1

Lake 2

...

Lake x

⋮

MC scenario 10,000

Lake 1

Lake 2

...

Lake x

In addition to the MC-generated scenarios, a scenario is created where all uncertainties are set to their best estimate.

best-estimate scenario

Lake 1

Lake 2

...

Lake x

MC sampling
generates scenarios of possible
(i.e., allowed by the uncertainties)
time series of lake status.

III

EOF PC 1

EOF PC 2

...

EOF PC x-1

EOF PC 1

EOF PC 2

...

EOF PC x-1

EOF PC 1

EOF PC 2

...

EOF PC x-1

⋮

EOF PC 1

EOF PC 2

...

EOF PC x-1



EOF PC 1
ensemble

EOF PC 2
ensemble

...

EOF PC x-1
ensemble


RSEOF
identifies dominant
patterns of variability
in each scenario.

EOF PC 1

EOF PC 2

...

EOF PC x-1




In GLSDB

- yes
- no




Lake type



- (seasonally) dry
- closed
- open

Elevation (m asl)




MAP/MAE






a


Dates used in new chronologies





b



c







ID	Site name	Coordinates		Current hydrology	Lake-status proxies used															Quantitative info available	References
		Lon	Lat		Paleo-shorelines	Non-shoreline geomorph.	Stratigraphy	Sedimentology	Mineralogy	Aquatic plants	Diatoms	Ostracodes	Algae	Molluscs	Chironomids	Isotopes	Archaeological data	Historical data	Instrumental/gauge data		
1	Lake Chilwa	35.5	-15.5	closed	x												x		x	Owen and Crossley (1990); Thomas et al. (2009)	
2	Lake Tritrivakely	46.92	-19.78	closed					x	x	x									Gasse et al. (1994); Williamson et al. (1998); Gasse and van Campo (1998); Gasse and van Campo (2001)	
3	Lake Lungué	33.6358	-24.7619	closed							x									Sitoe et al. (2017)	
4	Lake Cheshi	29.75	-9.08333	closed							x									Stager (1988)	
5	Ishiba Ngandu	31.741	-11.235	open			x				x									Livingstone (1971); Haberyan (2018)	
6	Lake Nhauhache	35.2947	-21.9807	closed							x									Holmgren et al. (2012)	
7	Lake Mapimbi	31.28	-22.4025	closed							x									Gillson and Ekblom (2009); Ekblom et al. (2012)	
8	Lake Sibaya	32.6121	-27.3441	closed							x									Neumann et al. (2008); Stager et al. (2013)	
9	Princess Vlei	18.483	-34.047	open, significant groundwater influence							x									Kirsten and Meadows (2016)	
10	Lake Chilau	34.9486	-23.9644	open							x									Norstrom et al. (2018)	
11	Lake Nhaucati	35.3120	-22.0377	closed							x									Ekblom and Stabell (2008); Norstrom et al. (2018b)	
12	Verlorenvlei	18.433	-32.35	open							x									Kirsten et al. (2020)	
13	Lake Naivasha	36.3548	-0.7636	closed, significant groundwater influence	x	x	x	x	x		x	x			x				x	Richardson and Richardson (1972); Washbourne-Kamau (1975); Richardson and Dussinger (1986); Verschuren et al. (2000); Verschuren (2001); Bergner et al. (2003); Bergner and Trauth (2004); Van der Meeren et al. (2019)	
14	Lake Ashenge	39.5013	12.5795	closed		x	x	x	x		x					x				Marshall et al. (2009)	
15	Lake Nabugabo	31.8995	-0.3639	open, significant groundwater influence	x			x	x											Stuiver et al. (1960); Stager et al. (2005)	
16	Lake Chibwera	30.1419	-0.1546	closed			x	x	x											Bessemis et al. (2008)	
17	Lake Emakat	35.8409	-2.9125	closed			x	x		x	x	x			x					Muzuka et al. (2004); Ryner et al. (2007); Ryner et al. (2008)	
18	Lake Victoria	33	-1	open, artificially controlled	x		x				x								x	Stuiver et al. (1960); Kendall (1969); Stager (1984); Stager et al. (1986); Sene and Plinston (1994); Johnson et al. (1996); Stager et al. (1997); Johnson et al. (1998); Johnson et al. (2000); Beuning et al. (2002); Stager et al. (2002); Stager et al. (2003); Stager et al. (2005); Stager et al. (2005b);	
19	Suguta	36.5	2.05	closed, seasonally filled	x	x	x	x						x					x	Truckle (1976); Garcin et al. (2009); Junginger et al. (2014)	

20	Lake Challa	37.7	-3.317	closed		x	x	x												Payne (1970); Verschuren et al. (2009); Moernaut et al. (2010); Wolff et al. (2011); Blaauw et al. (2011)
21	Nakuru-Elmenteita	36.17	-0.4	closed	x	x		x			x	x						x	x	Washbourn-Kamau (1971); Butzer et al. (1972); Vareschi (1982); Cohen et al. (1983); Richardson and Dussinger (1986); Dühnforth et al. (2006); De Cort et al. (2013)
22	Kitagata	29.9755	-0.0637	closed			x	x	x											Russell et al. (2007)
23	Kibengo	30.1783	-0.0817	open			x	x	x											Russell et al. (2007)
24	Lake Hayq	39.7167	11.35	closed	x	x	x											x		Lamb et al. (2007); Ghinassi et al. (2012); Ghinassi et al. (2015)
25	Lake Kasenda	30.29	0.4322	closed			x	x	x									x		Ssemmanda et al. (2005); Bessems (2007); Ryves et al. (2011); Mills and Ryves (2012)
26	Lake Nyamogusingiri	30.013	-0.2846	closed			x				x									Mills et al. (2014)
27	Lake Kyasanduka	30.0502	-0.2898	closed			x				x									Mills et al. (2014)
28	Lake Tana	37.25	12	open			x	x			x									Lamb et al. (2014); Marshall et al. (2011); Costa et al. (2014)
29	Lake Abhe	41.8	11.2	closed	x	x	x	x	x		x									Gasse (1977); Gasse and Street (1978)
30	Lake Abiyata (Ziway-Shala)	38.6	7.6	closed	x	x	x	x	x		x								x	Gasse and Street (1978); Gillespie et al. (1983); Chalié and Gasse (2002); Legesse et al. (2002)
31	Lake Duluti	36.7833	-3.3833	closed							x									Öberg et al. (2012); Öberg et al. (2013)
32	Lake Tilo	38.0958	7.0625	closed			x	x	x		x							x		Telford and Lamb (1999); Lamb et al. (2000); Lamb et al. (2004); Lamb et al. (2005)
33	Lake Sonachi	36.262	-0.782	closed			x	x	x											Damnati and Taieb (1996); Verschuren (1999); Verschuren et al. (1999)
34	Lake Albert	31	1.5	open			x	x	x	x	x		x							Hecky and Degens (1973); Harvey (1976); Stoffers and Singer (1979); Beuning et al. (1997); Lehman (1998)
35	Lake Turkana	36	3.5	closed	x	x	x	x	x									x	x	Butzer (1971); Butzer et al. (1972); Robbins (1972); Phillipson (1977); Owen et al. (1982); Halfman and Johnson (1989); Johnson et al. (1991); Halfman et al. (1992); Halfman et al. (1994); Ricketts and Johnson (1996); Mohammed et al. (1996); Nicholson (1998); Brown and Fuller (2008); Avery (2010); Velpuri et al. (2012); Garcin et al. (2012); Forman et al. (2014); Morrissey and Scholz (2014); Bloszies et al. (2015); Bloszies and Forman (2015)
36	Magadi-Natron	36.04	-2.12	closed, seasonally filled				x	x		x							x		Butzer et al. (1972); Hillaire-Marcel et al. (1986); Hillaire-Marcel and Casanova (1987); Barker et al. (1990); Taieb et al. (1991); Roberts et al. (1993); Williamson et al. (1993); Damnati and Taieb (1995); Hughes (2008)
37	Lake Edward	29.5833	-0.4167	open			x	x	x											Bishop (1969); Brooks and Smith (1987); Musisi (1991); de Heinzelin and Verniers (1996); Laerdal et al. (2002); Russell et al. (2003); Russell and Johnson (2005); Russell and Johnson (2007)
38	Lake Hora	41.95	9.43	closed	x														x	Williams et al. (1977)
39	Lake Besaka	39.87	8.86	closed, significant groundwater influence		x	x													Williams et al. (1977); Williams et al. (1981)
40	Loboi Swamp	36.05	0.3667	open, bog			x	x	x		x							x		Ashley et al. (2004); Driese et al. (2004)
41	Lake Rukwa	32.717	-8.417	closed	x		x	x	x		x							x	x	Clark et al. (1970); Haberyan (1987); Talbot and Livingstone (1989); Delvaux et al. (1998); Nicholson

																				(1999); Thevenon et al. (2002); Barker et al. (2002); Vincens et al. (2005)
42	Sacred Lake	37.5333	0.05	closed, significant groundwater influence			x													Coetzee (1964); Coetzee (1967); Huang et al. (1999); Olago et al. (1999); Olago et al. (2000); Olago et al. (2001); Loomis et al. (2012); Konecky et al. (2014)
43	Lake Baringo	36.0833	0.5333	closed	x		x										x	x	x	Williams and Johnson (1976); Tiercelin et al. (1987); Renaut et al. (2000); Bessems et al. (2008); Kiage and Liu (2009); Kiage and Liu (2009b); Obando et al. (2016)
44	Lake George	30.3	0	open			x													Greenwood (1976); Viner (1977); Laerdal et al. (2002)
45	Lake Tanganyika	29.5	-6	open	x	x	x	x			x	x					x	x		Stoffers and Hecky (1978); Livingstone (1965); Haberyan and Hecky (1987); Gasse et al. (1989); Casanova and Hillaire-Marcel (1992); Cohen et al. (1997); Nicholson (1999); Alin and Cohen (2003); Scholz et al. (2003); Cohen et al. (2005); Felton et al. (2007); Stager et al. (2009); Tierney et al. (2010)
46	Lake Manyara	35.82	-3.62	closed		x											x			Greenway and Vesey-Fitzgerald (1969); Holdship (1976); Casanova and Hillaire-Marcel (1992); Casanova and Hillaire-Marcel (1992b); Barker (1992); Barker and Gasse (2003)
47	Lake Masoko	33.755	-9.3333	closed					x											Williamson et al. (1999); Barker et al. (2000); Gibert et al. (2002); Barker et al. (2003); Barker and Gasse (2003); Delalande et al. (2005); Garcin et al. (2006a); Garcin et al. (2006b); Garcin et al. (2007a); Garcin et al. (2007b)
48	Lake Bogoria	36.1	0.25	closed			x	x	x								x	x		Young and Renaut (1979); Tiercelin et al. (1981); Carbonel et al. (1983); Vincens et al. (1986); Vincens et al. (1986b); Tiercelin et al. (1987); Hickley et al. (2003); Onyando et al. (2005); McCall (2010); De Cort et al. (2013); De Cort et al. (2018)
49	Chew Bahir	37	4.75	closed, seasonally filled		x											x			Grove et al. (1975); Foerster et al. (2012)
50	Lake Afrera	40.9	13.27	closed, significant groundwater influence		x	x				x									Bannert et al. (1970); Gasse et al. (1974); Delibrias et al. (1974); Bonatti et al. (2017)
51	Dobi	42	11.5	closed, significant groundwater influence		x	x													Gasse et al. (1974)
52	Lake Malawi	34.5	-12	open		x	x				x						x	x	x	Beadle (1981); Scholz and Rosendahl (1988); Specht and Rosendahl (1989); Owen et al. (1990); Finney and Johnson (1991); Scholz and Finney (1994); Ricketts and Johnson (1996); Wüest et al. (1996); Johnson et al. (2001); Barry et al. (2002); Gasse et al. (2002); Filippi and Talbot (2005); Johnson and McCave (2008); Scholz et al. (2011); Van Bockxlaer et al. (2012)
53	Lake Nkuruba	30.3031	0.5169	closed			x	x												Chapman et al. (1998); Saulnier-Talbot et al. (2018)
54	Urwi Pan	20.3658	-23.3419	closed, dry	x														x	Lancaster (1979)
55	Tsodilo Hills	21.7349	-18.7846	closed, dry	x			x			x			x					x	Brook et al. (1992); Thomas et al. (2003)
56	#Gi	21.0080	-19.6242	closed, dry			x	x												Helgren and Brooks (1983); Brooks et al. (1990)
57	Alexanders-fontein	24.7854	-28.8369	closed, dry	x	x		x											x	Butzer et al. (1973)

58	Ngami	22.7315	-20.4983	closed, seasonally filled	x	x								x					x		x	Shaw (1985); Shaw et al. (2003); Huntsman-Mapila et al. (2006); Burrough et al. (2007)
59	Makgadikgadi	25.4883	-20.7430	closed, dry	x	x					x			x							x	Cooke and Verstappen (1984); Shaw et al. (1997); Ringrose et al. (2005); Burrough et al. (2009); Schmidt et al. (2017)
60	Mababe	24.1585	-18.8482	closed, dry	x	x								x							x	Shaw (1985); Shaw and Cooke (1986); Shaw and Thomas (1988); Shaw and Thomas (1993); Burrough and Thomas (2008)
61	Etosha	16.3016	-18.8056	closed, dry	x	x															x	Buch and Zoller (1992); Brook et al. (2007); Brook et al. (2011); Brook et al. (2013); Hipondoka et al. (2014)
62	Tsondab	15.1606	-23.8850	closed, dry		x	x	x														Vogel and Visser (1981); Teller and Lancaster (1986); Teller et al. (1990); Stone et al. (2010)
63	Branddam-East Pan	18.8419	-26.3594	closed, dry		x		x	x													Schüller et al. (2018)
64	Omongwa Pan	19.3704	-23.7064	closed, dry		x		x	x													Schüller et al. (2018)
65	Kathu Pan	23.0083	-27.6619	closed, dry		x		x														Beaumont et al. (1984)
66	Lebatse Pan	23.8386	-23.8830	closed, dry		x		x														Holmgren and Shaw (1997)
67	Witpan	20.15	-26.6666	closed, dry		x		x														Holmgren and Shaw (1997); Telfer et al. (2009)

An uncertainty-focused database approach to extract spatiotemporal trends from qualitative and discontinuous lake-status histories – Supplementary Material

De Cort, Gijs; Chevalier, Manuel; Burrough, Sallie L.; Chen, Christine Y.; Harrison, Sandy P.

References for Table 1

- Alin, S.R., Cohen, A.S., 2003. Lake-level history of Lake Tanganyika, East Africa, for the past 2500 years based on ostracode-inferred water-depth reconstruction. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 199, 31–49.
- Ashley, G.M., Mworia, J.M., Muasya, A.M., Owen, R.B., Driese, S.G., Hover, V.C., Renaut, R.W., Goman, M.F., Mathai, S., Blatt, S.H., 2004. Sedimentation and recent history of a freshwater wetland in a semi-arid environment: Lobo Swamp, Kenya, East Africa. *Sedimentology* 51, 1301–1321.
- Avery, S., 2010. Hydrological impacts of Ethiopia's Omo Basin on Kenya's Lake Turkana water levels and fisheries. African Development Bank, Tunis.
- Bannert, O., Brinckmann, J., Kading, K. C., Knetsch, G., Kursten, M., Mayrhofer, H., 1970. Zur Geologie der Danakil Senke. *Geol. Rundsch.*, 59(2): 409-443.
- Barker, P., 1992. Differential diatom dissolution in Late Quaternary sediments from Lake Manyara, Tanzania: an experimental approach. *J. Paleolimnol.* 7, 235–251.
- Barker, P., Gasse, F., Roberts, N., Taieb, M., 1991. Taphonomy and diagenesis in diatom assemblages; a Late Pleistocene palaeoecological study from Lake Magadi, Kenya. *Hydrobiologia* 214, 267–272.
- Barker, P., Telford, R., Merdaci, O., Williamson, D., Taieb, M., Vincens, A., Gibert, E., 2000. The sensitivity of a Tanzanian crater lake to catastrophic tephra input and four millennia of climate change 10, 303–310.
- Barker, P., Telford, R., Gasse, F., Thevenon, F., 2002. Late pleistocene and holocene palaeohydrology of Lake Rukwa, Tanzania, inferred from diatom analysis. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 187, 295–305.
- Barker, P., Gasse, F., 2003. New evidence for a reduced water balance in East Africa during the Last Glacial Maximum: Implication for model-data comparison. *Quat. Sci. Rev.* 22, 823–837.

- Barker, P., Williamson, D., Gasse, F., Gibert, E., 2003. Climatic and volcanic forcing revealed in a 50,000-year diatom record from Lake Massoko, Tanzania. *Quat. Res.* 60, 368–376.
- Barry, S., Filippi, M., Talbot, M., Johnson, T., 2002. Sedimentology and geochronology of late Pleistocene and Holocene sediments from northern Lake Malawi. In: *The East African Great Lakes: Limnology, Palaeolimnology and Biodiversity*. Kluwer Academic Publishers, pp. 369–391.
- Beadle, L., 1981. *The inland waters of tropical Africa. An introduction to tropical limnology*. Longman, London.
- Beaumont, P.B., Van Zinderen Bakker Sr, E.M., Vogel, J.C. 1984. Environmental changes since 32 000 BP at Kathu Pan, northern Cape. Late Cainozoic palaeoclimates of the Southern Hemisphere. *Proc. SASQUA symposium, Swaziland, 1983*, pp. 329-338.
- Bergner, A. G. N., Trauth, M. H., Bookhagen, B. 2003. Paleoprecipitation estimates for the Lake Naivasha basin (Kenya) during the last 175 k.y. using a lake-balance model. *Glob. Planet. Change* 36, 117–136.
- Bergner, A.G.N., Trauth, M.H., 2004. Comparison of the hydrological and hydrochemical evolution of Lake Naivasha (Kenya) during three highstands between 175 and 60 kyr BP. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 215, 17–36.
- Bessemis, I., 2007. Late-Holocene climate reconstruction in equatorial East Africa: sedimentology and stable-isotope geochemistry of lake deposits. Unpubl. PhD thesis. Ghent University.
- Bessemis, I., Verschuren, D., Russell, J.M., Hus, J., Mees, F., Cumming, B.F., 2008. Palaeolimnological evidence for widespread late 18th century drought across equatorial East Africa. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 259, 107–120.
- Beuning, K.R.M., Talbot, M.R., Kelts, K., 1997. A revised 30,000-year paleoclimatic and paleohydrologic history of Lake Albert, East Africa. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 136, 259-279.
- Beuning, K.R.M., Kelts, K., Russell, J., Wolfe, B.B., 2002. Reassessment of Lake Victoria-Upper Nile River paleohydrology from oxygen isotope records of lake-sediment cellulose. *Geology* 30, 559–562.
- Bishop, W.W., 1969. Pleistocene stratigraphy of Uganda. *Geol. Surv. Uganda Mem.* X, 1-115.

- Blaauw, M., van Geel, B., Kristen, I., Plessen, B., Lyaruu, A., Engstrom, D.R., van der Plicht, J., Verschuren, D., 2011. High-resolution ¹⁴C dating of a 25,000-year lake-sediment record from equatorial East Africa. *Quat. Sci. Rev.* 30, 3043–3059.
- Bloszies, C., Forman, S.L., 2015. Potential relation between equatorial sea surface temperatures and historic water level variability for Lake Turkana, Kenya. *J. Hydrol.* 520, 489–501.
- Bloszies, C., Forman, S.L., Wright, D.K., 2015. Water level history for Lake Turkana, Kenya in the past 15,000 years and a variable transition from the African Humid Period to Holocene aridity. *Glob. Planet. Change* 132, 64–76.
- Bonatti, E., Gasperini, E., Vigliotti, L., Lupi, L., Vaselli, O., 2017. Lake Afrera, a structural depression in the Northern Afar Rift (Red Sea). *Heliyon* 3.
- Brook, G.A., Haberyan, K.A., Filippis, S., 1992. Evidence of a shallow lake at Tsodilo Hills, Botswana, 17,500 to 15,000 yr BP: further confirmation of a widespread late Pleistocene humid period in the Kalahari Desert. *Palaeoecol. Africa* 23, 165–175.
- Brook, G.A., Marais, E., Srivastava, P., Jordan, T., 2007. Timing of lake-level changes in Etosha Pan, Namibia, since the middle Holocene from OSL ages of relict shorelines in the Okondeka region. *Quaternary International* 175, 29–40.
- Brook, G.A., Railsback, L.B., Marais, E., 2011. Reassessment of carbonate ages by dating both carbonate and organic material from an Etosha Pan (Namibia) stromatolite: Evidence of humid phases during the last 20ka. *Quaternary International* 229, 24–37.
- Brook, G.A., Cherkinsky, A., Bruce Railsback, L., Marais, E., Hipondoka, M.H.T., 2013. ¹⁴C dating of organic residue and carbonate from stromatolites in Etosha Pan, Namibia: ¹⁴C reservoir effect, correction of published ages, and evidence of >8-m-deep lake during the Late Pleistocene. *Radiocarbon* 55, 1156–1163.
- Brooks, A.S., Smith, C.C., 1987. Ishango revisited: new age determinations and cultural interpretations. *Afr. Archaeol. Rev.*, 5, 65–78.
- Brooks, A.S., Hare, P.E., Kokis, J.E., Miller, G.H., Ernst, R.D., Wendorf, F., 1990. Dating pleistocene archeological sites by protein diagenesis in ostrich eggshell. *Science* (80-.). 248, 60–64.
- Brown, F.H., Fuller, C.R., 2008. Stratigraphy and tephra of the Kibish Formation, southwestern Ethiopia. *J. Hum. Evol.* 55, 366–403.

- Buch, M.W., Zöller, L., 1992. Pedostratigraphy and thermoluminescence-chronology of the western margin- (lunette-) dunes of Etosha Pan/Northern Namibia. *Würzburger Geographische Arbeiten* 84, 361-384.
- Burrough, S.L., Thomas, D.S.G., Shaw, P. a., Bailey, R.M., 2007. Multiphase Quaternary highstands at Lake Ngami, Kalahari, northern Botswana. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 253, 280–299.
- Burrough, S.L., Thomas, D.S.G., 2008. Late Quaternary lake-level fluctuations in the Mababe Depression: Middle Kalahari palaeolakes and the role of Zambezi inflows. *Quaternary Research* 69, 388-403
- Burrough, S.L., Thomas, D.S.G., Bailey, R.M., 2009. Mega-Lake in the Kalahari: A Late Pleistocene record of the Palaeolake Makgadikgadi system. *Quaternary Science Reviews* 28, 1392-1411.
- Butzer, K.W., 1971. Recent history of an Ethiopian delta: the Omo River and the level of Lake Rudolf. University of Chicago, Dept. of Geography, Chicago.
- Butzer, K.W., Isaac, G.L., Richardson, J.L., Washbourn-Kamau, C.K., 1972. Radiocarbon dating of East African lake levels. *Science* (80-). 175, 1069–1076.
- Butzer, K.W., Fock, G.J., Stuckenrath, R., Zilch, A., 1973. Paleohydrology of late Pleistocene lake Alexandersfontein, Kimberley, South Africa. *Nature* 243, 328–330.
- Carbonel, P., Grosdidier, E., Peypouquet, J.P., Tiercelin, J.J., 1983. Les Ostracodes, témoins de l'évolution hydrologique d'un lac de Rift. Exemple du lac Bogoria, Rift Gregory, Kenya. *Bull. Centres Rech. Explor. Prod. Elf-Aquitaine* 7(1), 301-13.
- Casanova, J., Hillaire-Marcel, C., 1992. Late holocene hydrological history of Lake Tanganyika, East Africa, from isotopic data on fossil stromatolites. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 91, 35–48.
- Casanova, J., Hillaire-Marcel, C., 1992b. Chronology and paleohydrology of late Quaternary high lake levels in the Manyara basin (Tanzania) from isotopic data (^{18}O , ^{13}C , ^{14}C , Th U) on fossil stromatolites. *Quat. Res.* 38, 205–226.
- Chalié, F. and Gasse, F.: Late Glacial-Holocene diatom record of water chemistry and lake level change from the tropical East African Rift Lake Abiyata (Ethiopia), *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 187(3–4), 259–283, 2002.

- Chapman, L.J., Chapman, C.A., Crisman, T.L., Nordlie, F.G., 1998. Dissolved oxygen and thermal regimes of a Ugandan crater lake. *Hydrobiologia* 385, 201–211.
- Clark, J.D., Haynes, C.V., Mawby, J.E., Gautier, A., 1970. Interim report on palaeoanthropological investigations in the Malawi Rift. *Quaternaria* 13, 305-354.
- Coetsee, J.A., 1964. Evidence for a Considerable Depression of the Vegetation Belts during the Upper Pleistocene on the East African Mountains. *Nature* 204, 564–566.
- Coetsee, J.A., 1967. Pollen analytical studies in East and Southern Africa. *Palaeoecol. Afr.* 3, 1-146.
- Cohen, A.S., Dussinger, R., Richardson, J., 1983. Lacustrine paleochemical interpretations based on Eastern and Southern african ostracodes. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 43, 129–151.
- Cohen, A.S., Talbot, M.R., Awramik, S.M., Dettman, D.L., Abell, P., 1997. Lake level and paleoenvironmental history of Lake Tanganyika, Africa, as inferred from late Holocene and modern stromatolites. *Geol. Soc. Am. Bull.* 109, 444–460.
- Cohen, A.S., Palacios-Fest, M.R., Msaky, E.S., Alin, S.R., McKee, B., O'Reilly, C.M., Dettman, D.L., Nkotagu, H., Lezzar, K.E., 2005. Paleolimnological investigations of anthropogenic environmental change in Lake Tanganyika: IX. Summary of paleorecords of environmental change and catchment deforestation at Lake Tanganyika and impacts on the Lake Tanganyika ecosystem. *J. Paleolimnol.* 34, 125–145.
- Cooke, H.J., T. Verstappen, H., 1984. The landforms of the western Makgadikgadi basin in northern Botswana, with a consideration of the chronology of the evolution of Lake Palaeo-Makgadikgadi. *Zeitschrift fur Geomorphologie* 28, 1-19.
- Costa, K., Russell, J., Konecky, B., Lamb, H., 2014. Isotopic reconstruction of the African Humid Period and Congo Air Boundary migration at Lake Tana, Ethiopia. *Quat. Sci. Rev.* 83, 58–67.
- Damnati, B., Taieb, M., 1995. Solar and ENSO signatures in laminated deposits from lake Magadi (Kenya) during the Pleistocene/Holocene transition. *J. African Earth Sci.* 21, 373–382.
- Damnati, B., Taieb, M., 1996. L évolution hydrologique du lac Sonachi (Kenya), à l Holocene (7400-0 ans BP). *C.R. Acad. Sci. Paris, t. 322, série II a*, 141-148.
- De Cort, G., Bessems, I., Keppens, E., Mees, F., Cumming, B., Verschuren, D., 2013. Late-Holocene and recent hydroclimatic variability in the central Kenya Rift Valley: The sediment record of hypersaline lakes Bogoria, Nakuru and Elementeita. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 388, 69–80.

- De Cort, G., Verschuren, D., Ryken, E., Wolff, C., Renaut, R.W., Creutz, M., Van der Meeren, T., Haug, G., Olago, D.O., Mees, F., 2018. Multi-basin depositional framework for moisture-balance reconstruction during the last 1300 years at Lake Bogoria, central Kenya Rift Valley. *Sedimentology* 38, 42–49.
- de Heinzelin, J., Verniers, J., 1996. Realm of the upper Semliki (eastern Zaire): an essay on historical geology. *Ann. Kon. Mus. Midd.-Afr.* 102, 3-83.
- Delalande, M., Bergonzini, L., Beal, F., Garcin, Y., Majule, A., Williamson, D., 2005. Contribution to the detection of Lake Masoko (Tanzania) groundwater outflow: isotopic evidence (^{18}O , D). *IAHS Publ.* 50, 867-880.
- Delibrias, G., Guillier, M., Labeyrie, J., 1974. GIF natural radiocarbon measurements VIII 16, 15–94.
- Delvaux, D., Kervyn, F., Vittori, E., Kajara, R.S.A., Kilembe, E., 1998. Late Quaternary tectonic activity and lake level change in the Rukwa Rift Basin. *J. African Earth Sci.* 26, 397–421.
- Driese, S.G., Ashley, G.M., Li, Z.H., Hover, V.C., Owen, R.B., 2004. Possible late Holocene equatorial palaeoclimate record based upon soils spanning the Medieval Warm Period and Little Ice Age, Lobo Plain, Kenya. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 213, 231–250.
- Dühnforth, M., Bergner, A.G.N., Trauth, M.H., 2006. Early Holocene water budget of the Nakuru-Elmenteita basin, Central Kenya Rift. *J. Paleolimnol.* 36, 281–294.
- Ekblom, A., Stabell, B., 2008. Paleohydrology of Lake Nhaucati (southern Mozambique), ~400 AD to present. *J. Paleolimnol.* 40, 1127–1141.
- Ekblom, A., Gillson, L., Risberg, J., Holmgren, K., Chidoub, Z., 2012. Rainfall variability and vegetation dynamics of the lower Limpopo Valley, Southern Africa, 500 AD to present. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 363–364, 69–78.
- Felton, A.A., Russell, J.M., Cohen, A.S., Baker, M.E., Chesley, J.T., Lezzar, K.E., McGlue, M.M., Pigati, J.S., Quade, J., Curt Stager, J., Tiercelin, J.J., 2007. Paleolimnological evidence for the onset and termination of glacial aridity from Lake Tanganyika, Tropical East Africa. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 252, 405–423.
- Filippi, M., Talbot, M., 2005. The palaeolimnology of northern Lake Malawi over the last 25ka based upon the elemental and stable isotopic composition of sedimentary organic matter. *Quat. Sci. Rev.* 24, 1303–1328.

- Finney, B.P., Johnson, T.C., 1991. Sedimentation in Lake Malawi (East Africa) during the past 10,000 years: a continuous paleoclimatic record from the southern tropics. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 85, 351–366.
- Foerster, V., Junginger, A., Langkamp, O., Gebru, T., Asrat, A., Umer, M., Lamb, H.F., Wennrich, V., Rethemeyer, J., Nowaczyk, N., Trauth, M.H., Schaebitz, F., 2012. Climatic change recorded in the sediments of the Chew Bahir basin, southern Ethiopia, during the last 45,000 years. *Quat. Int.* 274, 25–37.
- Forman, S.L., Wright, D.K., Blois, C., 2014. Variations in water level for Lake Turkana in the past 8500 years near Mt. Porro, Kenya and the transition from the African Humid Period to Holocene aridity. *Quat. Sci. Rev.* 97, 84–101.
- Garcin, Y., Williamson, D., Taieb, M., Vincens, A., Mathé, P.E., Majule, A., 2006. Centennial to millennial changes in maar-lake deposition during the last 45,000 years in tropical Southern Africa (Lake Masoko, Tanzania). *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 239, 334–354.
- Garcin, Y., Vincens, A., Williamson, D., Guiot, J., Buchet, G., 2006b. Wet phases in tropical southern Africa during the last glacial period. *Geophys. Res. Lett.* 33, 1–4.
- Garcin, Y., Williamson, D., Bergonzini, L., Radakovitch, O., Vincens, A., Buchet, G., Guiot, J., Brewer, S., Mathé, P.E., Majule, A., 2007. Solar and anthropogenic imprints on Lake Masoko (southern Tanzania) during the last 500 years. *J. Paleolimnol.* 37, 475–490.
- Garcin, Y., Vincens, A., Williamson, D., Buchet, G., Guiot, J., 2007b. Abrupt resumption of the African Monsoon at the Younger Dryas-Holocene climatic transition. *Quat. Sci. Rev.* 26, 690–704.
- Garcin, Y., Junginger, A., Melnick, D., Olago, D.O., Strecker, M.R., Trauth, M.H., 2009. Late Pleistocene-Holocene rise and collapse of Lake Suguta, northern Kenya Rift. *Quat. Sci. Rev.* 28, 911–925.
- Garcin, Y., Melnick, D., Strecker, M.R., Olago, D., Tiercelin, J.J., 2012. East African mid-Holocene wet-dry transition recorded in palaeo-shorelines of Lake Turkana, northern Kenya Rift. *Earth Planet. Sci. Lett.* 331–332, 322–334.
- Gasse, F., 1977. Evolution of Lake Abhé (Ethiopia and TFAI), from 70,000 b.p. *Nature* 265, 42–45.
- Gasse, F., Van Campo, E., 1998. A 40,000-yr pollen and diatom record from Lake Tritrivakely, Madagascar, in the southern tropics. *Quaternary Research*, 49(3), 299–311.

- Gasse, F., Van Campo, E., 2001. Late Quaternary environmental changes from a pollen and diatom record in the southern tropics (Lake Tritrivakely, Madagascar). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 167(3), 287-308.
- Gasse, F., Fontes, J., Rognon, P., 1974. Variations hydrologiques et extension des lacs Holocenes du desert Danakil. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 15, 109–148.
- Gasse, F., Street, F.A., 1978. Late Quaternary lake-level fluctuations and environments of the northern Rift Valley and Afar Region (Ethiopia and Djibouti). *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 24, 279-325.
- Gasse, F., Lédée, V., Massault, M., Fontes, J.-C., 1989. Water-level fluctuations of Lake Tanganyika in phase with oceanic changes during the last glaciation and deglaciation. *Nature* 342, 57–59.
- Gasse, F., Cortijo, E., Disnar, J. R., Ferry, L., Gibert, E., Kissel, C., Laggoun-Défarge, F., Lallier-Vergès, E., Miskovsky, J. C., Ratsimbazafy, B., Ranaivo, F., Tucholka, P., Saos, J. L., Siffedine, A., Taieb, M., Van Campo, E., and Williamson, D., 1994. A 36 kyr environmental record in the southern tropics: Lake Tritrivakely. *Comptes Rendus de l'Académie des Sciences (Paris) Série 2* 318, 1513–1519.
- Gasse, F., Barker, P., Johnson, T., 2002. A 24,000 yr diatom record from the northern basin of Lake Malawi. In: Odada, E., Olago, D. (Eds.), *The East African Great Lakes: Limnology, Palaeolimnology and Biodiversity*. Kluwer Academic Publishers, pp. 393–414.
- Ghinassi, M., D’Oriano, F., Benvenuti, M., Awramik, S., Bartolini, C., Fedi, M., Ferrari, G., Papini, M., Sagri, M., Talbot, M., 2012. Shoreline fluctuations of Lake Hayk (northern Ethiopia) during the last 3500 years: Geomorphological, sedimentary, and isotope records. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 365–366, 209–226.
- Ghinassi, M., D’oriano, F., Benvenuti, M., Fedi, M., Awramik, S., 2015. Lacustrine Facies In Response To Millennial-Century-Scale Climate Changes (Lake Hayk, Northern Ethiopia). *J. Sediment. Res.* 85, 381–398.
- Gibert, E., Bergonzini, L., Massault, M., Williamson, D., 2002. AMS-14C chronology of 40.0 cal ka BP continuous deposits from a crater lake (Lake Massoko, Tanzania) modern water balance and environmental implications. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 187, 307–322.
- Gillespie, R., Street-Perrott, F.A., Switsur, R., 1983. Post-glacial arid episodes in Ethiopia have implications for climate prediction. *Nature* 306, 680–683.
- Gillson, L., Ekblom, A., 2009. Untangling anthropogenic and climatic influence on riverine forest in the Kruger National Park, South Africa. *Veg. Hist. Archaeobot.* 18, 171–185.

- Greenway, P., DF, V.-F., 1969. The vegetation of Lake Manyara National Park. *J. Ecol.* 57, 127–149.
- Greenwood, P.H., 1976. Lake George, Uganda. *Philos. Trans. R. Soc. B Biol. Sci.* 274, 375–391.
- Grove, A.T., Street, A.F., Goudie, A.S., 1975. Former Lake Levels and Climatic Change in the Rift Valley of Southern Ethiopia Author (s): A . T . Grove , F . Alayne Street , A . S . Goudie Published by : Blackwell Publishing on behalf of The Royal Geographical Society (with the Institute of British. *Geogr. J.* 141, 177–194.
- Haberyan, K.A., 1987. Fossil diatoms and the paleolimnology of Lake Rukwa, Tanzania. *Freshw. Biol.* 17, 429–436.
- Haberyan, K.A., Hecky, R.E., 1987. The late pleistocene and holocene stratigraphy and paleolimnology of Lakes Kiva and Tanganyika. *Paleoecology* 61, 169–197.
- Haberyan, K.A., 2018. A 22,000 yr diatom record from the plateau of Zambia. *Quat. Res.* 89, 33–42.
- Halfman, J.D., Johnson, T.C., 1988. High-resolution record of cyclic climatic change during the past 4 ka from Lake Turkana, Kenya. *Geology* 16, 496.
- Halfman, J., Jacobson, D., Cannella, C., Haberyan, K., Finney, B., 1992. Fossil diatoms and the mid to late holocene paleolimnology of Lake Turkana, Kenya: a reconnaissance study. *J. Paleolimnol.* 7, 23–35.
- Halfman, J.D., Johnson, T.C., Finney, B.P., 1994. New AMS dates, stratigraphic correlations and decadal climatic cycles for the past 4 ka at Lake Turkana, Kenya. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 111, 83–98.
- Harvey, T.J., 1976. The Paleolimnology of Lake Mobutu Sese Seko, Uganda-Zaire : the Last 28,000 Years. Ph.D.thesis, Duke University, Durham, North Carolina,113pp.
- Hecky, R.E., Degens, E.T., 1973. Late Pleistocene-Holocene chemical stratigraphy and paleolimnology of the rift valley lakes of Central Africa. Woods Hole Oceanogr. Inst., WHOI-73-28, unpublished manuscript.
- Helgren, D.M., Brooks, A.S., 1983. Geoarchaeology at Gi, a middle stone age and later stone age site in the Northwest Kalahari. *J. Archaeol. Sci.* 10, 181–197.
- Hickley, P., Boar, R.R., Mavuti, K.M., 2003. Bathymetry of Lake Bogoria, Kenya. *J. East African Nat. Hist.* 92, 107–117.

- Hillaire-Marcel, C., Carro, O., Casanova, J., 1986. ^{14}C and Th/U dating of Pleistocene and Holocene stromatolites from East African paleolakes. *Quat. Res.* 25, 312–329.
- Hillaire-Marcel, C., Casanova, J., 1987. Isotopic hydrology and paleohydrology of the Madagi (Kenya)-Natron (Tanzania) basin during the late quaternary. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 58, 155–181.
- Hipondoka, M.H.T., Mauz, B., Kempf, J., Packman, S., Chiverrell, R.C., Bloemendal, J., 2014. Chronology of sand ridges and the Late Quaternary evolution of the Etosha Pan, Namibia. *Geomorphology* 204, 553–563.
- Holdship, S.A., 1976. The paleolimnology of Lake Manyara, Tanzania: a diatom analysis of a 56 meter sediment core. MSc thesis, Duke University, Durham, USA.
- Holmgren, K., Shaw, P. 1997. Palaeoenvironmental reconstruction from near-surface pan sediments: An example from Lebatse Pan, southeast Kalahari, Botswana. *Geografiska Annaler, Series A: Physical Geography*, 79, 83–93.
- Holmgren, K., Risberg, J., Freudendahl, J., Achimo, M., Ekblom, A., Mugabe, J., Norström, E., Siteo, S.R., 2012. Water-level variations in Lake Nhauhache, Mozambique, during the last 2,300 years. *J. Paleolimnol.* 48, 311–322.
- Huang, Y., Street-Perrott, F.A., Perrott, R.A., Metzger, P., Eglinton, G., 1999. Glacial-interglacial environmental changes inferred from molecular and compound-specific $\delta^{13}\text{C}$ analyses of sediments from Sacred Lake, Mt. Kenya. *Geochim. Cosmochim. Acta* 63, 1383–1404.
- Hughes, L., 2008. Mining the Maasai Reserve: The Story of Magadi. *J. East. African Stud.* 2, 134–164.
- Huntsman-Mapila, P., Ringrose, S., Mackay, A.W., Downey, W.S., Modisi, M., Coetzee, S.H., Tiercelin, J.-J., Kampunzu, A.B., Vanderpost, C., 2006. Use of the geochemical and biological sedimentary record in establishing palaeo-environments and climate change in the Lake Ngami basin, NW Botswana. *Quat. Int.*
- Johnson, O., Scholz, J.J.M. van, Talbot, F., Kelts, R.B., Ricketts, T.C., Ngobi, A., Beuning, T.D., Ssemmanda, P., McGill, A.C., Barel, C.D.N., Coulter, G.W., Meer, H.J. van der, Bowmaker, J.K., Zahavi, A., Lande, R., Hamilton, W.D., Zuk, M., Kirkpatrick, M., Basolo, A.L., Grafen, A., Kirkpatrick, M., Ryan, M.J., Archer, S.N., Endler, J.A., Lythgoe, J.N., Patridge, J.C., Turner, G.F., Burrows, M.T., Endler, J.A., Houde, A.E., Endler, J.A., ____, Zahavi, A., Pomiankowski, A.J., Møller, A.P., Luyten, P.H., Liley, N.R., Reimchen, T.E., Endler, J.A., 1996. Late Pleistocene Desiccation of Lake Victoria and Rapid Evolution of Cichlid Fishes. *Science* 273, 1091–3.

- Johnson, T.C., Halfman, J.D., Showers, W.J., 1991. Paleoclimate of the past 4000 years at Lake Turkana, Kenya, based on the isotopic composition of authigenic calcite. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 85, 189–198.
- Johnson, T.C., Chan, Y., Beuning, K.R.M., Kelts, K., Ngobi, G., Verschuren, D. 1998. Biogenic silica profiles in Holocene cores from Lake Victoria: implications for lake level history and initiation of the Victoria Nile. In: Lehman (ed.) *Environmental change and response in East African lakes*. Kluwer, Dordrecht, pp. 75-89.
- Johnson, T.C., Kelts, K., Odada, E., 2000. The Holocene History of Lake Victoria 29, 2–11.
- Johnson, T.C., Barry, S.L., Chan, Y., Wilkinson, P., 2001. Decadal record of climate variability spanning the past 700 yr in the Southern Tropics of East Africa. *Geology* 29, 83.
- Johnson, T.C., McCave, I.N., 2008. Transport mechanism and paleoclimatic significance of terrigenous silt deposited in varved sediments of an African rift lake. *Limnol. Oceanogr.* 53, 1622–1632.
- Junginger, A., Roller, S., Olaka, L.A., Trauth, M.H., 2014. The effects of solar irradiation changes on the migration of the Congo Air Boundary and water levels of paleo-Lake Suguta, Northern Kenya Rift, during the African Humid Period (15-5ka BP). *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 396, 1–16.
- Kendall, R.L., 1969. An Ecological History of the Lake Victoria Basin. *Ecol. Monogr.* 39, 121–176.
- Kiage, L.M., Liu, K.B., 2009. Paleoenvironmental changes in the lake Baringo Basin, Kenya, East Africa since AD 1650: Evidence from the paleorecord. *Prof. Geogr.* 61, 438–458.
- Kiage, L.M., Liu, K.B., 2009b. Palynological evidence of climate change and land degradation in the Lake Baringo area, Kenya, East Africa, since AD 1650. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 279, 60–72.
- Kirsten, K.L., Meadows, M.E., 2016. Late-Holocene palaeolimnological and climate dynamics at Princessvlei, South Africa: Evidence from diatoms. *Holocene* 26, 1371–1381.
- Kirsten, K.L., Kasper, T., Cawthra, H.C., Strobel, P., Quick, L.J., Meadows, M.E., Haberzettl, T., 2020. Holocene variability in climate and oceanic conditions in the winter rainfall zone of South Africa— inferred from a high resolution diatom record from Verlorenvlei. *J. Quat. Sci.* 1–10.

- Konecky, B., Russell, J., Huang, Y., Vuille, M., Cohen, L., Street-Perrott, F.A., 2014. Impact of monsoons, temperature, and CO₂ on the rainfall and ecosystems of Mt. Kenya during the Common Era. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 396, 17–25.
- Laerdal, T., Talbot, M.R., Russell, J.M., 2002. Late Quaternary sedimentation and climate in the lakes Edward and George area, Uganda-Congo. In: Odada, E.O., Olago, D.O. (eds.), *The East African Great Lakes: limnology, palaeolimnology, biodiversity*. Kluwer Academic, Dordrecht, pp. 429–470.
- Lamb, A.L., Leng, M.J., Lamb, H.F., Mohammed, M.U., 2000. A 9000-year oxygen and carbon isotope record of hydrological change in a small Ethiopian crater lake. *The Holocene* 10, 167–177.
- Lamb, A.L., Leng, M.J., Mohammed, M.U., Lamb, H.F., 2004. Holocene climate and vegetation change in the Main Ethiopian Rift Valley, inferred from the composition (C/N and $\delta^{13}\text{C}$) of lacustrine organic matter. *Quat. Sci. Rev.* 23, 881–891.
- Lamb, A.L., Leng, M.J., Sloane, H.J., Telford, R.J., 2005. A comparison of the palaeoclimate signals from diatom oxygen isotope ratios and carbonate oxygen isotope ratios from a low latitude crater lake. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 223, 290–302.
- Lamb, H.F., Leng, M.J., Telford, R.J., Ayenew, T., Umer, M., 2007. Oxygen and carbon isotope composition of authigenic carbonate from an Ethiopian lake: a climate record of the last 2000 years. *The Holocene* 17, 517–526.
- Lamb, H.F., Bates, C.R., Coombes, P. V., Marshall, M.H., Umer, M., Davies, S.J., Dejen, E., 2007b. Late Pleistocene desiccation of Lake Tana, source of the Blue Nile. *Quat. Sci. Rev.* 26, 287–299.
- Lancaster I.N., 1979. Evidence for a widespread late Pleistocene humid period in the Kalahari. *Nature* 279, 145–146.
- Legesse, D., Gasse, F., Radakovitch, O., Vallet-Coulomb, C., Bonnefille, R., Verschuren, D., Gibert, E., Barker, P., 2002. Environmental changes in a tropical lake (Lake Abiyata, Ethiopia) during recent centuries. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 187, 233–258.
- Lehman, J.T., 1998. *Environmental change and response in East African lakes*. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp. 236.
- Livingstone, D.A., 1965. Sedimentation and the history of water level change in Lake Tanganyika. *Limnol. Oceanogr.* 10, 607–610.
- Livingstone, D.A., 1971. A 22 000-year pollen record from the plateau of Zambia. *Limnol. Oceanogr.* 16, 349–356.

- Loomis, S.E., Russell, J.M., Ladd, B., Street-Perrott, F.A., Sinninghe Damsté, J.S., 2012. Calibration and application of the branched GDGT temperature proxy on East African lake sediments. *Earth Planet. Sci. Lett.* 357–358, 277–288.
- Marshall, M.H., Lamb, H.F., Davies, S.J., Leng, M.J., Kubsa, Z., Umer, M., Bryant, C., 2009. Climatic change in northern Ethiopia during the past 17,000 years: A diatom and stable isotope record from Lake Ashenge. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 279, 114–127.
- Marshall, M.H., Lamb, H.F., Huws, D., Davies, S.J., Bates, R., Bloemendal, J., Boyle, J., Leng, M.J., Umer, M., Bryant, C., 2011. Late Pleistocene and Holocene drought events at Lake Tana, the source of the Blue Nile. *Glob. Planet. Change* 78, 147–161.
- McCall, J., 2010. Lake Bogoria, Kenya: Hot and warm springs, geysers and Holocene stromatolites. *Earth-Science Rev.* 103, 71–79.
- Mills, K., Ryves, D.B., 2012. Diatom-based models for inferring past water chemistry in western Ugandan crater lakes. *J. Paleolimnol.* 48, 383–399.
- Mills, K., Ryves, D.B., Anderson, N.J., Bryant, C.L., Tyler, J.J., 2014. Expressions of climate perturbations in western ugandan crater lake sediment records during the last 1000 years. *Clim. Past* 10, 1581–1601.
- Moernaut, J., Verschuren, D., Charlet, F., Kristen, I., Fagot, M., De Batist, M., 2010. The seismic-stratigraphic record of lake-level fluctuations in Lake Challa: Hydrological stability and change in equatorial East Africa over the last 140 kyr. *Earth Planet. Sci. Lett.* 290, 214–223.
- Mohammed, M.U., Bonnefille, R., Johnson, T.C., 1996. Pollen and isotopic records in Late Holocene sediments from Lake Turkana, Kenya. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 119, 371–383.
- Morrissey, A., Scholz, C.A., 2014. Paleohydrology of Lake Turkana and its influence on the Nile River system. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 403, 88–100.
- Musisi, J.H., 1991. The Neogene-Quaternary geology of the Lake-George-Edward Basin. Unpublished PhD thesis, Vrije Universiteit Brussel, Belgium.
- Muzuka, A.N.N., Ryner, M., Holmgren, K., 2004. 12,000-Year, preliminary results of the stable nitrogen and carbon isotope record from the Empakai Crater lake sediments, Northern Tanzania. *J. African Earth Sci.* 40, 293–303.

- Neumann, F.H., Stager, J.C., Scott, L., Venter, H.J.T., Weyhenmeyer, C., 2008. Holocene vegetation and climate records from Lake Sibaya, KwaZulu-Natal (South Africa). *Rev. Palaeobot. Palynol.* 152, 113–128.
- Neumann, F.H., Scott, L., Bamford, M.K., 2011. Climate change and human disturbance of Fynbos vegetation during the late Holocene at Princess Vlei, Western Cape, South Africa. *The Holocene* 21, 1137–1149.
- Nicholson, S.E., 1998. Historical fluctuations of Lake Victoria and other lakes in the northern Rift Valley of East Africa. In: Lehman, J. (Ed.), *Environmental Change and Response in East African Lakes*. Kluwer Academic Publishers, pp. 7–35.
- Nicholson, S.E., 1999. Historical and Modern Fluctuations of Lakes Tanganyika and Rukwa and Their Relationship to Rainfall Variability. *Clim. Change* 41, 53–71.
- Norström, E., Öberg, H., Siteo, S.R., Ekblom, A., Westerberg, L.-O., Risberg, J., 2018. Vegetation dynamics within the savanna biome in southern Mozambique during the late Holocene. *The Holocene* 28, 277–292.
- Norström, E., Norén, G., Smittenberg, R.H., Massuanganhe, E.A., Ekblom, A., 2018b. Leaf wax δD inferring variable medieval hydroclimate and early initiation of Little Ice Age (LIA) dryness in southern Mozambique. *Glob. Planet. Change* 170, 221–233.
- Obando, J.A., Onywere, S., Shisanya, C., Ndubi, A., Masiga, D., Irura, Z., Mariita, N., Maragia, H., 2016. Impact of Short-Term Flooding on Livelihoods in the Kenya Rift Valley Lakes, *Geomorphology and Society, Advances in Geographical and Environmental Sciences*. Springer Japan.
- Öberg, H., Andersen, T.J., Westerberg, L.-O., Risberg, J., Holmgren, K., 2012. A diatom record of recent environmental change in Lake Duluti, northern Tanzania. *J. Paleolimnol.* 48, 401–416.
- Öberg, H., Norström, E., Malmström Ryner, M., Holmgren, K., Westerberg, L.-O., Risberg, J., Eddudóttir, S.D., Andersen, T.J., Muzuka, A., 2013. Environmental variability in northern Tanzania from AD 1000 to 1800, as inferred from diatoms and pollen in Lake Duluti. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 374, 230–241.
- Olago, D.O., Street-Perrott, F.A., Perrott, R.A., Ivanovich, M., Harkness, D.D., 1999. Late Quaternary glacial-interglacial cycle of climatic and environmental change on Mount Kenya, Kenya. *J. African Earth Sci.* 29, 593–618.

- Olago, D.O., Street-Perrott, F.A., Perrott, R.A., Ivanovich, M., Harkness, D.D., Odada, E.O., 2000. Long-term temporal characteristics of palaeomonsoon dynamics in equatorial Africa. *Glob. Planet. Change* 26, 159–171.
- Olago, D., Street-Perrott, F., Perrott, R., Ivanovich, M., Harkness, D., 2009. EU/Th AND 14C isotope dating of lake sediments from sacred lake and lake Nkunga, Kenya. *African J. Sci. Technol.* 2, 36–46.
- Onyando, J.O., Musila, F., Awer, M., 2005. The use of GIS and remote sensing techniques to analyse water balance of Lake Bogoria under limited data conditions. *J. Civ. Eng. Res. Pract.* 2, 53–65.
- Owen, R.B., Barthelme, J.W., Renaut, R.W., Vincens, A., 1982. Palaeolimnology and archaeology of Holocene deposits north-east of Lake Turkana, Kenya. *Nature* 298, 523–529.
- Owen, R.B., Crossley, R., Johnson, T.C., Tweddle, D., Kornfield, I., Davison, S., Eccles, D.H., Engstrom, D.E., 1990. Major Low Levels of Lake Malawi and their Implications for Speciation Rates in Cichlid Fishes. *Proc. R. Soc. B Biol. Sci.* 240, 519–553.
- Owen, R.B., Crossley, R., 1990. Recent sedimentation in lakes Chilwa and Chiuta, Malawi. *Palaeoecol. Afr* 20, 109–117.
- Payne, B.R., 1970. Water balance of Lake Chala and its relation to groundwater from tritium and stable isotope data. *J. Hydrol.* 11, 47–58.
- Phillipson, D.W., 1977. *The later prehistory of Eastern and Southern Africa*. Heinemann, London.
- Renaut, R.W., Owen, R.B., 2000. Lake Baringo, Kenya Rift Valley, and its Pleistocene Precursors. In: Gierlowski-Kordesch, E.H., Kelts, K.R. (Eds.), *Lake Basins through Space and Time: AAPG Studies in Geology*.
- Richardson, J. L., Richardson, A. E., 1972. History of an African Rift Lake and Its Climatic Implications. *Ecol. Monogr.* 42, 499–534.
- Richardson, J. L., Dussinger, R. A., 1986. Paleolimnology of mid-elevation lakes in the Kenya Rift Valley. *Hydrobiologia* 143, 167–174 (1986).
- Ricketts, R.D., Johnson, T.C., 1996. Climate change in the Turkana basin as deduced from a 4000 year long δO_{18} record. *Earth Planet. Sci. Lett.* 142, 7–17.
- Ringrose, S., Huntsman-Mapila, P., Kampunzu, A.B., Downey, W., Coetzee, S., Vink, B., Matheson, W., Vanderpost, C., 2005. Sedimentological and geochemical evidence for palaeo-environmental

- change in the Makgadikgadi subbasin, in relation to the MOZ rift depression, Botswana. *Palaeogeography, Palaeoclimatology, Palaeoecology* 217, 265–287.
- Robbins, L.H., 1972. Archeology in the Turkana District, Kenya. *Science* (80-.). 176, 359–366.
- Roberts, N., Taieb, M., Barker, P., Damnati, B., Icole, M., Williamson, D., 1993. Timing of the Younger Dryas event in East Africa from lake level changes. *Nature* 366, 146–148.
- Russell, J.M., Johnson, T.C., 2005. A high-resolution geochemical record from Lake Edward, Uganda Congo and the timing and causes of tropical African drought during the late Holocene. *Quat. Sci. Rev.* 24, 1375–1389.
- Russell, J.M., Johnson, T.C., 2007. Little ice age drought in equatorial Africa: Intertropical convergence zone migrations and El Niño-Southern oscillation variability. *Geology* 35, 21–24.
- Russell, J.M., Johnson, T.C., Kelts, K.R., Lærdal, T., Talbot, M.R., 2003. An 11 000-year lithostratigraphic and paleohydrologic record from Equatorial Africa: Lake Edward, Uganda-Congo. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 193, 25–49.
- Russell, J.M., Verschuren, D., Eggermont, H., 2007. Spatial complexity of “Little Ice Age” climate in East Africa: sedimentary records from two crater lake basins in western Uganda. *The Holocene* 17, 183–193.
- Ryner, M., Gasse, F., Rumes, B., Verschuren, D., 2007. Climatic and hydrological instability in semi-arid equatorial East Africa during the late Glacial to Holocene transition: A multi-proxy reconstruction of aquatic ecosystem response in northern Tanzania. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 248, 440–458.
- Ryner, M., Holmgren, K., Taylor, D., 2008. A record of vegetation dynamics and lake level changes from Lake Emakat, northern Tanzania, during the last c. 1200 years. *J. Paleolimnol.* 40, 583–601.
- Ryves, D.B., Mills, K., Bennike, O., Brodersen, K.P., Lamb, A.L., Leng, M.J., Russell, J.M., Ssemmanda, I., 2011. Environmental change over the last millennium recorded in two contrasting crater lakes in western Uganda, eastern Africa (Lakes Kasenda and Wandakara). *Quat. Sci. Rev.* 30, 555–569.
- Saulnier-Talbot, É., Chapman, L.J., Efitre, J., Simpson, K.G., Gregory-Eaves, I., 2018. Long-Term Hydrologic Fluctuations and Dynamics of Primary Producers in a Tropical Crater Lake. *Front. Ecol. Evol.* 6.
- Schmidt, M., Fuchs, M., Henderson, A.C.G., Kossler, A., Leng, M.J., Mackay, A.W., Shemang, E., Riedel, F., 2017. Paleolimnological features of a mega-lake phase in the Makgadikgadi Basin

- (Kalahari, Botswana) during Marine Isotope Stage 5 inferred from diatoms. *Journal of Paleolimnology* 58, 373-390.
- Scholz, C., King, J., Ellis, G., Swart, P.K., Stager, J.C., Colman, S.M., 2003. Paleolimnology of Lake Tanganyika, East Africa, over the past 100 kyr. *J. Paleolimnol.* 30, 139–150.
- Scholz, C.A., Cohen, A.S., Johnson, T.C., King, J., Talbot, M.R., Brown, E.T., 2011. Scientific drilling in the Great Rift Valley: The 2005 Lake Malawi Scientific Drilling Project — An overview of the past 145,000 years of climate variability in Southern Hemisphere East Africa. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 303, 3–19.
- Scholz, C.A., Rosendahl, B.R., 1988. Low lake stands in Lake Malawi and Tanganyika East Africa, delineated with multifold seismic data. *Science* (80-.). 240, 1645–1648.
- Scholz, C.A., Finney, B.P., 1994. Late Quaternary sequence stratigraphy of Lake Malawi (Nyasa), Africa. *Sedimentology* 41, 163–179.
- Schüller, I., Belz, L., Wilkes, H., Wehrmann, A. 2018. Late Quaternary shift in southern African rainfall zones: Sedimentary and geochemical data from Kalahari pans. *Zeitschrift für Geomorphologie*, 61, 339-362.
- Sene, K.J., Plinston, D.T. 1994. A review and update of the hydrology of Lake Victoria in East Africa. *Hydrological Sciences Journal* 39, 47-63.
- Shaw, P.A., 1985. Late quaternary landforms and environmental change in northwest Botswana: The evidence of Lake Ngami and the Mababe Depression. *Trans. Inst. Br. Geogr.*
- Shaw, P.A., Cooke, H.J. 1986. Geomorphic evidence for the late Quaternary palaeoclimates of the middle Kalahari of northern Botswana. *CATENA* 13, 349-359.
- Shaw, P., Thomas, D.S.G., 1988. Lake Caprivi: a late Quaternary link between the Zambezi and middle Kalahari drainage systems. *Zeitschrift für Geomorphologie* 32, 329-337.
- Shaw, P., Thomas, D.S.G., 1993. Geomorphological Processes, Environmental Change and Landscape Sensitivity in the Kalahari Region of Southern Africa., In: Thomas, D.S.G., Allison, R.J. (Eds.), *Landscape Sensitivity*. John Wiley and Sons Ltd, pp. 83-95.
- Shaw, P.A., Davies, F.B.M., Stokes, S., Thomas, D.S.G., Holmgren, K., 1997. Palaeoecology and age of a Quaternary high lake level in the Makgadikgadi Basin of the Middle Kalahari, Botswana. *South African Journal of Science* 93, 273-276.

- Shaw, P.A., Bateman, M.D., Thomas, D.S.G., Davies, F., 2003. Holocene fluctuations of Lake Ngami, Middle Kalahari: Chronology and responses to climatic change. *Quat. Int.* 111(1), 23-35.
- Sitoe, S.R., Risberg, J., Norström, E., Westerberg, L.O., 2017. Late Holocene sea-level changes and paleoclimate recorded in Lake Lungué, southern Mozambique. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 485, 305–315.
- Specht, T., Rosendahl, B., 1989. Architecture of the Lake Malawi Rift, East Africa. *J. African Earth Sci.* 8, 355–382.
- Ssemmanda, I., Ryves, D.B., Bennike, O., Appleby, P.G., 2005. Vegetation history in western Uganda during the last 1200 years: a sediment-based reconstruction from two crater lakes. *The Holocene* 15, 119–132.
- Stager, J.C. 1984. The diatom record of Lake Victoria (East Africa): The last 17,000 years. In: Mann (ed.) *Proceedings of the Seventh International Diatom Symposium*, Philadelphia, Strauss & Cramer, pp. 455-476
- Stager, J.C., 1988. Environmental Changes at Lake Cheshi, Zambia since 40,000 Years B.P. *Quat. Res.* 29, 54–65.
- Stager, J.C., Reinthal, P.N., Livingstone, D.A., 1986. A 25,000-year history for Lake Victoria, East Africa, and some comments on its significance for the evolution of cichlid fishes. *Freshw. Biol.* 16, 15–19.
- Stager, J.C., Cumming, B., Meeker, L., 1997. A High-Resolution 11,400-Yr Diatom Record from Lake Victoria, East Africa. *Quat. Res.* 47, 81–89.
- Stager, J.C., Mayewski, P.A., Meeker, L.D., 2002. Cooling cycles, Heinrich event 1, and the desiccation of Lake Victoria. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 183, 169–178.
- Stager, J.C., Cumming, B.F., Meeker, L.D., 2003. A 10,000-year high-resolution diatom record from Pilkington Bay, Lake Victoria, East Africa. *Quat. Res.* 59, 172–181.
- Stager, J.C., Westwood, J., Grzesik, D., Cumming, B.F., 2005. A 5500-year environmental history of Lake Nabugabo, Uganda. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 218, 347–354.
- Stager, J.C., Ryves, D., Cumming, B.F., David Meeker, L., Beer, J., 2005b. Solar variability and the levels of Lake Victoria, East Africa, during the last millenium. *J. Paleolimnol.* 33, 243–251.
- Stager, J.C., Cocquyt, C., Bonnefille, R., Weyhenmeyer, C., Bowerman, N., 2009. A late Holocene paleoclimatic history of Lake Tanganyika, East Africa. *Quat. Res.* 72, 47–56.

- Stager, J.C., Ryves, D.B., King, C., Madson, J., Hazzard, M., Neumann, F.H., Maud, R., 2013. Late Holocene precipitation variability in the summer rainfall region of South Africa. *Quat. Sci. Rev.* 67, 105–120.
- Stoffers, P., Hecky, R.E., 1978. Late Pleistocene–Holocene Evolution of the Kivu–Tanganyika Basin. In: Matter, A., Tucker, M.E. (Eds.), *Modern and Ancient Lake Sediments*. Blackwell Publishing Ltd., Oxford, UK, pp. 43–55.
- Stoffers, P., A. Singer, 1979. Clay minerals in Lake Mobutu Sese Seko (Lake Albert) - their diagenetic changes as an indicator of the paleoclimate. *Geologische Rundschau* 68: 1009-1024.
- Stone, A.E.C., Thomas, D.S.G., Viles, H.A. 2010. Late Quaternary palaeohydrological changes in the northern Namib Sand Sea: New chronologies using OSL dating of interdigitated aeolian and water-lain interdune deposits. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 288, 35-53.
- Stuiver, M., Deevey, E.S., Gjalenski, L.J., 1960. Yale natural radiocarbon measurements V 2, 49–61.
- Taieb, M., Barker, P., Bonnefille, R., Damnati, B., Gasse, F., Goetz, C., Hillaire-Marcel, C., Icole, M., Massault, M., Roberts, N., Vincens, A., Williamson, D., 1991. Histoire paleohydrologique du lac Magadi (Kenya) au Pleistocene superieur. *C.R. Acad. Sc. Paris*.
- Talbot, M.R., Livingstone, D.A., 1989. Hydrogen index and carbon isotopes of lacustrine organic matter as lake level indicators. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 70, 121–137.
- Telfer, M.W., Thomas, D.S.G., Parker, A.G., Walkington, H., Finch, A.A., 2009. Optically Stimulated Luminescence (OSL) dating and palaeoenvironmental studies of pan (playa) sediment from Witpan, South Africa. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 273, 50-60.
- Telford, R.J., Lamb, H.F., 1999. Groundwater-Mediated Response to Holocene Climatic Change Recorded by the Diatom Stratigraphy of an Ethiopian Crater Lake. *Quat. Res.* 52, 63–75.
- Teller, J.T., Lancaster, N. 1986 Lacustrine sediments at Narabeb in the central Namib Desert, Namibia. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 56, 177-195.
- Teller, J.T., Rutter, N., Lancaster, N. 1990. Sedimentology and paleohydrology of Late Quaternary lake deposits in the northern Namib Sand Sea, Namibia. *Quaternary Science Reviews*, 9, 343-364.
- Thevenon, F., Williamson, D., Taieb, M., 2002. A 22 kyr BP sedimentological record of Lake Rukwa (8°S, SW Tanzania): Environmental, chronostratigraphic and climatic implications. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 187, 285–294.

- Thomas, D.S.G., Brook, G., Shaw, P., Bateman, M., Appleton, C., Nash, D., McLaren, S., Davies, F., 2003. Late Pleistocene wetting and drying in the NW Kalahari: an integrated study from the Tsodilo Hills, Botswana. *Quat. Int.* 104, 53–67.
- Thomas, D.S.G., Bailey, R., Shaw, P.A., Durcan, J.A., Singarayer, J.S., 2009. Late Quaternary highstands at Lake Chilwa, Malawi: Frequency, timing and possible forcing
- Tiercelin, J.J., Renaut R.W., Delibrias G., LeFournier J., Bieda S., 1981. Late Pleistocene and Holocene lake level fluctuations in the Lake Bogoria basin, northern Kenya rift valley. *Palaeoecol. Afr.* 13, 105–120.
- Tiercelin, J.J., Vincens, A. (eds.), 1987. Le demi-graben de Baringo–Bogoria, Rift Gregory, Kenya: 30,000 ans d’histoire hydrologique et sedimentaire. *Bull. Centres Rech. Explor. Prod. Elf-Aquitaine*, 11, 249–540.
- Tierney, J.E., Russell, J.M., Huang, Y., 2010. A molecular perspective on Late Quaternary climate and vegetation change in the Lake Tanganyika basin, East Africa. *Quat. Sci. Rev.* 29, 787–800.
- Truckle, P.H., 1976. Geology and late Cainozoic lake sediments of the Suguta Trough, Kenya. *Nature* 263, 380–383.
- Van Bocxlaer, B., Salenbien, W., Praet, N., Verniers, J., 2012. Stratigraphy and paleoenvironments of the early to middle Holocene Chipalamawamba Beds (Malawi Basin, Africa). *Biogeosciences* 9, 4497–4512.
- Van der Meeren, T., Ito, E., Laird, K.R., Cumming, B.F., Verschuren, D., 2019. Ecohydrological evolution of Lake Naivasha (central Rift Valley, Kenya) during the past 1650 years, as recorded by ostracod assemblages and stable-isotope geochemistry. *Quat. Sci. Rev.* 223, 105906.
- Vareschi, E., 1982. The ecology of Lake Nakuru (Kenya) - III. Abiotic factors and primary production. *Oecologia* 55, 81–101.
- Velpuri, N.M., Senay, G.B., Asante, K.O., 2012. A multi-source satellite data approach for modelling Lake Turkana water level: Calibration and validation using satellite altimetry data. *Hydrol. Earth Syst. Sci.* 16, 1–18.
- Verschuren, D., 1999. Influence of depth and mixing regime on sedimentation in a small, fluctuating tropical soda lake. *Limnol. Oceanogr.* 44(4), 1103–1113.
- Verschuren, D., 2001. Reconstructing fluctuations of a shallow East African lake during the past 1800 yrs from sediment stratigraphy in a submerged crater basin. 297–311.

- Verschuren, D., Cocquyt, C., Tibby, J., Roberts, C.N., Leavitt, P.R., 1999. Long-term dynamics of algal and invertebrate communities in a small, fluctuating tropical soda lake. *Limnol. Oceanogr.* 44(5), 1216-1231.
- Verschuren, D., Laird, K. R., Cumming, B. F., 2000. Rainfall and drought in equatorial east Africa during the past 1,100 years. *Nature* 403, 410–414.
- Verschuren, D., Sinninghe Damsté, J.S., Moernaut, J., Kristen, I., Blaauw, M., Fagot, M., Haug, G.H., 2009. Half-precessional dynamics of monsoon rainfall near the East African Equator. *Nature* 462, 637–641.
- Vincens, A., 1986. Diagramme pollinique d'un sondage Pleistocene superieur-Holocene du Lac Bogoria (Kenya). *Rev. Palaeobot. Palynol.* 47, 169–192.
- Vincens, A., Casanova, J., Tiercelin, J.J., 1986. Palaeolimnology of Lake Bogoria (Kenya) during the 4500 BP high lacustrine phase. In: Frostick, L.E., Renaut, R.W., Reid I., Tiercelin, J.J. (eds.), *Sedimentation in the African Rifts*. Geol. Soc. London Spec. Publ., 25, 323–330.
- Vincens, A., Buchet, G., Williamson, D., Taieb, M., 2005. A 23,000 yr pollen record from Lake Rukwa (8°S, SW Tanzania): New data on vegetation dynamics and climate in Central Eastern Africa. *Rev. Palaeobot. Palynol.* 137, 147–162.
- Viner, A.B., 1977. The sediments of Lake George (Uganda), IV: Vertical distribution of chemical features in relation to ecological history and nutrient recycling. *Arch. Hydrobiol.* 80, 40–69.
- Vogel, J.C., Visser, E. 1981. Pretoria radiocarbon dates II. *Radiocarbon*, 23, 43-80.
- Washbourn-Kamau, C.K., 1971. Late Quaternary Lakes in the Nakuru-Elmenteita Basin , Kenya 137, 522–535.
- Washbourn-Kamau, C. K., 1975. Late Quaternary Shorelines of Lake Naivasha, Kenya. *Azania Archaeol. Res. Africa* 10, 77–92.
- Williams, M.A.J., Bishop, P.M., Dakin, F.M., Gillespie, R., 1977. Late Quaternary lake levels in southern Afar and the adjacent Ethiopian Rift. *Nature* 267, 690-693.
- Williams, M.A.J., Williams, F.M., Bishop, P.M., 1981. Late Quaternary history of Lake Besaka, Ethiopia. *Paleoecol. Afr.* 13, 93-104.
- Williams, R.E.G., Johnson, A.S., 1976. Birmingham University Radiocarbon Dates X. *Radiocarbon* 18, 249–267.

- Williamson, D., Jelinowska, A., Kissel, C., Tucholka, P., Gibert, E., Gasse, F., Massault, M., Taieb, M., Van Campo, E., and Wieckowski, K., 1998. Rock magnetic proxies of erosion/oxidation cycles in Late Quaternary maar lake sediments (Lake Tritrivakely Madagascar): paleoenvironmental implications. *Earth and Planetary Science Letters* 155, 205–219.
- Williamson, D., Jackson, M.J., Banerjee, S.K., Marvin, J., Merdaci, O., Thouveny, N., Decobert, M., Gibert-Massault, E., Massault, M., Mazaudier, D., Taieb, M., 1999. Magnetic signatures of hydrological change in a tropical maar-lake (Lake Massoko, Tanzania): Preliminary results. *Phys. Chem. Earth, Part A Solid Earth Geod.* 24, 799–803.
- Williamson, D., Taieb, M., Damnati, B., Icole, M., Thouveny, N., 1993. Equatorial extension of the younger Dryas event: rock magnetic evidence from Lake Magadi (Kenya). *Glob. Planet. Change* 7, 235–242.
- Wolff, C., Haug, G.H., Timmermann, A., Damste, J.S.S., Brauer, A., Sigman, D.M., Cane, M.A., Verschuren, D., 2011. Reduced Interannual Rainfall Variability in East Africa During the Last Ice Age. *Science* 333, 743–747.
- Wüest, A., Piepke, G., Halfman, J., 1996. Combined effects of dissolved solids and temperature on the density stratification of Lake Malawi. In: Johnson, T., Odada, E. (Eds.), *The Limnology, Climatology and Paleoclimatology of the East African Lakes*. Gordon and Breach, Amsterdam, pp. 183–202.
- Young, J.A.T. and Renaut, R.W., 1979. A radiocarbon date from Lake Bogoria, Kenya Rift Valley. *Nature* 278, 243–245.