

Vitamin D and covid-19

Article

Accepted Version

Vimaleswaran, K. S. ORCID: <https://orcid.org/0000-0002-8485-8930>, Forouhi, N. G. and Khunti, K. (2021) Vitamin D and covid-19. British Medical Journal, 372. n544. ISSN 1468-5833 doi: 10.1136/bmj.n544 Available at <https://centaur.reading.ac.uk/97086/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1136/bmj.n544>

Publisher: BMJ Publishing Group Ltd

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Can current guidelines on vitamin D supplementation prevent or treat SARS-CoV-2 infection?

Karani S Vimalleswaran¹, Nita G Forouhi², Kamlesh Khunti³

¹ Hugh Sinclair Unit of Human Nutrition, Department of Food and Nutritional Sciences, University of Reading, Reading, UK. Email: v.karani@reading.ac.uk

² MRC Epidemiology Unit, University of Cambridge School of Clinical Medicine, Cambridge, UK. Email: nita.forouhi@mrc-epid.cam.ac.uk

³ Primary Care Diabetes & Vascular Medicine, Diabetes Research Centre and The Centre for Black Minority Health, University of Leicester, Trustee, South Asian Health Foundation, Leicester, UK. Email: kk22@leicester.ac.uk

No. of words: 800 (Text)

No. of references: 16

Corresponding author:

Karani S Vimalleswaran, Professor in Nutrigenetics and Nutrigenomics, Hugh Sinclair Unit of Human Nutrition, Department of Food and Nutritional Sciences, University of Reading, Reading, UK. Email: v.karani@reading.ac.uk

SARS-CoV-2 infection-related health crisis has led to unfounded or exaggerated claims on treatments. One area of controversy has been the role and dose of vitamin D supplementation in COVID-19^{1 2}. Given the potential importance but prevailing uncertainty, the joint National Institute for Health and Care Excellence (NICE), Public Health England and Scientific Advisory Committee on Nutrition rapid guideline was recently published. It concluded that there was little evidence, highlighted the need for further research, and supported the existing government advice predicated on musculoskeletal health on vitamin D supplementation of 400IU/day for adults and children between October and March, when people in the UK do not make sufficient vitamin D from sunlight. It also stressed that certain populations such as minority ethnic groups, should consider taking this dose throughout the year³. The guidance is timely, but questions remain.

What is the evidence for a link between vitamin D and COVID-19?

Vitamin D supplementation of 400-1000IU/day has a modest protective effect for acute respiratory infections⁴, providing indirect evidence for SARS-CoV-2 infection. There is, however, sparse research on a direct link.

The NICE review included one small randomised controlled trial (RCT) for COVID-19 treatment⁵, no RCTs for COVID-19 prevention, and 12 observational studies of associations between vitamin D status and COVID-19 incidence or treatment. Among 76 Spanish patients hospitalised with COVID-19, high-dose supplementation equivalent to 21,280IU cholecalciferol on admission day and 10,640IU on day 3 and 7, then weekly until discharge reduced disease severity⁵. We found two further RCTs with conflicting findings. Among 240 Brazilian hospitalised patients with COVID-19, a single oral vitamin D₃ dose of 200,000IU versus placebo did not reduce the hospital length of stay⁶. In contrast, in 40 Indian SARS-CoV-2 positive Indians, infection vitamin D 60,000IU daily for 7 days, followed by the same dose either weekly or daily led to greater negative tests at 21 days⁷. These discrepancies highlight

the challenges of RCT design, with varying selection criteria, initial vitamin D status, the type, dose, and duration of supplementation, the endpoints studied, and risk of bias and study quality.

The observational evidence is inconsistent with some, but not all, studies reporting an association between vitamin D insufficiency and SARS-CoV-2 infection^{3 8}. However, the observed link could be attributed to several confounding factors including age⁹, ethnicity¹⁰, genetic heterogeneity¹¹, and obesity^{12 13} which are incompletely or not accounted for in different studies.

High-quality evidence is currently lacking, but there is suggestive evidence for an immunomodulatory role of vitamin D for respiratory infections and contextual evidence of the shared risk factors between vitamin D deficiency and COVID-19 severity: older age, obesity, and minority ethnicity. There is also a correlation between seasonal decline of vitamin D and higher COVID-19 burden in high-latitude countries¹⁴. The available evidence cannot be ignored and makes a compelling case for further research.

What next with the UK guidance on use of vitamin D supplementation?

The recommendation of 400IU seems justifiable to maintain 25(OH)D levels >25nmol/l in 97.5% of the UK population, but it is unclear whether this level is appropriate for immunomodulatory actions for COVID-19. Guidance recommending 400IU/day vitamin D supplementation in the UK has been in existence for a while, but its implementation has not been ensured. Raising awareness of the relevance of vitamin D is therefore appropriate for musculoskeletal health, particularly during lockdowns. It may also be relevant for COVID-19, given the suggestive though not conclusive evidence of its potential role and the precautionary principle. We also need clear guidance on how to obtain vitamin D for vulnerable groups. Health care professionals can point people to the free NHS vitamin D supplement provisions for people at high risk¹⁵ and for women and children who qualify for the Healthy Start scheme.

Clinicians also need to be aware that vegetarians or vegans would need guidance on appropriate sources of vitamin D supplements.

The public health emergency posed by COVID-19 demands the use of all promising solutions, therefore vitamin D remains a plausible candidate. However, policy recommendations need to ensure that the public are not falsely reassured regarding the role of vitamin D for COVID-19. These guidelines therefore must be accompanied with continued messaging on hand hygiene, face coverings, physical distancing, and the importance of vaccine uptake in culturally and linguistically adapted campaigns through local community groups.

The published guidelines have clearly articulated the currently unconvincing evidence of vitamin D for COVID-19; therefore, it is vital that ongoing and future trials evaluate the effect of vitamin D supplementation with improved design including attention to comparing different dosing regimens, initial vitamin D status, inclusion of different population subgroups, older participants and those with morbidities, and in settings including hospitalised patients and population-based samples. Ongoing trials such as COVIT-TRIAL¹⁶ and CORONAVIT (NCT04579640), which is comparing the national recommendation with higher dosage (800IU/day and 3200IU/day), will be important to inform future guidance.

Acknowledgements

NGF acknowledges funding from MRC Epidemiology Unit (MC_UU_12015/5) and NIHR Biomedical Research Centre Cambridge: Nutrition, Diet, and Lifestyle Research Theme (IS-BRC-1215-20014). KK is supported by the National Institute for Health Research (NIHR) Applied Research Collaboration East Midlands (ARC EM) and the NIHR Leicester Biomedical Research Centre (BRC).

Duality of Interest

NGF is an honorary consultant public health physician with Public Health England. The views expressed are her own. KK is Director of the University of Leicester Centre for Black Minority Ethnic Health, Trustee of the South Asian Health Foundation, Chair of the Ethnicity Subgroup of SAGE and Member of Independent SAGE.

References

1. Ali N. Role of vitamin D in preventing of COVID-19 infection, progression and severity. *J Infect Public Health* 2020;13(10):1373-80. doi: 10.1016/j.jiph.2020.06.021 [published Online First: 2020/07/02]
2. Pereira M, Dantas Damascena A, Galvao Azevedo LM, et al. Vitamin D deficiency aggravates COVID-19: systematic review and meta-analysis. *Crit Rev Food Sci Nutr* 2020;1-9. doi: 10.1080/10408398.2020.1841090 [published Online First: 2020/11/05]
3. NICE. Vitamin D for COVID-19: Evidence reviews for the use of vitamin D supplementation as prevention and treatment of COVID-19 2020 [Available from: <https://www.nice.org.uk/guidance/ng187/evidence/evidence-reviews-for-the-use-of-vitamin-d-supplementation-as-prevention-and-treatment-of-covid19-pdf-8957587789>].
4. Jolliffe DA, Camargo CA, Sluyter JD, et al. Vitamin D supplementation to prevent acute respiratory infections: systematic review and meta-analysis of aggregate data from randomised controlled trials. *medRxiv* 2020 doi: 10.1101/2020.07.14.20152728 [published Online First: 2020/12/04]
5. Entrenas Castillo M, Entrenas Costa LM, Vaquero Barrios JM, et al. "Effect of calcifediol treatment and best available therapy versus best available therapy on intensive care unit admission and mortality among patients hospitalized for COVID-19: A pilot randomized clinical study". *J Steroid Biochem Mol Biol* 2020;203:105751. doi: 10.1016/j.jsbmb.2020.105751 [published Online First: 2020/09/02]
6. Murai IH, Fernandes AL, Sales LP, et al. Effect of a Single High Dose of Vitamin D3 on Hospital Length of Stay in Patients With Moderate to Severe COVID-19: A Randomized Clinical Trial. *JAMA* 2021 doi: 10.1001/jama.2020.26848 [published Online First: 2021/02/18]

7. Rastogi A, Bhansali A, Khare N, et al. Short term, high-dose vitamin D supplementation for COVID-19 disease: a randomised, placebo-controlled, study (SHADE study). *Postgrad Med J* 2020 doi: 10.1136/postgradmedj-2020-139065 [published Online First: 2020/11/14]
8. Kaufman HW, Niles JK, Kroll MH, et al. SARS-CoV-2 positivity rates associated with circulating 25-hydroxyvitamin D levels. *PLoS One* 2020;15(9):e0239252. doi: 10.1371/journal.pone.0239252 [published Online First: 2020/09/18]
9. Baktash V, Hosack T, Patel N, et al. Vitamin D status and outcomes for hospitalised older patients with COVID-19. *Postgrad Med J* 2020 doi: 10.1136/postgradmedj-2020-138712 [published Online First: 2020/08/29]
10. Khunti K, Singh AK, Pareek M, et al. Is ethnicity linked to incidence or outcomes of covid-19? *BMJ* 2020;369:m1548. doi: 10.1136/bmj.m1548 [published Online First: 2020/04/22]
11. Anastassopoulou C, Gkizarioti Z, Patrinos GP, et al. Human genetic factors associated with susceptibility to SARS-CoV-2 infection and COVID-19 disease severity. *Hum Genomics* 2020;14(1):40. doi: 10.1186/s40246-020-00290-4 [published Online First: 2020/10/24]
12. Syed AA, Soran H, Adam S. Obesity and covid-19: the unseen risks. *BMJ* 2020;370:m2823. doi: 10.1136/bmj.m2823 [published Online First: 2020/07/18]
13. Vimalaswaran KS, Berry DJ, Lu C, et al. Causal relationship between obesity and vitamin D status: bi-directional Mendelian randomization analysis of multiple cohorts. *PLoS Med* 2013;10(2):e1001383. doi: 10.1371/journal.pmed.1001383 [published Online First: 2013/02/09]

- 144 14. Kohlmeier M. Avoidance of vitamin D deficiency to slow the COVID-19 pandemic. *BMJ*
145 *Nutr Prev Health* 2020;3(1):67-73. doi: 10.1136/bmjnp-2020-000096 [published
146 Online First: 2020/11/25]
- 147 15. NHS. Get vitamin D supplements 2020 [Available from:
148 [https://www.nhs.uk/conditions/coronavirus-covid-19/people-at-higher-risk/get-](https://www.nhs.uk/conditions/coronavirus-covid-19/people-at-higher-risk/get-vitamin-d-supplements/)
149 [vitamin-d-supplements/](https://www.nhs.uk/conditions/coronavirus-covid-19/people-at-higher-risk/get-vitamin-d-supplements/).
- 150 16. Annweiler C, Beaudenon M, Gautier J, et al. COvid-19 and high-dose VITamin D
151 supplementation TRIAL in high-risk older patients (COVIT-TRIAL): study protocol
152 for a randomized controlled trial. *Trials* 2020;21(1):1031. doi: 10.1186/s13063-020-
153 04928-5 [published Online First: 2020/12/30]