

# *Using ecological and field survey data to establish a national list of the wild bee pollinators of crops*

Article

Accepted Version

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Hutchinson, L. A., Oliver, T. H. ORCID: <https://orcid.org/0000-0002-4169-7313>, Breeze, T. D. ORCID: <https://orcid.org/0000-0002-8929-8354>, Bailes, E. J., Brünjes, L. ORCID: <https://orcid.org/0000-0002-1344-1476>, Campbell, A. J., Erhardt, A., de Groot, G. A., Földesi, R., García, D., Goulson, D., Hainaut, H., Hambäck, P. A., Holzschuh, A., Jauker, F., Klatt, B. K., Klein, A.-M., Kleijn, D. ORCID: <https://orcid.org/0000-0003-2500-7164>, Kovács-Hostyánszki, A. ORCID: <https://orcid.org/0000-0001-5906-4816>, Krimmer, E., McKerchar, M., Miñarro, M., Phillips, B. B., Potts, S. G. ORCID: <https://orcid.org/0000-0002-2045-980X>, Pufal, G., Radzevičiūtė, R., Roberts, S. P. M., Samnegård, U., Schulze, J. ORCID: <https://orcid.org/0000-0002-6944-0130>, Shaw, R. F., Tscharntke, T., Vereecken, N. J., Westbury, D. B., Westphal, C. ORCID: <https://orcid.org/0000-0002-2615-1339>, Wietzke, A., Woodcock, B. A. and Garratt, M. P. D. ORCID: <https://orcid.org/0000-0002-0196-6013> (2021) Using ecological and field survey data to establish a national list of the wild bee pollinators of crops. *Agriculture, Ecosystems & Environment*, 315. 107447. ISSN 0167-8809 doi: 10.1016/j.agee.2021.107447 Available at

<https://centaur.reading.ac.uk/97939/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1016/j.agee.2021.107447>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

## **CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

**Using ecological and field survey data to establish a national list of the wild bee  
pollinators of crops.**

Louise A. Hutchinson<sup>1\*</sup>, Tom H. Oliver<sup>2</sup>, Tom D. Breeze<sup>1</sup>, Emily J. Bailes<sup>3,4,5</sup>, Lisa Brünjes<sup>6</sup>, Alistair J. Campbell<sup>7</sup>, Andreas Erhardt<sup>8</sup>, G. Arjen de Groot<sup>9</sup>, Rita Földesi<sup>10</sup>, Daniel García<sup>11</sup>, Dave Goulson<sup>12</sup>, Hélène Hainaut<sup>13</sup>, Peter A. Hambäck<sup>14</sup>, Andrea Holzschuh<sup>15</sup>, Frank Jauker<sup>16</sup>, Björn K. Klatt<sup>17,18</sup>, Alexandra-Maria Klein<sup>19</sup>, David Kleijn<sup>20</sup>, Anikó Kovács-Hostyánszki<sup>21</sup>, Elena Krimmer<sup>15</sup>, Megan McKerchar<sup>22</sup>, Marcos Miñarro<sup>23</sup>, Benjamin B. Phillips<sup>24</sup>, Simon G. Potts<sup>1</sup>, Gesine Pufal<sup>19</sup>, Rita Radzevičiūtė<sup>25,26,27</sup>, Stuart P. M. Roberts<sup>1</sup>, Ulrika Samnegård<sup>13,17</sup>, Jürg Schulze<sup>28</sup>, Rosalind F. Shaw<sup>24</sup>, Teja Tscharnke<sup>29</sup>, Nicolas J. Vereecken<sup>13</sup>, Duncan B. Westbury<sup>22</sup>, Catrin Westphal<sup>18</sup>, Alexander Wietzke<sup>30</sup>, Ben A. Woodcock<sup>31</sup>, Michael P. D. Garratt<sup>1</sup>

<sup>1</sup>Centre for Agri-Environmental Research, School of Agriculture, Policy and Development, University of Reading, United Kingdom

<sup>2</sup>School of Biological Sciences, University of Reading, United Kingdom

<sup>3</sup>Department of Molecular Biology and Biotechnology, University of Sheffield, United Kingdom

<sup>4</sup>Department of Plant Sciences, University of Cambridge, Cambridge, United Kingdom

<sup>5</sup>National Institute of Agricultural Botany, Cambridge, United Kingdom

<sup>6</sup>Plant Breeding Methodology, Department of Crop Sciences, University of Göttingen, Göttingen, Germany

<sup>7</sup>Embrapa Amazônia Oriental, Travessa Enéas Pinheiro, Marco, Belém, CEP 66095-903, Pará, Brazil

<sup>8</sup>University of Basel, Department of Environmental Sciences, Botany, Schönbeinstrasse 6, CH-4056, Basel, Switzerland

<sup>9</sup>Wageningen Environmental Research, Wageningen UR, P.O. Box 47, 6700 AA Wageningen, the Netherlands

<sup>10</sup>Lendület Ecosystem Services Research Group, Institute of Ecology and Botany, Centre for Ecological Research, 2163 Vácrátót, Hungary

<sup>11</sup>Depto. Biología de Organismos y Sistemas, Universidad de Oviedo, and Unidad Mixta de Investigación en Biodiversidad (CSIC-Uo-PA). C/Catedrático Rodrigo Uría s/n, E-33006 Oviedo, Asturias, Spain

<sup>12</sup>School of Life Sciences, University of Sussex, Brighton, UK

- <sup>13</sup>Agroecology Lab, Université Libre de Bruxelles (ULB), Boulevard du Triomphe CP 264/2, B-1050 Brussels, Belgium
- <sup>14</sup>Department of Ecology, Environment and Plant Sciences, Stockholm University, 106 91 Stockholm, Sweden
- <sup>15</sup>Animal Ecology and Tropical Biology, Biocenter, University of Würzburg, 97074 Würzburg, Germany
- <sup>16</sup>Department of Animal Ecology, Justus Liebig University Giessen, Heinrich-Buff-Ring 26-32, D-35392 Giessen, Germany
- <sup>17</sup>Department of Biology, Lund University, SE-223 62 Lund, Sweden
- <sup>18</sup>Functional Agrobiodiversity, Department of Crop Sciences, University of Göttingen, Göttingen, Germany
- <sup>19</sup>Nature Conservation and Landscape Ecology, Faculty of Environment and Natural Resources, University of Freiburg, Freiburg, Germany
- <sup>20</sup>Plant Ecology and Nature Conservation Group, Wageningen University, Droevendaalsesteeg 3a, 6708PB, Wageningen, The Netherlands
- <sup>21</sup>Lendület Ecosystem Services Research Group, Institute of Ecology and Botany, Centre for Ecological Research, Alkotmány str. 2-4, 2163 Vácrátót, Hungary
- <sup>22</sup>School of Science & the Environment, University of Worcester, Worcester, United Kingdom
- <sup>23</sup>Servicio Regional de Investigación y Desarrollo Agroalimentario (SERIDA). Apdo. 13, E-33300 Villaviciosa, Asturias, Spain
- <sup>24</sup>Environment and Sustainability Institute, University of Exeter, Penryn Campus, Penryn, Cornwall TR10 9FE, United Kingdom
- <sup>25</sup>General Zoology, Institute for Biology, Martin Luther University Halle-Wittenberg, Hoher Weg 8, D-06120 Halle (Saale), Germany
- <sup>26</sup>Molecular Evolution and Animal Systematics, Institute for Biology, Leipzig University, Talstraße 33, D-04103 Leipzig, Germany
- <sup>27</sup>Life Sciences Center, Vilnius University, Saulėtekio al. 7, LT-10223 Vilnius, Lithuania
- <sup>28</sup>Agency for Environment and Energy Canton Basel-City, Hochbergerstr. 157, 4019 Basel, Switzerland
- <sup>29</sup>Agroecology, Dept. of Crop Sciences, University of Göttingen, Grisebachstrasse 6, 37077 Göttingen, Germany

<sup>30</sup>Plant Ecology and Ecosystems Research, University of Goettingen, Untere Karspüle 2, 37073 Göttingen, Germany

<sup>31</sup>UK Centre for Ecology & Hydrology, Crowmarsh Gifford, Wallingford, Oxfordshire, United Kingdom

\*Corresponding author. E-mail: l.hutchinson@pgr.reading.ac.uk

## **Abstract**

The importance of wild bees for crop pollination is well established, but less is known about which species contribute to service delivery to inform agricultural management, monitoring and conservation. Using sites in Great Britain as a case study, we use a novel qualitative approach combining ecological information and field survey data to establish a national list of crop pollinating bees for four economically important crops (apple, field bean, oilseed rape and strawberry). A traits data base was used to establish potential pollinators, and combined with field data to identify both dominant crop flower visiting bee species and other species that could be important crop pollinators, but which are not presently sampled in large numbers on crops flowers. Whilst we found evidence that a small number of common, generalist species make a disproportionate contribution to flower visits, many more species were identified as potential pollinators, including rare and specialist species. Furthermore, we found evidence of substantial variation in the bee communities of different crops. Establishing a national list of crop pollinators is important for practitioners and policy makers, allowing targeted management approaches for improved ecosystem services, conservation and species monitoring. Data can be used to make recommendations about how pollinator diversity could be promoted in agricultural landscapes. Our results suggest agri-environment schemes need to support a higher diversity of species than at present, notably of solitary bees. Management would also benefit from targeting specific species to enhance crop pollination services to particular crops. Whilst our study is focused upon Great Britain, our methodology can easily be applied to other countries, crops and groups of pollinating insects.

## **Key-words**

Agri-environment Schemes, Apple, Biodiversity, Crop pollination, Dominant Pollinators, Ecosystem Services, Field Bean, Oilseed Rape, Rare Species, Strawberry.

## 1. Introduction

Insect pollination is key to global agricultural productivity (IPBES, 2016) due to growing demand for entomophilous crops (Godfray and Garnett 2014). The nutritional and economic importance of insect pollinated crops (Vanbergen et al., 2014), and the inability of managed pollinators (e.g., *Apis mellifera*) to meet service demand, mean agriculture is highly dependent upon wild pollinators (Aizen and Harder 2009; Breeze et al., 2014). Yet conventional agricultural practices are a key driver of pollinator declines (Senapathi et al., 2015). Whilst agri-environment scheme options have had positive impacts (Tonietto et al., 2018), most benefit a limited suite of common species (Scheper et al., 2013) and homogeneous communities provide less reliable pollination services (Grab et al., 2019). Currently agri-environment schemes tend preferentially to benefit bumblebee populations (Wood et al., 2015a; Wood et al., 2015b, 2016a, b), yet solitary bee species are more important pollinators of some crops (Woodcock et al., 2013). As such, current agri-environment schemes may not be optimally designed to increase pollination services to many crops. Identifying key pollinating species to individual crops, and ones which may provide additional pollination and insurance against declines in other species, would help inform agricultural management for bee pollinators (Garratt et al., 2014a). Yet there is insufficient information on bee communities for many crops (Kremen and Chaplin-Kramer, 2007) and no studies have attempted to establish a 'national list' of crop pollinators to advise management or monitoring programmes.

Whilst the majority of crop flower visitation is attributed to a small proportion of bee species (Kleijn et al., 2015), species-rich communities have been shown to positively influence crop yields and pollination service stability (Hoehn et al., 2008; Garibaldi et al., 2011; Martins et al., 2015; Dainese et al., 2019; Woodcock et al., 2019). Biodiversity conservation and ecosystem service management are often seen as distinct objectives (Sutter et al., 2017), however management that only targets common crop pollinators will not safeguard production if it fails to encompass species that supplement service provision (Fijen et al., 2018). High species turnover means that diverse communities, including rare and specialist species, are required

to maintain crop pollination service at regional scales (Winfree et al., 2018). With climate change reducing the occupancy and richness of some wild bee species (Soroye et al., 2020), supporting wider species diversity may be crucial for crop pollination service stability under the substantial future environmental change that is predicted (Oliver et al., 2015; Dainese et al., 2019). Additionally, different crops have distinct pollinator communities and it will be beneficial to identify the pollinating taxa of individual crops and target management accordingly (Garratt et al., 2014a). Furthermore, a national list of crop pollinators can inform monitoring schemes to ensure they include important crop pollinating species (Carvell et al., 2017; Garratt et al., 2019).

In order to inform pollinator management and monitoring, our study aimed to compile the bee species visiting four crops: apple (*Malus domestica*), field bean (*Vicia faba*), oilseed rape (*Brassica napus*) and strawberry (*Fragaria x ananassa*). Insect pollination has been shown to enhance yield quantity and quality in all four crops (Bartomeus et al., 2014; Garratt et al., 2014b). Additionally, they differ in flower phenology and morphology (Garibaldi et al., 2015) and likely show corresponding differences in their pollinator community composition (Garratt et al., 2014a). We use sites in Great Britain as a case study because its bee fauna is comprehensively described and their occupancy is well recorded over a long time period (Powney et al. 2019). We compiled a list of all British bee species and their available physiological and ecological traits, and combined it with field survey data in order to devise an approach to generate lists of (i) definite flower visitors to each crop (ii) likely flower visitors, which are expected to also contribute to crop pollination (iii) possible crop flower visitors whose contribution to pollination is not well understood and merits further investigation. Our aim was to compile these lists for reference purposes, but not to statistically compare pollinator communities between crops, due to the unstandardised nature of the datasets used to generate the lists of bee species. Additionally, we identify dominant crop pollinating species, and assess the contribution of wild bees compared to honey bees for crop flower visitation.

## **2. Materials and Methods**

### **2.1 Potential crop pollinators.**

First, a species database of all extant, resident wild bee species in Great Britain was established using the most recent checklist of UK species (Else et al., 2016). For each species, data on the following were collated: flight period (months); sociality (cleptoparasite, eusocial or solitary); lecty (oligolectic or polylectic, including if any of the target crop plant families are visited for pollen and/or nectar), tongue length (short/long), geographic coverage (distribution and habitat) (based on trait information compiled by Stuart Roberts for the EU- FP6 ALARM-project and BWARS, 2020) and conservation status (Webb et al., 2018). Potential crop pollinators, as defined here, are those bee species which, based upon these ecological traits, such as flight period, lecty, sociality and tongue length, could pollinate our target crops. Habitat specialists that are not coincident with cropland were initially excluded i.e., primarily coastal, heathland species. The known floral ecology of each species was then used to refine lists for each crop. Cleptoparasitic species, species that are oligolectic on plant families other than the target crop or polylectic, but not documented as foraging on the relevant plant family for pollen or nectar and species whose flight period does not overlap with the relevant crops flowering period were excluded. For field bean, only 'long-tongued' species (Michener, 2000) were considered as its flowers have deep corollas and most visits by 'short-tongued' species involve nectar robbing rather than legitimate visitation (Garratt et al., 2014a).

### **2.2 Field survey data**

Field studies were sourced through literature searches in google scholar and existing datasets held by the authors. Fifty-seven datasets from across England, Scotland and eight other European countries were available to combine with the potential crop pollinator lists in order to establish shortlists of crop flower visitors (Figure 1 and Table S3).





Lists of bee species recorded in crop fields were compiled using three types of survey data:

- i) British flower visitation studies (e.g. transect walks, observation plots).
- ii) British pan trap studies in crop fields.
- iii) Other European flower visitation studies (used to validate crop flower visitation for species sampled in British pan traps only).

For every bee species the total number of reported legitimate flower visits and number of studies recorded in were calculated for each crop. If studies did not include quantitative data then a conservative approach was taken whereby each bee species listed was taken as representing a single crop flower visit. As pan trap catches do not provide information on floral associations (Westphal et al., 2008), these data were only used, in combination with trait data, to generate the list of possible pollinators.

### **2.3 Crop flower visitors**

The lists of potential crop pollinators were combined with the field survey data to categorize bee species into one of three flower visitor categories (Figure 2; Full details in Supplementary Methods 1):

- i) **Definite Flower Visitors** – Species recorded visiting crop flowers in British flower visitation studies.
- ii) **Likely Flower Visitors** - Species recorded in British pan trap crop studies and recorded as making at least two flower visits in other European studies.
- iii) **Possible Flower Visitors** - Species only recorded in British pan trap studies, or in other European flower visitor studies only, and classified as a potential crop flower visitor.




Figure 2: Methodology by which bee species were categorised as definite, likely and possible flower visitors.

## 2.4 Dominant crop flowers visitors

As visitation rate to crop flowers is a good proxy of relative contribution to pollination service delivery (Vazquez et al., 2005), we identified the dominant British flower visiting bee species per crop by approximating the species attributed with a combined total of 80% of flower visits, the proportion identified as corresponding to the dominant flower visitors by Kleijn et al. (2015). Only British flower visitation datasets where bee species were either all identified to species or genus were included in the analysis (Supplementary Methods 2). Additionally, we calculated the average proportion of visits to crop flowers attributed to wild bees compared to honey bees for all crops (Supplementary Methods 2).

### 3. Results

#### 3.1 Potential crop pollinators

A preliminary list of 229 extant, resident British wild bee species was compiled. Of those 132 species were excluded due to ecological and lecty traits that were deemed incompatible with these bees being present in crop fields and/or crop flower visitors (Table S1). Four species were treated as an aggregate – *Bombus terrestris* aggregate – due to the difficulties of separating their workers in the field (Wolf et al., 2010; Bossert, 2015). Therefore, a total of 97 species were initially identified as potential crop pollinators. Accounting for their documented foraging ecology and flight period, the following number of species were considered as potential pollinators per crop: apple- 83, bean- 30, oilseed- 60, and strawberry – 90 (Table S2).

#### 3.2 Field survey data

The total number of studies sourced per crop were as follows: apple – 17; bean – 10; oilseed – 19; strawberry – 11. The number of studies per survey type for each crop is provided in Figure S1.

#### 3.3 Crop flower visitors

Seventy-three species from ten genera were categorised as flower visitors of one or more crops, 63 of which were recorded in British crop field studies (Table 1, Figure 3). Fourteen species were included in flower visitor categories that were not initially identified as potential crop pollinators. Ten of those were widely polylectic *Bombus* or *Lasioglossum* species, all recorded in oilseed datasets, but not documented in the literature as foraging on *Brassicaceae*. The remaining species were three short-tongued *Andrena* species recorded visiting bean flowers, two of which are oligolectic on Fabaceae and a *Colletes* species, recorded in a single strawberry dataset, that is documented as being oligolectic on another plant family. The majority of species identified as potential pollinators, but not recorded in crop field surveys were either rare species or polylectic species documented as having distinct

preferences for plant families other than the target crop. The remaining species were overwhelmingly smaller species from the genera *Hylaeus* and *Lasioglossum* or cavity nesting *Megachilidae*. Most species identified as crop flower visitors were geographically widespread (BWARS, 2020) and polylectic species. However, a quarter (n=18) of species included in flower visitor categories, currently have a designated conservation status in Britain. Full details of all species in crop flower visitor categories are given in tables S4a-d and S5a – S8d.

Table 1: Number of bee species, based upon field datasets and trait information, that were assigned to each category of flower visitor per crop

Crop	Flower Visitor Category			Total
	Definite	Likely	Possible	
Apple	19	13	25	57
Field Bean	11	0	3	14
Oilseed Rape	37	11	3	51
Strawberry	9	6	18	33

#### *Apple*

All five British apple flower visitor studies recorded every bee to species level. *Andrena* were the most speciose genus of flower visitor, both overall (n=22) and in the definite flower visitor category (n=10). *Bombus* species were the next most commonly represented genus in the latter category (n=6), but were less frequent overall (n=9) than *Lasioglossum* species (n=16). Within the definite flower visitor category 80% of flower visits were attributed to eight species, only half of which were recorded in all studies. Most likely and possible flower visitors were *Andrena* or *Lasioglossum* species.

#### *Bean*

Three of the five British bean flower visitor studies recorded all bee to species level, the remainder only recorded *Bombus* to species, which was both the most common genus overall (n=9) and in the definite flower visitor category (n=7). Three short-tongued *Andrena* sp. were identified as definite flower visitors, but all were recorded as very low numbers of flower visits


( $\leq 10$ ). Four *Bombus* species and *Anthophora plumipes* accounted for 95% of all visits recorded in British flower visitation studies. However, all the *A. plumipes* records derived from one study (Bond and Kirby, 1999) carried out at a single site. The four *Bombus* were the only species recorded in four or more studies. No species met the criteria for the likely flower visitor category. The possible flower visitor category included two *Bombus* and one *Osmia* species.

#### *Oilseed*

Six of the nine British oilseed flower visitor studies recorded bees to species level, but only two included quantitative data on all bee species. *Andrena* was the most speciose genus of bee, both overall ( $n=27$ ) and within the definite flower visitor category ( $n=15$ ). *Bombus* and *Lasioglossum* species were equally represented in the definite flower visitor category ( $n=9$ ), but *Lasioglossum* were more frequent overall ( $n=14$ ). Within the definite flower visitor category 80% of recorded flower visits were attributed to six species, only two of which were recorded in all nine studies, with the remainder only recorded in between five and eight studies, despite all being large *Andrena* or *Bombus* species, generally identified and quantified in all field studies. The likely and possible visitor categories were entirely comprised of *Andrena* or *Halictidae* species, two of which are oligolectic on *Brassicaceae*.

#### *Strawberry*

Two British strawberry flower visitor studies recorded all bees to species level. The remaining three only recorded a group of large *Andrena* and *Bombus* to species. *Bombus* species were the most common genus of bee within the definite flower visitor category ( $n=5$ ), but joint second as the most frequent genus overall, alongside *Lasioglossum* ( $n=7$ ), with *Andrena* species being the most prevalent genus across all categories ( $n=14$ ). Within the definite flower visitor category 80% of recorded flower visits were attributed to just two *Bombus* species, which along with two other *Bombus*, were the only species recorded in more than two studies. The likely visitor category was almost exclusively represented by *Andrena* species. The possible visitor category was largely comprised of solitary bees from five different genera.



119

120 Figure 3: The number of bee species from each genus which were categorised as definite  likely  or possible  flower visitors per crop

### 3.4 Dominant crop flower visitors

Ten bee species were attributed with 80% of flower visits across the four crops (Figure S2; Figure 4). There were differences however in the number and composition of those species making up the 80% of flower visits on a per crop basis. Differences in crop communities were even more distinct when considering the entire suite of bee species included in the characterisation of each crops' total flower visiting community (Figure 3; Figure 4). Wild bees were attributed with an average of between 63 and 83 percent of crop flower visits compared to honey bees (Apple: solitary bee visits = 68%; Bean: solitary bee visits = 83%; Oilseed: solitary bee visits = 63%; Strawberry: solitary bee = 77%).






Figure 4: Dominant crop visiting bee species (attributed with ~80% of flower visits in field studies per crop) shown as photographs, with number of bee species in each genus that are 'definite' flower visitors for each crop.

## 4. Discussion

### 4.1 Crop pollinator species

Our study is the one of the first to evaluate the entire wild bee community of multiple crops on a national basis and can be used as model approach for other countries, crops and pollinators. With the identification of bee species important for pollinating crops we build the basis to better sustainably manage services with changing climate and land use. Whilst in accordance with other studies (Rader et al., 2012; Kleijn et al., 2015) our results indicate that a small proportion of common, generalist bee species do make the majority of crop flower visits, many more species were evidenced as crop flower visitors. Additionally, our results suggest that the contribution of wild bee species to crop flower visitation may be even greater than previously thought. Whereas previous estimates indicate that wild bees make a similar overall contribution to honey bees (Kleijn et al. 2015), when considering the entire suite of flower visiting species our results indicate that wild bees make on average between 63 and 83% of flower visits to our target crops. Given the benefits of biodiverse communities for current and future crop pollination services (Kremen et al., 2002; Hoehn et al., 2008; Garibaldi et al., 2011; Rader et al., 2012), interventions to support crop pollinators should target a more significant proportion of the bee fauna than at present (Wood et al., 2015b, 2016a; Gresty et al., 2018). Establishing a list of currently important, but also potentially relevant crop pollinators, is necessary to help target monitoring and conservation (Carvell et al., 2017).

Our results also support prior evidence of distinct differences in individual crop pollinator communities (Garratt et al., 2014a). The overwhelming majority of field bean and strawberry flower visits were attributed to bumblebees. However, whereas field bean was visited by the three longest tongued species in Britain, strawberry crops were almost exclusively visited by two other bumblebee species, with relatively shorter tongues. This supports a link between trait matching of bees and flowers in crop pollination (Garibaldi et al., 2015). *Bombus* species were also recorded visiting apple and oilseed rape. However, due to their low abundance in early spring during apple flowering (Martins et al., 2015), and lower rate of pollen transfer

when visiting oilseed flowers (Woodcock et al., 2013) they are less important pollinators of these crops compared to solitary species. *Andrena* and *Lasioglossum* species were prevalent across both apple and oilseed flower visitor categories. *Andrena* are known to be highly efficient pollinators of both crops (Martins et al., 2015; Woodcock et al., 2013), especially apple (Russo et al., 2017). Most *Lasioglossum* species however, generally emerge later than many *Andrena* species, and peak after apple flowering, whereas oilseed tends to flower later and longer, and *Lasioglossum* are likely important pollinators of this crop (Perrot et al., 2018; Catarino et al., 2019). Furthermore, we almost certainly significantly underestimated the diversity and abundance of *Lasioglossum* bees visiting oilseed rape, given that many studies did not include detailed quantitative data on this genus.

Our datasets also indicate that rare and specialist species may visit crop flowers when they are locally abundant or are especially attracted to crop flowers (MacLeod et al., 2020). Several rare species recorded in apple orchards are most common in south-east England, Britain's principal apple growing region, and bee species that are oligolectic on Brassicaceae were recorded in oilseed rape studies. Given that biodiversity benefits pollination (Dainese et al., 2019), strategies to support biodiverse crop communities may prove critical to sustain ecosystem service provision. Yet current agri-environment schemes options rarely consider rare species (Senapathi et al., 2015). There is however, a significant overlap in the floral resources used by common and rare crop pollinators (Sutter et al., 2017; MacLeod et al., 2020), and thus there are opportunities to promote both biodiversity and conservation in agricultural landscapes.

Our findings also offer an opportunity to anticipate potentially important future crop pollinators. For example, whilst a number of European crop flower visitors not presently recorded in British crop fields are currently geographically restricted, should they expand their range in the future, they could ameliorate the threat of ecological mismatches between current pollinators and crops due to climate change (Polce et al., 2013; Polce et al., 2014; Settele et al., 2016). Taken further, this information could be used to refine existing models of bee populations used to

project pollinator populations at large spatial scales (e.g. Gardner et al., 2020), which can assist in larger scale planning of pollinator management.

Identifying specific bee crop pollinating species, as we have done, can inform refinements to agri-environment schemes to promote more biodiverse communities in agricultural landscapes. For example, *Andrena* were the most speciose genus of bees identified across flower visitor categories in three of the crops. Currently European agri-environment measures to boost pollinator populations have focused on the creation of flower-rich habitats, including wildflower buffer strips (Wratten et al., 2012). Yet evidence suggests these are primarily visited by bumblebees, with solitary bees preferring non-sown, wild plants (Wood et al., 2015). In apple orchards for example, early-flying *Andrena* species have been positively associated with dandelions (*Taraxacum* agg.) rather than sown species, which often bloom later than apple flowers (Campbell et al., 2017). Reduced mowing regimes in orchards, and other crop areas, particularly in early spring could boost *Andrena* numbers and hence pollination. Such interventions are also likely to benefit early flying *Lasioglossum*, many species of which are known be attracted to yellow flowers in the family *Asteraceae*. *Osmia* species have also been demonstrated as efficient pollinators of apple, oilseed and strawberry crops (Abel et al., 2003; Garratt et al., 2016; Horth and Campbell, 2018), but as in this study, are frequently recorded in low numbers, likely due to a lack of suitable nesting and floral resources in agricultural landscapes for cavity nesting species (Gardner and Ascher, 2006; Blitzer et al., 2016). Incorporating hedgerow species such as Dog Rose and Bramble, alongside, areas of old and dead wood, around crop areas would provide both forage and nesting resources (Else and Edwards 2018; Gresty et al., 2018) for these and other cavity nesting bees. Future management to support long-tongued solitary bees could benefit field bean pollination. *Anthophora plumipes*, for example, prefers to nest in vertical soil profiles, which are not currently a common feature in agricultural landscapes.

## **4.2 Data constraints and limitations**

There are caveats to using foraging ecology to identify potential bee pollinators, as done here and elsewhere (Ahrenfeldt et al., 2015). There is a lack of published data for many bee species and others visit a wider range of flowers than can be realistically documented (Else and Edwards, 2018). As such, determining the status of bee species as crop flower visitors requires field survey data for confirmation. Yet comprehensive crop pollinator data is currently lacking as sampling is irregular, undertaken almost exclusively as part of bespoke research projects rather than systematic monitoring (Breeze et al., 2020). Furthermore, whilst census methods can provide information on floral associations, they require experienced surveyors to comprehensively record species richness (O'Connor et al., 2019). Across all four crops the only bees which were consistently identified to species level were large, conspicuous ones from the genera *Bombus* and *Andrena*. Small and inconspicuous species, particularly from the genus *Lasioglossum*, were often only extensively sampled in the pan trap surveys. Additionally, whilst the visitation rate of dominant species is strongly correlated to pollination service delivery (Winfree et al., 2015; Fijen et al., 2018), the assumption here and elsewhere that quantitative visitation data can be used to infer pollination (Kleijn et al., 2015), neglects to factor in that flower visitation alone is not a perfect proxy for pollination (King et al., 2013; Senapathi et al., 2015; Ollerton, 2017). Certain physiological and behavioural traits also influence pollination service delivery (Martins et al., 2015). Further detailed data and research is required before any definitive conclusions can be made about the contributions of individual bee species to crop pollination.

## 5. Conclusions

Given the importance of wild pollinators and the detrimental impacts of conventional agriculture on their populations it is unsurprising that the management of wild and managed pollinating insects is considered a critical step for future food security (Garibaldi et al., 2019; Kleijn et al., 2019; Rollin and Garibaldi et al., 2019; Reilly et al., 2020). Yet information on which species contribute most to ecosystem service delivery has long been elusive (Kremen and Chaplin-Kramer, 2007) despite its critical importance for both monitoring and conservation

measures. Here we combine ecological and field data to provide a uniquely comprehensive overview of the crop pollinating bees of a single region, Great Britain. Whilst we have focused on Great Britain, a similar approach would be applicable across Europe, and could also be applied to non-bee species that have been identified as important crop pollinators (Rader et al., 2016). Our research bolsters evidence that many wild bee species, including rare and specialised ones, may contribute to crop pollination (Klein et al., 2003; Sutter et al., 2017; Winfree et al., 2018; MacLeod et al., 2020), thus it can be argued that agri-environment scheme options should not focus solely on dominant crop pollinators.

Future climatic changes threaten to further deplete already impoverished bee populations (Soroye et al., 2020) and create spatial mismatches between crops and their pollinators, which could exacerbate existing pollination deficits (Polce et al., 2014). To that end, the species identified as possible crop pollinators could represent an as yet untapped pollinator resource. Whilst some species may not currently visit crops due to ecological or environmental constraints, they could be assisted to expand by dedicated conservation measures in agricultural landscapes, allowing them to compensate for any declines in current crop pollinating species. Many such species are solitary, which presently benefit much less from agri-environment schemes than social species (Wood et al., 2015b, 2016a, 2016b; Gresty et al., 2018). As such land managers may need to re-evaluate existing pollinator management interventions and consider a broader range of species to safeguard the ecosystem service of crop pollination in an uncertain future.

#### **Declaration of Competing Interest**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

#### **Authors' contributions**

LH conceived the ideas, analysed the data and wrote the manuscript. MG, TB and TO contributed to the conceptual development and manuscript revisions. All other authors provided data and contributed to manuscript revisions.

## **Funding**

LH was funded by NERC QMEE CDT. EJB was funded by a BBSRC PhD studentship under grant BB/F016581/1. LB was supported by the Scholarship Program of the German Federal Environmental Foundation (Deutsche Bundesstiftung Umwelt, DBU, AZ 20014/302). AJC was funded by the BBSRC and Syngenta UK as part of a case award PhD (grant no. 1518739). AE was funded by the Swiss National Science Foundation (grant number 405940-115642). DG and A-MK were funded by grant PCIN2014-145-C02-02 (MinECo; EcoFruit project BiodivERsA-FACCE2014-74). MG was supported by Establishing a UK Pollinator Monitoring and Research Partnership (PMRP) a collaborative project funded by Defra, the Welsh and Scottish Governments, JNCC and project partners. GAdG was funded via research projects BO-11-011.01-051 and BO-43-011.06-007, commissioned by the Dutch Ministry of Agriculture, Nature and Food Quality. DK was funded by the Dutch Ministry of Economic Affairs (BO-11-011.01-011). AK-H was funded by the NKFIH project (FK123813), the Bolyai János Fellowship of the MTA, the ÚNKP-19-4-SZIE-3 New National Excellence Program of the Ministry for Innovation and Technology, and together with RF by the Hungarian Scientific Research Fund OTKA 101940. MM was funded by Waitrose & Partners, Fruition PO, and the University of Worcester. MM was funded by grant INIA-RTA2013-00139-C03-01 (MinECo and FEDER). BBP and RFS were funded by the UK Natural Environment Research Council as part of Wessex BESS (ref. NE/J014680/1). NJV was funded by the Walloon Region (Belgium) Direction générale opérationnelle de l'Agriculture, des Ressources naturelles et de l'Environnement (DGO3) for the "Modèle permaculturel" project on biodiversity in micro-farms, FNRS/FWO joint programme EOS — Excellence Of Science CliPS: Climate change and its impact on Pollination Services (project 30947854)". CW was funded by the Deutsche Forschungsgemeinschaft (DFG) (Project number 405945293). BW was funded by the Natural

Environment Research Council (NERC) under research programme NE/N018125/1 ASSIST – Achieving Sustainable Agricultural Systems [www.assist.ceh.ac.uk](http://www.assist.ceh.ac.uk). TB and TO are supported by BBSRC, NERC, ESRC and the Scottish Government under the Global Food Security Programme (Grant BB/R00580X/1). MG and data collection was funded by Insect Pollinators Initiative funded jointly by a grant from BBSRC, Defra, NERC, the Scottish Government and the Wellcome Trust, under the Living with Environmental Change Partnership and the Sustainable Management of Orchard Pollination Services Project.

## Acknowledgements

Thank you to Jean-Marc Molenberg for assistance in field surveys and specimen curation for the apple data collection in Belgium. Thanks to Ignasi Bartomeus for providing additional field survey data. Also thank you to Samantha Ardin, D.A. Bond, Hannah Feltham, Harriet Griffin, E.J.M. Kirby, Jean-Noel Tasei and Verena Riedinger for published crop field survey data that were used in analyses.

Figure 1 – Crop map courtesy of Alice Haughan, University of Reading.

Figure 4 – Bee photographs courtesy of Nicolas J. Vereecken and Stéphane De Greef.

## References

- Abel, C.A., Wilson, R.L. and Luhman, R.L., 2003. Pollinating efficacy of *Osmia cornifrons* and *Osmia lignaria* subsp. *lignaria* (Hymenoptera: Megachilidae) on three Brassicaceae species grown under field cages. *Journal of Entomological Science*, 38(4), 545-552.
- Ahrenfeldt, E.J., Klatt, B.K., Arildsen, J., Trandem, N., Andersson, G.K.S., Tscharrntke, T., ... and Sigsgaard, L., 2015. Pollinator communities in strawberry crops–variation at multiple spatial scales. *Bulletin of Entomological Research*, 105(4), 497-506.
- Aizen, M. A. and Harder, L. D., 2009. The global stock of domesticated honey bees is growing slower than agricultural demand for pollination. *Current Biology*, 19, 915–918.



330 Bartomeus, I., Potts, S.G., Steffan-Dewenter, I., Vaissiere, B.E., Woyciechowski, M.,  
 331 Krewenka, K.M., ... and Bommarco, R., 2014. Contribution of insect pollinators to crop yield  
 332 and quality varies with agricultural intensification. *PeerJ*, 2, 328.

333 Blitzer, E.J., Gibbs, J., Park, M.G. and Danforth, B.N., 2016. Pollination services for apple  
 334 are dependent on diverse wild bee communities. *Agriculture, Ecosystems & Environment*,  
 335 221, 1-7.

336 Bond, D.A. and Kirby, E.J.M., 1999. *Anthophora plumipes* (Hymenoptera: Anthophoridae) as  
 337 a pollinator of broad bean (*Vicia faba major*). *Journal of Apicultural Research*, 38(3-4), 199-  
 338 203.

339 Bossert, S., 2015. Recognition and identification of species in the *Bombus lucorum*-complex-  
 340 A review and outlook. *bioRxiv*, 011379.

341 Breeze, T.D., Vaissière, B.E., Bommarco, R., Petanidou, T., Seraphides, N., Kozák, L., ...  
 342 and Moretti, M., 2014. Agricultural policies exacerbate honeybee pollination service supply-  
 343 demand mismatches across Europe. *PloS one*, e82996.

344 Breeze T.D., Bailey A.P., Balcombe K.G., Brereton T., Comont R., Edwards M., ... and  
 345 Carvell C., 2020. Pollinator Monitoring More than Pays for Itself. *Journal of Applied Ecology*,  
 346 In Press

347 BWARS., 2020. Bees, Wasps & Ants Recording Society. <https://www.bwars.com/home>

348 Campbell, A. J., Wilby, A., Sutton, P. and Wäckers, F. L., 2017. Do sown flower strips boost  
 349 wild pollinator abundance and pollination services in a spring-flowering crop? A case study  
 350 from UK cider apple orchards. *Agriculture, Ecosystems & Environment*, 239, 20-29.

351 Carvell, C., Isaac, N., Jitlal, M., Peyton, J., Powney, G., Roy, D., ... and Roy, H., 2017.  
 352 Design and testing of a national pollinator and pollination monitoring framework. Technical  
 353 Report. Department for Environment, Food and Rural Affairs.

354 Catarino, R., Bretagnolle, V., Perrot, T., Vialloux, F. and Gaba, S., 2019. Bee pollination  
 355 outperforms pesticides for oilseed crop production and profitability. *Proceedings of the Royal*  
 356 *SocietyB*, 286 (1912), 20191550.

357 Dainese, M., Martin, E.A., Aizen, M., Albrecht, M., Bartomeus, I., Bommarco, R., ... and  
 358 Ghazoul, J., 2019. A global synthesis reveals biodiversity-mediated benefits for crop  
 359 production. *bioRxiv*, 554170.

360 Else, G.R., Bolton, B. and Broad, G.R., 2016. Checklist of British and Irish Hymenoptera-  
 361 aculeates (Apoidea, Chrysidoidea and Vespoidea). *Biodiversity Data Journal*, 4.

362 Else, G.R. & Edwards, M., 2018. *Handbook of the Bees of the British Isles: Volume 2*. Ray  
 363 Society.

364 Fijen, T.P., Scheper, J.A., Boom, T.M., Janssen, N., Raemakers, I. and Kleijn, D., 2018.  
 365 Insect pollination is at least as important for marketable crop yield as plant quality in a seed  
 366 crop. *Ecology Letters*, 21 (11), 1704-1713.

367 Gardner, K.E. and Ascher, J.S., 2006. Notes on the native bee pollinators in New York apple  
 368 orchards. *Journal of the New York Entomological Society*, 114 (1), 86-91.

369 Gardner E., Breeze T.D., Clough Y., Smith H., Baldock K., Campbell A., ... and Oliver T.,  
 370 2020. Reliably Predicting Pollinator Abundance: Challenges of Process Based Ecological  
 371 Models; *Methods in Ecology and Evolution*, In Press

372 Garibaldi, L.A., Steffan-Dewenter, I., Kremen, C., Morales, J.M., Bommarco, R.,  
 373 Cunningham, S.A., ... and Holzschuh, A., 2011. Stability of pollination services decreases  
 374 with isolation from natural areas despite honey bee visits. *Ecology Letters*, 14 (10), 1062-  
 375 1072.

376 Garibaldi, L.A., Bartomeus, I., Bommarco, R., Klein, A.M., Cunningham, S.A., Aizen, M.A.,  
 377 ... and Morales, C.L., 2015. Trait matching of flower visitors and crops predicts fruit set  
 378 better than trait diversity. *Journal of Applied Ecology*, 52, 1436-1444.

379 Garibaldi, L.A., Pérez-Méndez, N., Garratt, M.P., Gemmill-Herren, B., Miguez, F.E. and  
 380 Dicks, L.V., 2019. Policies for ecological intensification of crop production. *Trends in ecology*  
 381 & *evolution*, 34(4), pp.282-286.

382 Garratt, M.P., Coston, D.J., Truslove, C.L., Lappage, M.G., Polce, C., Dean, R., ... and  
 383 Potts, S.G., 2014a. The identity of crop pollinators helps target conservation for improved  
 384 ecosystem services. *Biological Conservation*, 169, 128-135.

385 Garratt, M.P., Breeze, T.D., Jenner, N., Polce, C., Biesmeijer, J.C. and Potts, S.G., 2014b.  
 386 Avoiding a bad apple: Insect pollination enhances fruit quality and economic value.  
 387 *Agriculture, ecosystems & environment*, 184, pp.34-40.

388 Garratt, M.P.D., Breeze, T.D., Boreux, V., Fountain, M.T., Mckerchar, M., Webber, S.M., ...  
 389 and Biesmeijer, J.C., 2016. Apple pollination: demand depends on variety and supply  
 390 depends on pollinator identity. *PloS one*, 11(5), e0153889.

391 Garratt, M.P.D., Potts, S.G., Banks, G., Hawes, C., Breeze, T.D., O'Connor, R.S. and  
 392 Carvell, C., 2019. Capacity and willingness of farmers and citizen scientists to monitor crop  
 393 pollinators and pollination services. *Global Ecology and Conservation*, 20, e00781.

394 Godfray, H. C. J. and Garnett, T., 2014. Food security and sustainable intensification.  
 395 *Philosophical Transactions of the Royal Society B: Biological Sciences*, 369, 20120273.

396 Grab, H., Branstetter, M.G., Amon, N., Urban-Mead, K.R., Park, M.G., Gibbs, J., ... and  
 397 Danforth, B.N., 2019. Agriculturally dominated landscapes reduce bee phylogenetic diversity  
 398 and pollination services. *Science*, 363(6424), 282-284.

399 Gresty, C.E., Clare, E., Devey, D.S., Cowan, R.S., Csiba, L., Malakasi, P., ... and Willis,  
 400 K.J., 2018. Flower preferences and pollen transport networks for cavity-nesting solitary bees:  
 401 Implications for the design of agri-environment schemes. *Ecology and evolution*, 8(5), 7574-  
 402 7587.

403 Hoehn, P., Tschardtke, T., Tylianakis, J.M. and Steffan-Dewenter, I., 2008. Functional group  
 404 diversity of bee pollinators increases crop yield. *Proceedings of the Royal Society of London*  
 405 *B: Biological Sciences*, 275(1648), 2283-2291.

406 Horth, L. and Campbell, L.A., 2018. Supplementing small farms with native mason bees  
 407 increases strawberry size and growth rate. *Journal of Applied Ecology*, 55(2), 591-599.

408 IPBES., 2016. Deliverable 3a: Thematic assessment of pollinators, pollination and food  
 409 production.  
 410 [http://www.ipbes.net/sites/default/files/downloads/pdf/3a\\_pollination\\_individual\\_chapters\\_20](http://www.ipbes.net/sites/default/files/downloads/pdf/3a_pollination_individual_chapters_20161124.pdf)  
 411 [161124.pdf](http://www.ipbes.net/sites/default/files/downloads/pdf/3a_pollination_individual_chapters_20161124.pdf)

412 King, C., Ballantyne, G. and Willmer, P.G., 2013. Why flower visitation is a poor proxy for  
 413 pollination: measuring single-visit pollen deposition, with implications for pollination networks  
 414 and conservation. *Methods in Ecology and Evolution*, 4(9), 811-818.

415 Klein, A.M., Steffan-Dewenter, I. and Tschardtke, T., 2003. Fruit set of highland coffee  
 416 increases with the diversity of pollinating bees. *Proceedings of the Royal Society of London*.  
 417 *Series B: Biological Sciences*, 270(1518), 955-961.

418 Kleijn, D., Winfree, R., Bartomeus, I., Carvalheiro, L.G., Henry, M., Isaacs, R., ... and  
 419 Ricketts, T.H., 2015. Delivery of crop pollination services is an insufficient argument for wild  
 420 pollinator conservation. *Nature Communications*, 6, 7414.

421 Kleijn, D., Bommarco, R., Fijen, T.P., Garibaldi, L.A., Potts, S.G. and van der Putten, W.H.,  
 422 2019. Ecological intensification: bridging the gap between science and practice. *Trends in*  
 423 *ecology & evolution*, 34 (2), pp.154-166.

424 Kremen, C., Williams, N.M. and Thorp, R.W., 2002. Crop pollination from native bees at risk  
 425 from agricultural intensification. *Proceedings of the National Academy of Sciences*, 99(26),  
 426 16812-16816.

427 Kremen, C. and Chaplin-Kramer, R., 2007. Insects as providers of ecosystem services: crop  
 428 pollination and pest control. In: Insect conservation biology: proceedings of the royal  
 429 entomological society's 23rd symposium, 349-382, Wallingford, UK: CABI Publishing.

430 MacLeod, M., Reilly, J., Cariveau, D.P., Genung, M.A., Roswell, M., Gibbs, J. and Winfree,  
 431 R., 2020. How much do rare and crop-pollinating bees overlap in identity and flower  
 432 preferences? *Journal of Applied Ecology*, 57(2), 413-423.

433 Martins, K.T., Gonzalez, A. and Lechowicz, M.J., 2015. Pollination services are mediated by  
 434 bee functional diversity and landscape context. *Agriculture, Ecosystems &*  
 435 *Environment*, 200, 12-20.

436 Michener, C.D., 2000. *The bees of the world* (Vol. 1). JHU press.

437 O'Connor, R.S., Kunin, W.E., Garratt, M.P., Potts, S.G., Roy, H.E., Andrews, C., ... and  
 438 Morris, R.K., 2019. Monitoring insect pollinators and flower visitation: the effectiveness and  
 439 feasibility of different survey methods. *Methods in Ecology and Evolution*, 10(12), 2129-  
 440 2140.

441 Oliver, T.H., Heard, M.S., Isaac, N.J., Roy, D.B., Procter, D., Eigenbrod, F., ... and Proença,  
 442 V., 2015. Biodiversity and resilience of ecosystem functions. *Trends in Ecology &*  
 443 *Evolution*, 30(11), 673-684.

444 Ollerton, J., 2017. Pollinator diversity: distribution, ecological function, and conservation.  
 445 *Annual Review of Ecology, Evolution, and Systematics*, 48, 353-376.

446 Perrot, T., Gaba, S., Roncoroni, M., Gautier, J.L. and Bretagnolle, V., 2018. Bees increase  
 447 oilseed rape yield under real field conditions. *Agriculture, Ecosystems & Environment*, 266,  
 448 39-48.

449 Polce, C., Termansen, M., Aguirre-Gutiérrez, J., Boatman, N.D., Budge, G.E., Crowe, A., ...  
 450 and Somerwill, K.E., 2013. Species distribution models for crop pollination: a modelling  
 451 framework applied to Great Britain. *PloS one*, 8(10), e76308.

452 Polce, C., Garratt, M.P., Termansen, M., Ramirez-Villegas, J., Challinor, A.J., Lappage,  
 453 M.G., ... and Somerwill, K.E., 2014. Climate-driven spatial mismatches between British  
 454 orchards and their pollinators: increased risks of pollination deficits. *Global Change Biology*,  
 455 20(9), 2815-2828.

456 Powney, G.D., Carvell, C., Edwards, M., Morris, R.K., Roy, H.E., Woodcock, B.A. and Isaac,  
 457 N.J., 2019. Widespread losses of pollinating insects in Britain. *Nature Communications*,  
 458 10(1), 1018.

459 Rader, R., Howlett, B.G., Cunningham, S.A., Westcott, D.A. and Edwards, W., 2012. Spatial  
 460 and temporal variation in pollinator effectiveness: do unmanaged insects provide consistent  
 461 pollination services to mass flowering crops? *Journal of Applied Ecology*, 49(1), 126-134.

462 Rader, R., Bartomeus, I., Garibaldi, L.A., Garratt, M.P., Howlett, B.G., Winfree, R., ... and  
 463 Bommarco, R., 2016. Non-bee insects are important contributors to global crop pollination.  
 464 *Proceedings of the National Academy of Sciences*, 113(1), pp.146-151.

465 Reilly, J.R., Artz, D.R., Biddinger, D., Bobiwash, K., Boyle, N.K., Brittain, C., ... and Ellis,  
 466 J.D., 2020. Crop production in the USA is frequently limited by a lack of pollinators.  
 467 *Proceedings of the Royal Society B*, 287(1931), p.20200922.

468 Russo, L., Park, M.G., Blitzer, E.J. and Danforth, B.N., 2017. Flower handling behavior and  
 469 abundance determine the relative contribution of pollinators to seed set in apple  
 470 orchards. *Agriculture, Ecosystems & Environment*, 246, 102-108.

471 Scheper, J., Holzschuh, A., Kuussaari, M., Potts, S.G., Rundlöf, M., Smith, H.G. and Kleijn,  
 472 D., 2013. Environmental factors driving the effectiveness of European agri-environmental  
 473 measures in mitigating pollinator loss—a meta-analysis. *Ecology Letters*, 16(7), 912-920.

474 Senapathi, D., Biesmeijer, J.C., Breeze, T.D., Kleijn, D., Potts, S.G. and Carvalheiro, L.G.,  
 475 2015. Pollinator conservation—the difference between managing for pollination services and  
 476 preserving pollinator diversity. *Current Opinion in Insect Science*, 12, 93-101.

477 Settele, J., Bishop, J. and Potts, S.G., 2016. Climate change impacts on pollination. *Nature*  
 478 *Plants*, 2(7),16092.

479 Soroye, P., Newbold, T. and Kerr, J., 2020. Climate change contributes to widespread  
 480 declines among bumble bees across continents. *Science*, 367(6478), 685-688.

481 Sutter, L., Jeanneret, P., Bartual, A.M., Bocci, G. and Albrecht, M., 2017. Enhancing plant  
 482 diversity in agricultural landscapes promotes both rare bees and dominant crop-pollinating  
 483 bees through complementary increase in key floral resources. *Journal of Applied*  
 484 *Ecology*, 54(6),1856-1864.

485 Tonietto, R.K. and Larkin, D.J., 2018. Habitat restoration benefits wild bees: A meta-  
 486 analysis. *Journal of Applied Ecology*, 55(2), 582-590.

487 Vanbergen, A.J., Heard, M.S., Breeze, T., Potts, S.G. and Hanley, N., 2014. Status and  
 488 value of pollinators and pollination services. <http://nora.nerc.ac.uk/id/eprint/505259/>

489 Vázquez, D.P., Morris, W.F. and Jordano, P., 2005. Interaction frequency as a surrogate for  
 490 the total effect of animal mutualists on plants. *Ecology Letters*, 8(10), 1088-1094.

491 Webb, J., Heaver, D., Lott, D., Dean, H.J., van Breda, J., Curson, ... and Foster, G., 2018.  
 492 Pantheon - database version 3.7.6. <https://www.brc.ac.uk/pantheon/>

493 Westphal, C., Bommarco, R., Carré, G., Lamborn, E., Morison, N., Petanidou, T., ... and  
 494 Vaissière, B.E., 2008. Measuring bee diversity in different European habitats and  
 495 biogeographical regions. *Ecological Monographs*, 78(4), 653-671.

496 Winfree, R., W. Fox, J., Williams, N.M., Reilly, J.R. and Cariveau, D.P., 2015. Abundance of  
 497 common species, not species richness, drives delivery of a real-world ecosystem  
 498 service. *Ecology letters*, 18(7), 626-635.

499 Winfree, R., Reilly, J.R., Bartomeus, I., Cariveau, D.P., Williams, N.M. and Gibbs, J., 2018.  
 500 Species turnover promotes the importance of bee diversity for crop pollination at regional  
 501 scales. *Science*, 359(6377), 791-793.

502 Wolf, S., Rohde, M. and Moritz, R.F., 2010. The reliability of morphological traits in the  
 503 differentiation of *Bombus terrestris* and *B. lucorum* (Hymenoptera:  
 504 Apidae). *Apidologie*, 41(1), 45-53.

505 Wood, T.J., Holland, J.M., Hughes, W.O. and Goulson, D., 2015a. Targeted agri-  
 506 environment schemes significantly improve the population size of common farmland  
 507 bumblebee species. *Molecular Ecology*, 24(8), 1668-1680.

508 Wood, T.J., Holland, J.M. and Goulson, D., 2015b. Pollinator-friendly management does not  
 509 increase the diversity of farmland bees and wasps. *Biological Conservation*, **187**, pp.120-  
 510 126.

511 Wood, T.J., Holland, J.M. and Goulson, D., 2016a. Providing foraging resources for solitary  
 512 bees on farmland: current schemes for pollinators benefit a limited suite of species. *Journal*  
 513 *of Applied Ecology*, 54(1),323-333.

514 Wood, T.J., Holland, J.M. and Goulson, D., 2016b. Diet characterisation of solitary bees on  
 515 farmland: dietary specialisation predicts rarity. *Biodiversity and Conservation*, 25(13), 2655-  
 516 2671.

517 Woodcock, B.A., Edwards, M., Redhead, J., Meek, W.R., Nuttall, P., Falk, S., ... and Pywell,  
 518 R.F., 2013. Crop flower visitation by honeybees, bumblebees and solitary bees: Behavioural  
 519 differences and diversity responses to landscape. *Agriculture, Ecosystems &*  
 520 *Environment*, 171, 1-8.

521 Woodcock, B.A., Garratt, M.P.D., Powney, G.D., Shaw, R.F., Osborne, J.L., Soroka, J., ...  
 522 and Jauker, F., 2019. Meta-analysis reveals that pollinator functional diversity and abundance  
 523 enhance crop pollination and yield. *Nature Communications*, 10(1) 1-10.

524 Wratten, S.D., Gillespie, M., Decourtye, A., Mader, E. and Desneux, N., 2012. Pollinator  
 525 habitat enhancement: benefits to other ecosystem services. *Agriculture, Ecosystems &*  
 526 *Environment*, 159, pp.112-122.