

The relationship between sensory reactivity, intolerance of uncertainty and anxiety subtypes in preschool-age autistic children

Article

Published Version

Creative Commons: Attribution-Noncommercial 4.0

Open Access

MacLennan, K. ORCID: <https://orcid.org/0000-0002-3148-3450>, Rossow, T. and Tavassoli, T. ORCID: <https://orcid.org/0000-0002-7898-2994> (2021) The relationship between sensory reactivity, intolerance of uncertainty and anxiety subtypes in preschool-age autistic children. *Autism*, 25 (8). pp. 2305-2316. ISSN 1362-3613 doi: 10.1177/13623613211016110 Available at <https://centaur.reading.ac.uk/98478/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1177/13623613211016110>

Publisher: Sage

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in

the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

The relationship between sensory reactivity, intolerance of uncertainty and anxiety subtypes in preschool-age autistic children

Keren MacLennan^{id}, Timothy Rossow and Teresa Tavassoli

Autism

1–12

© The Author(s) 2021



Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/13623613211016110

journals.sagepub.com/home/aut

Abstract

Sensory reactivity differences are a diagnostic criterion of autism. Sensory hyperreactivity has been linked to intolerance of uncertainty and anxiety in autistic children. However, research is yet to explore the mediating relationships or sensory hyporeactivity, seeking and anxiety subtypes in preschool-age autistic children. Therefore, we aimed to elucidate the relationships between sensory reactivity, intolerance of uncertainty and anxiety subtypes in a heterogeneous group of 54 preschool-age autistic children, age 3–5 years, using observation and parent-report assessments. Correlational analysis found sensory hyperreactivity, intolerance of uncertainty and total anxiety were significantly intercorrelated. In addition, sensory hyperreactivity was significantly correlated with separation anxiety when controlling for autism traits. Serial mediation analyses indicated significant full mediation between sensory hyperreactivity and anxiety through intolerance of uncertainty, and significant full mediation between sensory hyperreactivity and intolerance of uncertainty through anxiety. Our results suggest that sensory hyperreactivity is a key early factor in the development of anxiety, and supports that intolerance of uncertainty is an important interrelated construct in the development and maintenance of anxiety in autism. Our findings have important implications for the development of effective interventions. However, due to limitations with the measures, our research also highlights a pressing need for objective assessments of anxiety and intolerance of uncertainty that can be used with preschool-age autistic children.

Lay abstract

This study found links between greater sensory hyperreactivity (e.g., over-sensitive to sensory input), intolerance of uncertainty and anxiety, including separation anxiety, in autistic pre-schoolers. Sensory hyperreactivity may predict both anxiety and intolerance of uncertainty, and anxiety and intolerance of uncertainty may both be mutually important, mediating factors. These findings have implications for early anxiety interventions. But there is a pressing need for objective assessments that can be used with preschool-age autistic children.

Keywords

anxiety, autism, children, mental health, preschool, sensory, uncertainty

Introduction

Autism spectrum conditions (ASC) are neurodevelopmental conditions with characteristics related to restricted and repetitive behaviours and interests (RRBs), and social interaction and communication differences (*Diagnostic and Statistical Manual of Mental Disorders* (5th ed.; DSM-5; American Psychiatric Association, 2013)). Sensory reactivity differences, including sensory hyperreactivity (e.g.

strong reaction to touch), hyporeactivity (e.g. no or slower reaction to sounds) and seeking (e.g. fascinated by lights)

University of Reading, UK

Corresponding author:

Keren MacLennan, The Centre for Autism, University of Reading, Earley, Reading, RG6 6DZ, UK.

Email: k.r.maclennan@pgr.reading.ac.uk

are commonly experienced by approximately 65%–85% of autistic¹ children (Leekam et al., 2007; Miller et al., 2007; Tavassoli et al., 2016). Due to this prevalence, sensory reactivity differences have become a diagnostic criterion for ASC, under the RRBs domain (DSM-5; American Psychiatric Association, 2013). However, they are also common across other neurodevelopmental conditions, such as attention deficit hyperactivity disorder (ADHD; Miller et al., 2012). Sensory reactivity differences can be experienced differentially and across a range of modalities, such as vision, hearing and touch (Miller et al., 2007). Although some sensory experiences may be pleasant, others can be extremely distressing, which can greatly impact developmental outcomes and quality of life (Pfeiffer et al., 2005; Uljarević et al., 2016). Sensory hyperreactivity has previously been linked to anxiety and intolerance of uncertainty (IU) in autistic individuals (Carpenter et al., 2019; Green et al., 2012; Hwang et al., 2019; Neil et al., 2016; Pfeiffer et al., 2005; Uljarević et al., 2016). However, research has yet to explore the relationship between sensory reactivity differences, including hyperreactivity, hyporeactivity and seeking, IU and anxiety subtypes in preschool-age autistic children. Understanding the early emergence of these relationships has important implications for early anxiety interventions.

Sensory reactivity differences and anxiety

Autistic children are twice as likely to develop an anxiety condition compared to non-autistic children (van Steensel et al., 2011). Research has suggested that around 49% of autistic children will have at least one co-occurring anxiety condition: commonly generalised anxiety disorder (GAD), social anxiety, or specific phobia, or anxiety-related conditions such as obsessive-compulsive disorder (OCD) (Simonoff et al., 2008; van Steensel et al., 2011; White et al., 2009). Anxiety diagnoses are not always straight forward due to some degree of overlap between traits common in autistic children and anxiety symptomology, for instance, social avoidance which can be present in both autism and social anxiety (Kerns & Kendall, 2012; White et al., 2009). As well as presenting with traditional anxiety symptomology, autistic people can also present with some unique symptoms, such as unusual phobias, sensory fears, social distress unrelated to fears of negative evaluation and compulsive behaviours that are not driven by distress relief (Kerns et al., 2014, 2017, 2020; South & Rodgers, 2017). Anxiety symptoms develop early (i.e. before the age of 6 years) and are enhanced in autistic children and increase over time (Vasa et al., 2020). Thus, there is an imperative need to identify early risk factors.

Research has repeatedly shown links between sensory hyperreactivity and anxiety in autistic children (Green & Ben-Sasson, 2010; Green et al., 2012; Pfeiffer et al., 2005;

Vasa et al., 2020; Wigham et al., 2015). Sensory hyperreactivity may be an early risk factor for the development of anxiety as it has been suggested to emerge earlier and to unidirectionally predict anxiety in both autistic and non-autistic samples (Carpenter et al., 2019; Green et al., 2012). Green et al. (2012) explored the bidirectional relationship between sensory hyperreactivity and anxiety in autistic toddlers in a longitudinal study. They found that while sensory hyperreactivity remained stable over time, anxiety increased. Furthermore, sensory hyperreactivity predicted an increase in anxiety; however, anxiety did not predict changes in sensory hyperreactivity (Green et al., 2012). There is also research in autistic children and adolescents to suggest that sensory hyperreactivity is implicated in specific anxiety conditions, such as separation anxiety, social anxiety and specific phobia (Black et al., 2017; MacLennan et al., 2020; Pickard et al., 2020). Understanding if sensory reactivity is an early predictor for certain anxiety symptomology pertaining to the current diagnostic criteria has important clinical implications for early identification and targeted intervention therapies.

There is limited and inconclusive research into the relationships between both sensory hyporeactivity, and sensory seeking and anxiety in autistic children. Some research has found links between sensory hyporeactivity and anxiety in autistic children and adolescents (Glod et al., 2019), as well as specific links to social anxiety (MacLennan et al., 2020). However, links between sensory hyporeactivity and anxiety have not been consistently found in research (Wigham et al., 2015). Sensory seeking is often neglected in research and direct links to anxiety have rarely been found (Levit-Binnun et al., 2014; Lidstone et al., 2014; MacLennan et al., 2020). There is evidence to suggest that sensory seeking may be indirectly linked with anxiety through sensory hyperreactivity and over-focusing behaviours (Liss et al., 2006). Therefore, it would be advantageous to continue to explore how both sensory hyporeactivity and seeking relate to anxiety in autistic children.

IU, sensory reactivity differences and anxiety

IU is broadly characterised by a difficulty with situations or contexts that are unforeseen or unpredictable (Buhr & Dugas, 2006). It is considered to be a key cognitive component in the development and maintenance of anxiety through negatively appraising ambiguous or uncertain events (Dugas et al., 1998; Freeston et al., 1994). Greater IU has been linked with anxiety in non-autistic populations, and may be a key transdiagnostic feature across anxiety conditions (Carleton et al., 2012). Although argued to be a distinct construct, there is also some degree of similarity between the characteristics of IU and some autistic traits, such as difficulty with change and unpredictability (Boulter et al., 2014; Chamberlain et al., 2013). However,

IU has been identified as a recognisable construct in autistic children and has been found to be higher in autistic groups (Boulter et al., 2014; Hodgson et al., 2017; Vasa et al., 2018).

Existing research has aimed to elucidate the relationship between sensory reactivity, anxiety and IU. A key theoretical model of anxiety in autism by South and Rodgers (2017) proposes that IU is a principal mediator in the relationship between sensory reactivity and anxiety. In support of this, research has found IU to partially mediate the relationship between both sensory hyperreactivity, and hyporeactivity and anxiety in autistic adults (Hwang et al., 2019). However, there is currently mixed, and sometimes conflicting, evidence to support this. Research has found IU to mediate the relationship between autism traits and anxiety, as well as autism traits and sensory hyperreactivity (Boulter et al., 2014; Hwang et al., 2019; Neil et al., 2016). In addition, research has found anxiety to mediate the relationship between IU and sensory hyperreactivity in autistic children (Neil et al., 2016). Therefore, research is still needed to elucidate the relationship between sensory reactivity, IU and anxiety in autistic children.

The present study

Symptoms of anxiety are present in autistic children at an early age and there is an early, unidirectional association between sensory hyperreactivity and anxiety in autistic toddlers. Although there may be a differential relationship between sensory hyperreactivity and symptoms associated with anxiety subtypes, such as social anxiety and specific phobia, this is an under-researched area. In addition, sensory hyporeactivity and seeking are often neglected in research, and there is inconclusive evidence of the relationships between sensory reactivity, IU and anxiety.

Fundamentally, the links between sensory reactivity, IU and anxiety have only been explored in autistic children above the age of 6 years, without intellectual disability (ID) and with proficient language skills, and research with preschool-age autistic children (age 3–5 years) is yet to be done. Targeting risk factors for anxiety in its earliest stages is important to improve later outcomes (Rapee et al., 2010; Yap et al., 2016). Ages 3–5 years are a critical time when sensory reactivity differences and anxiety symptoms are emerging in autistic children, and early interventions for anxiety already show efficacy (Ben-Sasson et al., 2007, 2008; Driscoll et al., 2020; Green et al., 2012; Vasa et al., 2020).

Therefore, we aimed to elucidate the relationship between sensory reactivity differences, IU and anxiety in preschool-age autistic children. First, in line with previous findings (Carpenter et al., 2019; Neil et al., 2016; Pfeiffer et al., 2005; Uljarević et al., 2016; Wigham et al., 2015), we expected to find greater sensory hyperreactivity to be significantly related to greater anxiety symptoms, and for IU to be a mediator in the predictive relationship. Secondly,

as previous work including ours has found a differential relationship between sensory hyperreactivity and anxiety subtypes in older children (Black et al., 2017; MacLennan et al., 2020), we explored if sensory reactivity and IU were differentially related to anxiety subtypes, including GAD, separation anxiety symptoms, OCD, physical injury fears and social anxiety symptoms.

Moreover, due to conflicting theoretical models in existing literature, we aimed to test the predictive relationship between sensory reactivity, anxiety and IU. We expected that sensory hyperreactivity would predict anxiety, as hyperreactivity has been shown to predict anxiety unidirectionally (Green et al., 2012). But that the relationship between sensory hyperreactivity and anxiety would be mediated by IU, as IU has consistently been found to be a mediating, transdiagnostic construct (Boulter et al., 2014; Hwang et al., 2019; Neil et al., 2016; Wigham et al., 2015).

Methods

Participants

In total, 65 children with (1) a clinical diagnosis of autism or (2) suspected autism were recruited for this study. Fifty-four children (41 males, 13 females) aged 3–5 years (mean (M)=4.02, standard deviation (SD)=0.77) met inclusion criteria of (1) having a formal clinical diagnosis of autism which included a direct observation as well as developmental history ($n=42$) or (2) having suspected autism and meeting cut-off scores on the Autism Diagnostic Observation Schedule (ADOS-2; Lord et al., 2012) (Module 1 few-to-no words cut-off: 11; Module 1 some words cut-off: 8; Module 2 <5 years cut-off: 7; Module 2 ≥ 5 years cut-off: 8), and the Autism Spectrum Quotient: Child Version (AQ; Auyeung et al., 2008) (cut-off ≥ 76) ($n=12$), and were therefore included in the final analysis. Thirty-one children completed a Module 1, 19 completed a Module 2 and 4 were unable to complete the ADOS-2 during testing; however, these children already had a formal diagnosis (see Table 1 for additional demographic information).

To indicate the cognitive level of the sample, we conducted the Matrix reasoning subtests of the Wechsler Non-Verbal (WNV; Wechsler & Naglieri, 2006) and the British Ability Scale (BAS; Elliott et al., 1983). However, 31 participants were unable to complete the intelligence quotient (IQ) tasks either due to cognitive delay or behavioural difficulties (e.g. demand avoidance). Of these participants, 25 completed an ADOS-2 Module 1, 2 completed a Module 2 and 4 were also unable to complete the ADOS-2 assessment. IQ and verbal language level was taken into consideration in the analyses. Signed consent was obtained from parents, and verbal assent was obtained from children who were able to, otherwise, voluntary task compliance was

Table 1. Demographic characteristics of participants.

Characteristics	N	Range	M	SD
Age (years)	54	3–5	4.0	0.8
IQ	22	52–141	108.4	23.6
Sex		Percentage		
Male	41	75.9%		
Female	13	24.1%		
Ethnicity				
White British	32	59.2%		
White Swedish	1	1.9%		
Mixed Race – White Asian	5	9.2%		
Mixed Race – Other	3	5.5%		
Asian Indian	10	18.5%		
Asian Pakistani	1	1.9%		
North African	1	1.9%		
Undeclared	1	1.9%		

N: number, M: mean, SD: standard deviation; IQ: intelligence quotient.

considered assent. Families were provided with a social story for the children prior to testing to outline what their participation would entail. We recruited participants through the University of Reading Centre for Autism participant database, social media platforms, local NHS CAMHS services and through Autism Berkshire, a local autism organisation. Ethical approval was granted prior to the commencement of this study by the University of Reading Ethics Committee and the Berkshire Healthcare NHS Foundation Trust Health Research Authority.

Measures

Sensory Processing Scale Inventory. The Sensory Processing Scale Inventory (SPSI; Schoen et al., 2017) was used to measure sensory reactivity differences. It is a standardised, 96-item parent-report questionnaire that assesses the degree of sensory reactivity differences in children. It produces subscales for sensory hyperreactivity (47 items, e.g. ‘These garments bother my child: Seams in clothing’), sensory hyporeactivity (21 items, e.g. ‘My child has a less intense response than others to: Getting hurt, bruises or cuts’) and sensory seeking (28 items, e.g. ‘My child has difficulty disengaging from: Looking at spinning objects’). Items are scored dichotomously (1=yes, 0=no), with higher scores indicating a greater extent of sensory reactivity differences. The SPSI has strong internal reliability (hyperreactivity=0.89; hyporeactivity=0.88; sensory seeking=0.93) (Schoen et al., 2017). As in our previous work (MacLennan et al., 2020), we chose to use the SPSI over other commonly used measures such as the Sensory Profile (SP; Dunn, 1999) as the SPSI has a comparatively greater representation of hyporeactivity and seeking compared to the SP, which more heavily represents hyperreactivity (Neil et al., 2017; Uljarević et al., 2016).

Sensory Assessment for Neurodevelopmental Differences. The Sensory Assessment for Neurodevelopmental Differences (SAND; Siper et al., 2017) was also used to measure sensory reactivity differences. It is a standardised observational assessment and associated caregiver interview, which assesses sensory reactivity differences in line with the ASC diagnostic criteria (DSM-5; American Psychiatric Association, 2013). In the direct observation, children are presented with a range of sensory toys with tactile (e.g. textured toy), auditory (e.g. musical toy) and visual (e.g. flashing toy) components. The assessment commences and concludes with 1–2 min of unstructured play to help the child familiarise with the situation, followed by presentation of the sensory toys. The children are scored for behavioural responses consistent with sensory hyperreactivity, hyporeactivity and seeking. The administration is conducted by a trained examiner, and behaviours are scored dichotomously (i.e. 1=present, 0=not present) across 36 items, and if a behaviour is present, there is the additional coding of a severity score (1=mild, 2=moderate to severe).

The subsequent interview component requires the caregiver to indicate whether their child presents with the various sensory reactivity differences in visual, tactile and auditory domains (1=present, 0=not present), and to rate the severity if present (1=mild, 2=moderate to severe). The items of this interview map onto the scoring for the observation. Observed and reported scores can then combined to produce a total score (0–90) and subscales for sensory hyperreactivity, hyporeactivity and seeking (0–30), with higher scores reflecting greater sensory reactivity differences. The SAND has been found to have strong internal consistency (0.90), test–retest reliability (0.97, $p < 0.001$) and moderate convergent validity (−0.63) (Siper et al., 2017).

Preschool Anxiety Scale. The Preschool Anxiety Scale (PAS; Edwards et al., 2010) was used to measure anxiety symptoms. It is a standardised 30-item parent-report questionnaire that assesses anxiety symptoms in children aged 3–6 years, adapted from the Spence Children’s Anxiety Scale (SCAS; Spence, 1998). Items are scored on the scale 0 (no true at all) to 4 (very often true), with higher scores indicating a greater extent of anxiety symptoms, and scores above cut-off indicate clinically elevated symptoms. The PAS produces a score for total anxiety (total score 120) and subscales for symptoms related to GAD (five items, total score of 20, e.g. ‘Has difficulty stopping him/herself from worrying’), social anxiety (six items, total score of 24, e.g. ‘Worries that he or she will do something embarrassing in front of other people’), separation anxiety (five items, total score of 20, e.g. ‘Is reluctant to go to sleep without you or to sleep away from home’), physical injury fears (seven items, total score of 28, e.g. ‘Is afraid of crowded or closed-in places’) and OCD (five items, total score of 20, e.g. ‘Washes his or her hands over and over many times each day’). The PAS has moderate internal

consistency ($\alpha=0.70$) (Edwards et al., 2010). The PAS was selected over other measures of anxiety as it has subscales for early symptomology relating to anxiety subtypes.

Responses to Uncertainty and Low Environmental Structure questionnaire. The Responses to Uncertainty and Low Environmental Structure (RULES; Sanchez et al., 2017) was used to measure IU. It is a parent-report questionnaire and is validated to assess IU in children aged 3–10 years. It contains 17 items (e.g. ‘My child has a hard time coping with even minor changes’), rated on a scale of 1 (not at all) to 5 (very much). Scores can range from 17 to 85 with higher scores reflecting greater IU. The RULES has strong internal consistency ($\alpha=0.93$) and convergent validity in (0.67) (Sanchez et al., 2017).

Analysis

The data were analysed using SPSS 27 (IBM Corp, 2020) and JASP (JASP Team, 2020). Following normality checks, we conducted a Spearman’s bivariate correlation analysis to elucidate the correlational relationship between sensory reactivity differences, anxiety, IU and potential covariates. We then conducted partial correlation analyses and subgroup comparisons to ensure the relationships were independent of the effects of covariates. Significance was indicated by a Bonferroni’s corrected p -value of $p \leq 0.01$.

We then conducted a series of simple mediation analyses to understand the mediating relationship between sensory reactivity, anxiety and IU. We defined the analysis to conduct bootstrapping with 1000 resamples to adjust for measurement error and generated accelerated 95% confidence intervals (CIs). Full mediation is present if the indirect effect between the predictor and outcome variable is significant, and the direct effect is not significant. Partial mediation is present if both the direct and indirect effects between the predictor and outcome variable are significant. Significance is indicated by CIs not overlapping with zero and Bonferroni’s corrected p -value of $p \leq 0.008$.

Community involvement

The research questions for this project aligned with Autistica’s research priorities, which were identified by stakeholders. Throughout our project, we sought feedback from the families who took part in this study on our testing sessions and procedures, interpretations of our results and the language used in the article.

Results

Descriptive analyses

Descriptive analysis showed that the data were not normally distributed as a Shapiro–Wilk test of normality was

Table 2. Descriptive statistics for sensory reactivity differences (SPSI and SAND raw scores; z-score composites), anxiety (PAS raw scores), intolerance of uncertainty (RULES raw score) and autism traits (AQ and ADOS raw comparison scores).

	Mean	SD	Clinically elevated (%)
SPSI			
Hyperreactivity	12.8	5.5	–
Hyporeactivity	11.3	4.7	–
Seeking	10.0	5.0	–
SAND			
Hyperreactivity	9.2	3.9	–
Hyporeactivity	8.4	5.6	–
Seeking	17.4	3.6	–
PAS			
Total anxiety	33.2	24.3	46%
GAD	7.6	6.4	60%
Social anxiety	7.6	6.4	34%
OCD	3.9	4.1	46%
Physical injury fears	7.8	5.9	22%
Separation anxiety	6.3	5.1	42%
IU	50.4	19.6	–
ADOS-2 comparison score	6.2	2.8	–
AQ	100.1	17.6	–

SD: standard deviation; SPSI: Sensory Processing Scale Inventory; SAND: Sensory Assessment for Neurodevelopmental Differences; PAS: Preschool Anxiety Scale; RULES: Responses to Uncertainty and Low Environmental Structure; GAD: generalised anxiety disorder; OCD: obsessive–compulsive disorder; IU: intolerance of uncertainty; ADOS: Autism Diagnostic Observation Schedule-2;

significant for total anxiety and the anxiety subtypes ($p < 0.05$). However, no outliers were present in the data (see Table 2 for descriptive statistics). To avoid inflated correlations due to bias and multiple comparisons, parent-reported and observed sensory reactivity measures were combined (Kim & Lord, 2012; Spain et al., 2018). SPSI and SAND construct scores were significantly correlated for sensory hyperreactivity ($r(52)=0.60, p < 0.001$), hyporeactivity ($r(52)=0.38, p=0.005$) and seeking ($r(52)=0.50, p < 0.001$), suggesting both sensory measures are assessing related constructs. Therefore, SPSI and SAND z-scores were averaged to create standardised sensory composite scores for sensory hyperreactivity, hyporeactivity and seeking (Song et al., 2013).

Correlational analyses

Spearman’s correlation analysis showed that sensory hyperreactivity was significantly positively correlated with total anxiety, GAD, social anxiety, OCD, physical injury fears, separation anxiety and IU. IU was also significantly positively correlated with total anxiety, GAD, social anxiety, OCD, physical injury fears and separation anxiety (see Table 3 for correlations).

Table 3. Spearman's correlations between sensory reactivity differences (SPSI and SAND composite z-scores), anxiety (PAS), intolerance of uncertainty (RULES), autism traits (AQ) and age.

	Hyperreactivity	Hyporeactivity	Seeking	IU	AQ	Age
Total anxiety	0.520* (0.000)	-0.193 (0.180)	0.115 (0.426)	0.796* (0.000)	0.539* (0.000)	0.299 (0.035)
GAD	0.451* (0.001)	-0.212 (0.140)	0.099 (0.496)	0.728* (0.000)	0.461* (0.000)	0.337 (0.017)
Social anxiety	0.422* (0.002)	-0.222 (0.121)	0.130 (0.369)	0.708* (0.000)	0.534* (0.000)	0.143 (0.321)
OCD	0.414* (0.003)	-0.048 (0.742)	0.246 (0.085)	0.706* (0.000)	0.473* (0.000)	0.172 (0.232)
Physical injury fears	0.435* (0.002)	-0.280 (0.049)	-0.086 (0.551)	0.544* (0.000)	0.397* (0.004)	0.224 (0.118)
Separation anxiety	0.472* (0.000)	-0.181 (0.208)	0.132 (0.361)	0.727* (0.000)	0.435* (0.002)	0.453* (0.000)
IU	0.554* (0.000)	-0.025 (0.867)	0.221 (0.131)	–	–	–
AQ	0.413* (0.002)	0.059 (0.680)	0.268 (0.054)	0.641* (0.000)	–	–
Age	0.305 (0.028)	-0.266 (0.057)	-0.094 (0.509)	0.144 (0.328)	0.014 (0.924)	–

SPSI: Sensory Processing Scale Inventory; SAND: Sensory Assessment for Neurodevelopmental Differences; PAS: Preschool Anxiety Scale; RULES: Responses to Uncertainty and Low Environmental Structure; GAD: generalised anxiety disorder; OCD: obsessive-compulsive disorder; IU: intolerance of uncertainty;

*Correlation is significant at the $p \leq 0.01$ level (two-tailed) after a Bonferroni's correction.

Spearman's correlation analysis also showed that age was significantly positively correlated with separation anxiety. In addition, autism traits were significantly positively correlated with sensory hyperreactivity, total anxiety, GAD, social anxiety, OCD, physical injury fears, separation anxiety and IU. Post hoc partial correlation analyses controlling for age found that the correlations between sensory reactivity, anxiety subtypes and IU sustained their significance ($p \leq 0.01$) or non-significance ($p > 0.01$), suggesting the relationships are independent from the effects of age. However, when controlling for autism traits, the relationships between sensory hyperreactivity and GAD, social anxiety, OCD and physical injury fears were no longer significant ($p > 0.01$). Fisher's z -test was conducted to test if the significant correlation coefficient between sensory hyperreactivity and separation anxiety was significantly different from the non-significant correlation coefficients between sensory hyperreactivity and the other anxiety subtypes. This showed that the significant relationship with separation anxiety was significantly stronger than GAD, OCD and physical injury fears ($p < 0.05$), but was not significantly stronger than social anxiety.

In addition, we explored the effects of IQ and verbal language ability. As we only had IQ scores for 41% of the sample, we conducted a Spearman's correlation analyses independently for the group with IQ scores and the group without to see if correlations between sensory reactivity and anxiety differed between groups. The analysis showed that sensory hyperreactivity was significantly related to anxiety in those who were unable to do an IQ test (without IQ score) ($r(30)=0.56$, $p=0.001$), but not in those who were able to conduct an IQ test (with IQ score) ($r(20)=0.38$, $p=0.09$). However, Fisher's z -test confirmed that these correlation coefficients were not significantly different ($p > 0.05$). As for verbal language, correlation analysis showed that sensory hyperreactivity was significantly

related to anxiety in those who had few-to-no words and completed the ADOS-2 Module 1 ($r(28)=0.62$, $p < 0.001$), but not in those who were able to use sentences and completed the ADOS-2 Module 2 ($r(18)=0.29$, $p > 0.05$). However, Fisher's z -test confirmed that these correlation coefficients were not significantly different ($p > 0.05$). Furthermore, independent samples t -test indicated that there were no significant group differences in sensory hyperreactivity or anxiety scores for those with or without an IQ score, or those who did the ADOS-2 Module 1 or Module 2 ($p > 0.05$). Therefore, we did not take IQ or verbal language ability into account in further analyses as the non-significant subgroup findings are likely due to lack of power.

Mediation analyses

As sensory hyperreactivity, IU and total anxiety were intercorrelated, we conducted a series of simple mediation analyses to understand the predictive relationship between these constructs. To achieve this, we tested all possible iterations of the model, including sensory hyperreactivity, IU and total anxiety as predictor, outcome and mediating variables (see Table 4 for mediation analyses). As correlational analyses indicated that these relationships were independent from the effects of age, autism traits, IQ and verbal language level, we did not control for these in the analyses. We applied a Bonferroni's significance correction to control for type 1 error due to multiple analyses ($p \leq 0.008$).

Mediation analysis indicated significant indirect effects between sensory hyperreactivity and anxiety through IU ($B=0.47$, lower limit (LL)=0.28, upper limit (UL)=0.66, $p < 0.001$). A significant indirect effect was also found between sensory hyperreactivity and IU through anxiety ($B=0.42$, LL=0.23, UL=0.72, $p < 0.001$). As the direct effects were not significant for both of these models

Table 4. Mediation analyses between sensory hyperreactivity (SPSI and SAND composite z-score), anxiety (PAS) and intolerance of uncertainty (RULES).

Predictor	Mediator	Outcome	Total effect			Direct effect			Indirect effect								
			B	SE	LL	UL	p	B	SE	LL	UL	p					
Hyper	IU	Anxiety	0.63	0.13	0.35	0.92	< 0.001	0.16	0.12	-0.00	0.40	0.184	0.47	0.12	0.28	0.66	< 0.001
	Anxiety	IU	0.66	0.13	0.38	0.89	< 0.001	0.24	0.12	-0.03	0.55	0.036	0.42	0.12	0.23	0.72	< 0.001
Hyper	IU	Hyper	0.50	0.10	0.32	0.69	< 0.001	0.22	0.17	0.01	0.52	0.184	0.28	0.14	-0.02	0.56	0.040
Anxiety	Hyper	IU	0.79	0.09	0.60	0.99	< 0.001	0.67	0.10	0.41	0.94	< 0.001	0.12	0.06	0.01	0.32	0.055
Anxiety	Anxiety	Hyper	0.53	0.11	0.25	0.75	< 0.001	0.36	0.17	0.02	0.66	0.036	0.18	0.14	-0.03	0.42	0.192
IU	Hyper	Anxiety	0.80	0.09	0.60	0.96	< 0.001	0.71	0.11	0.51	0.90	< 0.001	0.08	0.06	0.00	0.23	0.202

Hyper: sensory hyperreactivity; IU: intolerance of uncertainty; SE: standard error; LL and UL: 95% confidence intervals lower limit and upper limit. Significant relationships at the $p < 0.008$ level after a Bonferroni's correction highlighted in bold.

(Bonferroni's corrected $p > 0.008$ and 95% CIs containing 0), this indicates full mediation (Figure 1). No mediation was indicated when IU was entered as the predictor for sensory hyperreactivity through anxiety, and for anxiety through sensory hyperreactivity, or when anxiety was entered as the predictor for IU through sensory hyperreactivity, or for sensory hyperreactivity through IU, as the indirect effects were not significant (Bonferroni's corrected $p > 0.008$ and 95% CIs containing 0).

Discussion

This study has been the first to explore the relationship between sensory reactivity differences, IU and anxiety in a heterogenous group of preschool-age autistic children. In line with our expectations, we found sensory hyperreactivity, IU and anxiety to be interrelated. In addition, we found that IU fully mediates the relationship between sensory hyperreactivity and anxiety, but that anxiety also fully mediates the relationship between sensory hyperreactivity and IU. We did not find sensory hyporeactivity or seeking to be significantly related to anxiety or IU.

We found, in line with previous findings (Hwang et al., 2019; Neil et al., 2016; Wigham et al., 2015), sensory hyperreactivity to be significantly related to total anxiety in autistic children. In our initial correlation analysis, we did not find sensory hyperreactivity to be differentially related to anxiety subtypes. However, when controlling for autism traits, we found sensory hyperreactivity to be significantly related to separation anxiety. As suggested in our previous work, controlling for AQ may limit measurement or response bias, or variability from trait-symptom overlap (MacLennan et al., 2020). Although existing findings have also shown links between sensory hyperreactivity and separation anxiety (Black et al., 2017), our previous work with older children found sensory hyperreactivity to be related to symptoms associated with physical injury fears and specific phobia (MacLennan et al., 2020). Inconsistencies across studies could be due to the use of different measures or the differing age and developmental level of the participants, as research has found differential prevalence of anxiety conditions across ages of autistic children and adolescents (van Steensel et al., 2011). This may be due to differential conditioning implicated in the development of certain anxiety conditions, such as context conditioning in GAD (Green & Ben-Sasson, 2010). It would be beneficial for future research to explore differential relationship over time and to further understand the association of IU when relationships are differential.

As previous literature has proposed disputing theoretical models that have included sensory hyperreactivity, IU and anxiety, here we tested a series of mediation models. We found that IU fully mediates the relationships between sensory hyperreactivity and anxiety in preschool-age autistic children, and that anxiety fully mediates the relationship




Figure 1. (a) Depicting the relationship between sensory hyperreactivity and anxiety fully mediated by IU and (b) the relationship between sensory hyperreactivity and IU fully mediated by anxiety.

Direct relationships are indicated by a solid line, and indirect relationships are indicated by a dashed line. Significant relationships are indicated by a black line, non-significant with a grey. B coefficients are shown for the indirect effects.

between sensory hyperreactivity and IU. Critically, in line with our predictions, we found that sensory hyperreactivity was the predictor in both significant mediation models. This reflects previous longitudinal research in younger children that found sensory hyperreactivity to unidirectionally predict anxiety symptoms, suggesting that sensory hyperreactivity emerges earlier than anxiety (Green et al., 2012). However, our results also suggest that IU is an important early factor in the development and maintenance of anxiety in autism. Aversive sensory experiences can be unpredictable and not always specific to an object or situation, which is associated with greater avoidance and anxious behavioural responses, and associated amygdala activation (Green et al., 2015; Herry et al., 2007). Altered sensory interpretation in autism can enhance the perceived unpredictability of sensory input (Fiser et al., 2010; Pellicano & Burr, 2012). As IU is associated with alerting in attentional networks, individuals with higher IU may be more hypervigilant and biased to process uncertain information as threatening (Dugas et al., 2005; Fergus & Carleton, 2016). Thus, IU and anxiety may reciprocally be maintained in response to sensory hyperreactivity. Due to the constraints of cross-sectional mediation models, next steps in research would be to longitudinally test these mediating relationships.

As our results support that sensory hyperreactivity is a predictive factor for anxiety early in development, this could be an important target for early interventions. Sensory integration-based approaches are widely used and are deemed to be effective in targeting early sensory reactivity difficulties (Schaaf et al., 2017; Schoen et al., 2019), but little is known about how these impact anxiety development. Alternatively, Cognitive Behavioural Therapy (CBT) could be adapted to target sensory hyperreactivity (Edgington et al., 2016), as CBT has shown efficacy in young autistic children (Driscoll et al., 2020). However, IU may also be an important transdiagnostic mechanism to target in intervention (Hallett et al., 2021; Rodgers et al., 2017). There is evidence that targeting IU in clinical

interventions effectively reduces anxiety in non-autistic cohorts (Boswell et al., 2013; Buhr & Dugas, 2006; Dugas et al., 2010; McEvoy & Mahoney, 2012). This has recently been extended into autism research, with the development of a manualised parent group-based intervention programme, such as Predictive Parenting (Hallett et al., 2021) and Coping with Uncertainty in Everyday Situations (CUES[®]; Rodgers et al., 2018, 2017). These interventions aim to enable children to become more tolerant of uncertainty and have shown promise in their efficacy.

In contrast to previous research (Glod et al., 2019; Hwang et al., 2019; MacLennan et al., 2020), we did not find sensory hyporeactivity to relate to anxiety or IU in this study. This may be due to the varying measures and age groups across studies. Alternatively, sensory hyporeactivity may be more related to depressive conditions, characterised by suppressed arousal, rather than anxiety (Lane, 2002; Neal et al., 2002; Pfeiffer et al., 2005). Or it may be more related to other autistic traits, such as communication differences, emotional, cognitive and behavioural differences, or RRBs (Glod et al., 2015; South & Rodgers, 2017; Wigham et al., 2015). As for sensory seeking, we did not find it to be significantly related to anxiety in autistic children, which reflects previous findings (MacLennan et al., 2020). Sensory seeking may be indirectly linked to anxiety or interlinked with sensory hyperreactivity and hyporeactivity, such as being a stimulatory strategy in response to sensory hyporeactivity (Lidstone et al., 2014; Liss et al., 2006). Future directions should aim to further understand how sensory hyporeactivity and seeking are related to anxiety.

There is a pressing need for research to include participants with few-to-no words or ID (Russell et al., 2019), thus, our heterogeneous sample is a key strength of the study. However, our relatively small sample size may have impacted the power of our subgroup analysis, as well as relationships with anxiety subtypes. In addition, there are a few suitable objective assessments for IU and anxiety,

especially for this demographic, and therefore, we had to rely on parent-reports. Although total anxiety scores have been found to be reliable in autistic populations, subscales for anxiety on parent-reports may need to be interpreted with caution in autistic children, especially those with co-occurring ID and who speak few-to-no words (Kerns et al., 2020; Toscano et al., 2020). Understanding the risk factors for traditional anxiety symptomology has clinical significance for existing anxiety interventions; however, it is also important that distinct anxiety symptoms in autistic children are recognised (Kerns et al., 2020). Therefore, there is a need for the development of autism-specific, measures that can be used to screen for early anxiety symptomology in young autistic children. But also, further research examining anxiety in autistic subgroups, such as those with few-to-no words or ID, is needed.

In conclusion, this study is the first to explore the relationship between sensory reactivity differences, anxiety subtypes and IU in preschool-age autistic children. Our findings indicate that sensory hyperreactivity, IU and anxiety are interrelated in preschool-age autistic children, and that sensory hyperreactivity may be specifically related to separation anxiety when taking broader autism traits into account. In addition, they suggest sensory hyperreactivity is an early predictor of anxiety and IU, which may both be reciprocally mediating factors in the maintenance of anxiety and IU in autism. The results of this study have important clinical implications for early anxiety interventions in preschool-age autistic children. However, there is still an imperative need to develop objective assessments, especially for IU and anxiety, for preschool-age autistic children, and those who speak few-to-no words. This is fundamental for research and clinical services that aim to support autistic individuals.

Acknowledgements

The authors specially thank all the families who participated. The authors also thank Francesca Englezou, Charlotte Daniels and Khadija Kapadia of the University of Reading for assisting with data collection, and the Centre for Autism at the University of Reading, Autism Berkshire and NHS Berkshire Healthcare CAMHS for helping with recruitment.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship and/or publication of this article: The authors are funded by MQ and Autistica.

ORCID iD

Keren MacLennan  <https://orcid.org/0000-0002-3148-3450>

Note

1. The terminology and language used in this article are important for the reduction of stigma (Gernsbacher, 2017). It is informed by preferred terms that have been outlined by the autistic community and their families in research (Kenny et al., 2016), as well as feedback from the families who took part in this study.

References

- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders (DSM-5)*. American Psychiatric Publishing.
- Auyeung, B., Baron-Cohen, S., Wheelwright, S., & Allison, C. (2008). The Autism Spectrum Quotient: Children's Version (AQ-Child). *Journal of Autism and Developmental Disorders*, 38(7), 1230–1240. <https://doi.org/10.1007/s10803-007-0504-z>
- Ben-Sasson, A., Cermak, S. A., Orsmond, G. I., Tager-Flusberg, H., Carter, A. S., Kadlec, M. B., & Dunn, W. (2007). Extreme sensory modulation behaviors in toddlers with autism spectrum disorders. *The American Journal of Occupational Therapy: Official Publication of the American Occupational Therapy Association*, 61(5), 584–592. <https://doi.org/10.5014/ajot.61.5.584>
- Ben-Sasson, A., Cermak, S. A., Orsmond, G. I., Tager-Flusberg, H., Kadlec, M. B., & Carter, A. S. (2008). Sensory clusters of toddlers with autism spectrum disorders: Differences in affective symptoms. *Journal of Child Psychology and Psychiatry*, 49(8), 817–825. <https://doi.org/10.1111/j.1469-7610.2008.01899.x>
- Black, K. R., Stevenson, R. A., Segers, M., Ncube, B. L., Sun, S. Z., Philipp-Muller, A., Bebkko, J. M., Barensse, M. D., & Ferber, S. (2017). Linking anxiety and insistence on sameness in autistic children: The role of sensory hypersensitivity. *Journal of Autism and Developmental Disorders*, 47(8), 2459–2470. <https://doi.org/10.1007/s10803-017-3161-x>
- Boswell, J. F., Thompson-Hollands, J., Farchione, T. J., & Barlow, D. H. (2013). Intolerance of uncertainty: A common factor in the treatment of emotional disorders. *Journal of Clinical Psychology*, 69(6), 630–645. <https://doi.org/10.1002/jclp.21965>
- Boulter, C., Freeston, M., South, M., & Rodgers, J. (2014). Intolerance of uncertainty as a framework for understanding anxiety in children and adolescents with autism spectrum disorders. *Journal of Autism and Developmental Disorders*, 44(6), 1391–1402. <https://doi.org/10.1007/s10803-013-2001-x>
- Buhr, K., & Dugas, M. J. (2006). Investigating the construct validity of intolerance of uncertainty and its unique relationship with worry. *Journal of Anxiety Disorders*, 20(2), 222–236. <https://doi.org/10.1016/j.janxdis.2004.12.004>
- Carleton, R. N., Mulvogue, M. K., Thibodeau, M. A., McCabe, R. E., Antony, M. M., & Asmundson, G. J. G. (2012). Increasingly certain about uncertainty: Intolerance of uncertainty across anxiety and depression. *Journal of Anxiety Disorders*, 26(3), 468–479. <https://doi.org/10.1016/j.janxdis.2012.01.011>
- Carpenter, K. L. H., Baranek, G. T., Copeland, W. E., Compton, S., Zucker, N., Dawson, G., & Egger, H. L. (2019). Sensory over-responsivity: An early risk factor for anxi-

- ety and behavioral challenges in young children. *Journal of Abnormal Child Psychology*, 47, 1075–1088. <https://doi.org/10.1007/s10802-018-0502-y>
- Chamberlain, P. D., Rodgers, J., Crowley, M. J., White, S. E., Freeston, M. H., & South, M. (2013). A potentiated startle study of uncertainty and contextual anxiety in adolescents diagnosed with autism spectrum disorder. *Molecular Autism*, 4(1), 31. <https://doi.org/10.1186/2040-2392-4-31>
- Driscoll, K., Schonberg, M., Stark, M. F., Carter, A. S., & Hirshfeld-Becker, D. (2020). Family-centered cognitive behavioral therapy for anxiety in very young children with autism spectrum disorder. *Journal of Autism and Developmental Disorders*, 50(11), 3905–3920. <https://doi.org/10.1007/s10803-020-04446-y>
- Dugas, M. J., Brillon, P., Savard, P., Turcotte, J., Gaudet, A., Ladouceur, R., Leblanc, R., & Gervais, N. J. (2010). A randomized clinical trial of cognitive-behavioral therapy and applied relaxation for adults with generalized anxiety disorder. *Behavior Therapy*, 41(1), 46–58. <https://doi.org/10.1016/j.beth.2008.12.004>
- Dugas, M. J., Gagnon, F., Ladouceur, R., & Freeston, M. H. (1998). Generalized anxiety disorder: A preliminary test of a conceptual model. *Behaviour Research and Therapy*, 36(2), 215–226. [https://doi.org/10.1016/S0005-7967\(97\)00070-3](https://doi.org/10.1016/S0005-7967(97)00070-3)
- Dugas, M. J., Hedayati, M., Karavidas, A., Buhr, K., Francis, K., & Phillips, N. A. (2005). Intolerance of uncertainty and information processing: Evidence of biased recall and interpretations. *Cognitive Therapy and Research*, 29(1), 57–70. <https://doi.org/10.1007/s10608-005-1648-9>
- Dunn, W. (1999). *Sensory profile* (Vol. 555). Psychological Corporation.
- Edgington, L., Hill, V., & Pellicano, E. (2016). The design and implementation of a CBT-based intervention for sensory processing difficulties in adolescents on the autism spectrum. *Research in Developmental Disabilities*, 59, 221–233. <https://doi.org/10.1016/j.ridd.2016.09.004>
- Edwards, S. L., Rapee, R. M., Kennedy, S. J., & Spence, S. H. (2010). The assessment of anxiety symptoms in preschool-aged children: The revised Preschool Anxiety Scale. *Journal of Clinical Child & Adolescent Psychology*, 39(3), 400–409. <https://doi.org/10.1080/15374411003691701>
- Elliott, C. D., Murray, D. J., & Pearson, L. S. (1983). *British ability scales*. NFER-Nelson.
- Fergus, T. A., & Carleton, R. N. (2016). Intolerance of uncertainty and attentional networks: Unique associations with alerting. *Journal of Anxiety Disorders*, 41, 59–64. <https://doi.org/10.1016/j.janxdis.2016.03.010>
- Fiser, J., Berkes, P., Orbán, G., & Lengyel, M. (2010). Statistically optimal perception and learning: From behavior to neural representations. *Trends in Cognitive Sciences*, 14(3), 119–130. <https://doi.org/10.1016/j.tics.2010.01.003>
- Freeston, M. H., Rhéaume, J., Letarte, H., Dugas, M. J., & Ladouceur, R. (1994). Why do people worry? *Personality and Individual Differences*, 17(6), 791–802. [https://doi.org/10.1016/0191-8869\(94\)90048-5](https://doi.org/10.1016/0191-8869(94)90048-5)
- Gernsbacher, M. A. (2017). Editorial perspective: The use of person-first language in scholarly writing may accentuate stigma. *Journal of Child Psychology and Psychiatry*, 58(7), 859–861. <https://doi.org/10.1111/jcpp.12706>
- Glod, M., Riby, D. M., Honey, E., & Rodgers, J. (2015). Psychological correlates of sensory processing patterns in individuals with autism spectrum disorder: A systematic review. *Review Journal of Autism and Developmental Disorders*, 2(2), 199–221. <https://doi.org/10.1007/s40489-015-0047-8>
- Glod, M., Riby, D. M., & Rodgers, J. (2019). Short report: Relationships between sensory processing, repetitive behaviors, anxiety, and intolerance of uncertainty in autism spectrum disorder and Williams syndrome. *Autism Research*, 12(5), 759–765. <https://doi.org/10.1002/aur.2096>
- Green, S. A., & Ben-Sasson, A. (2010). Anxiety disorders and sensory over-responsivity in children with autism spectrum disorders: Is there a causal relationship? *Journal of Autism and Developmental Disorders*, 40(12), 1495–1504. <https://doi.org/10.1007/s10803-010-1007-x>
- Green, S. A., Ben-Sasson, A., Soto, T. W., & Carter, A. S. (2012). Anxiety and sensory over-responsivity in toddlers with autism spectrum disorders: Bidirectional effects across time. *Journal of Autism and Developmental Disorders*, 42(6), 1112–1119. <https://doi.org/10.1007/s10803-011-1361-3>
- Green, S. A., Hernandez, L., Tottenham, N., Krasileva, K., Bookheimer, S. Y., & Dapretto, M. (2015). Neurobiology of sensory overresponsivity in youth with autism spectrum disorders. *JAMA Psychiatry*, 72(8), 778–786. <https://doi.org/10.1001/jamapsychiatry.2015.0737>
- Hallett, V., Mueller, J., Breese, L., Hollett, M., Beresford, B., Irvine, A., Pickles, A., Slonims, V., Scott, S., Charman, T., & Simonoff, E. (2021). Introducing ‘predictive parenting’: A feasibility study of a new group parenting intervention targeting emotional and behavioral difficulties in children with autism spectrum disorder. *Journal of Autism and Developmental Disorders*, 51(1), 323–333. <https://doi.org/10.1007/s10803-020-04442-2>
- Herry, C., Bach, D. R., Esposito, F., Di Salle, F., Perrig, W. J., Scheffler, K., Luthi, A., & Seifritz, E. (2007). Processing of temporal unpredictability in human and animal amygdala. *Journal of Neuroscience*, 27(22), 5958–5966. <https://doi.org/10.1523/JNEUROSCI.5218-06.2007>
- Hodgson, A. R., Freeston, M. H., Honey, E., & Rodgers, J. (2017). Facing the unknown: Intolerance of uncertainty in children with autism spectrum disorder. *Journal of Applied Research in Intellectual Disabilities*, 30(2), 336–344. <https://doi.org/10.1111/jar.12245>
- Hwang, Y. I., (Jane) Arnold, S., Srasuebkul, P., & Trollor, J. (2020). Understanding anxiety in adults on the autism spectrum: An investigation of its relationship with intolerance of uncertainty, sensory sensitivities and repetitive behaviours. *Autism*, 24, 411–422. <https://doi.org/10.1177/1362361319868907>
- IBM Corp. (2020). *IBM SPSS Statistics for Mac* (27.0) [Computer software].
- JASP team. (2020). *JASP (Version 0.14)* [Computer software]. <https://jasp-stats.org/>
- Kenny, L., Hattersley, C., Molins, B., Buckley, C., Povey, C., & Pellicano, E. (2016). Which terms should be used to describe autism? Perspectives from the UK autism community. *Autism*, 20(4), 442–462. <https://doi.org/10.1177/1362361315588200>
- Kerns, C. M., & Kendall, P. C. (2012). The presentation and classification of anxiety in autism spectrum disorder. *Clinical*

- Psychology: Science and Practice*, 19(4), 323–347. <https://doi.org/10.1111/cpsp.12009>
- Kerns, C. M., Kendall, P. C., Berry, L., Souders, M. C., Franklin, M. E., Schultz, R. T., Miller, J., & Herrington, J. (2014). Traditional and atypical presentations of anxiety in youth with autism spectrum disorder. *Journal of Autism and Developmental Disorders*, 44(11), 2851–2861. <https://doi.org/10.1007/s10803-014-2141-7>
- Kerns, C. M., Renno, P., Kendall, P. C., Wood, J. J., & Storch, E. A. (2017). Anxiety disorders interview schedule–autism addendum: Reliability and validity in children with autism spectrum disorder. *Journal of Clinical Child & Adolescent Psychology*, 46(1), 88–100. <https://doi.org/10.1080/15374416.2016.1233501>
- Kerns, C. M., Winder-Patel, B., Iosif, A. M., Nordahl, C. W., Heath, B., Solomon, M., & Amaral, D. G. (2020). Clinically significant anxiety in children with autism spectrum disorder and varied intellectual functioning. *Journal of Clinical Child & Adolescent Psychology*. Advance online publication. <https://doi.org/10.1080/15374416.2019.1703712>
- Kim, S. H., & Lord, C. (2012). Combining information from multiple sources for the diagnosis of autism spectrum disorders for toddlers and young preschoolers from 12 to 47 months of age: Combining information for the diagnosis of ASD for toddlers and young preschoolers. *Journal of Child Psychology and Psychiatry*, 53(2), 143–151. <https://doi.org/10.1111/j.1469-7610.2011.02458.x>
- Lane, S. J. (2002). Sensory modulation. In A. C. Bundy, S. J. Lane, & E. A. Murray (Eds.), *Sensory integration: Theory and practice* (2nd ed., pp. 101–122). F. A. Davis.
- Leekam, S. R., Nieto, C., Libby, S. J., Wing, L., & Gould, J. (2007). Describing the sensory abnormalities of children and adults with autism. *Journal of Autism and Developmental Disorders*, 37(5), 894–910. <https://doi.org/10.1007/s10803-006-0218-7>
- Levit-Binnun, N., Szepsenwol, O., Stern-Ellran, K., & Engel-Yeger, B. (2014). The relationship between sensory responsiveness profiles, attachment orientations, and anxiety symptoms: Sensory responsiveness, attachment, and anxiety. *Australian Journal of Psychology*, 66(4), 233–240. <https://doi.org/10.1111/ajpy.12064>
- Lidstone, J., Uljarević, M., Sullivan, J., Rodgers, J., McConachie, H., Freeston, M., Le Couteur, A., Prior, M., & Leekam, S. (2014). Relations among restricted and repetitive behaviors, anxiety and sensory features in children with autism spectrum disorders. *Research in Autism Spectrum Disorders*, 8(2), 82–92. <https://doi.org/10.1016/j.rasd.2013.10.001>
- Liss, M., Saulnier, C., Fein, D., & Kinsbourne, M. (2006). Sensory and attention abnormalities in autistic spectrum disorders. *Autism*, 10(2), 155–172. <https://doi.org/10.1177/13623613060062021>
- Lord, C., Rutter, M., DiLavore, P., Risi, S., Gotham, K., & Bishop, S. (2012). *Autism diagnostic observation schedule* (2nd ed.). Western Psychological Services.
- MacLennan, K., Roach, L., & Tavassoli, T. (2020). The relationship between sensory reactivity differences and anxiety subtypes in autistic children. *Autism Research*, 13(5), 785–795. <https://doi.org/10.1002/aur.2259>
- McEvoy, P. M., & Mahoney, A. E. J. (2012). To be sure, to be sure: Intolerance of uncertainty mediates symptoms of various anxiety disorders and depression. *Behavior Therapy*, 43(3), 533–545. <https://doi.org/10.1016/j.beth.2011.02.007>
- Miller, L. J., Anzalone, M. E., Lane, S. J., Cermak, S. A., & Osten, E. T. (2007). Concept evolution in sensory integration: A proposed nosology for diagnosis. *The American Journal of Occupational Therapy: Official Publication of the American Occupational Therapy Association*, 61(2), 135–140. <https://doi.org/10.5014/ajot.61.2.135>
- Miller, L. J., Nielsen, D. M., & Schoen, S. A. (2012). Attention deficit hyperactivity disorder and sensory modulation disorder: A comparison of behavior and physiology. *Research in Developmental Disabilities*, 33(3), 804–818. <https://doi.org/10.1016/j.ridd.2011.12.005>
- Neal, J. A., Edelmann, R. J., & Glachan, M. (2002). Behavioural inhibition and symptoms of anxiety and depression: Is there a specific relationship with social phobia? *British Journal of Clinical Psychology*, 41(4), 361–374. <https://doi.org/10.1348/014466502760387489>
- Neil, L., Green, D., & Pellicano, E. (2017). The psychometric properties of a new measure of sensory behaviors in autistic children. *Journal of Autism and Developmental Disorders*, 47(4), 1261–1268. <https://doi.org/10.1007/s10803-016-3018-8>
- Neil, L., Olsson, N. C., & Pellicano, E. (2016). The relationship between intolerance of uncertainty, sensory sensitivities, and anxiety in autistic and typically developing children. *Journal of Autism and Developmental Disorders*, 46(6), 1962–1973. <https://doi.org/10.1007/s10803-016-2721-9>
- Pellicano, E., & Burr, D. (2012). When the world becomes ‘too real’: A Bayesian explanation of autistic perception. *Trends in Cognitive Sciences*, 16(10), 504–510. <https://doi.org/10.1016/j.tics.2012.08.009>
- Pfeiffer, B., Kinnealey, M., Reed, C., & Herzberg, G. (2005). Sensory modulation and affective disorders in children and adolescents with Asperger’s disorder. *The American Journal of Occupational Therapy: Official Publication of the American Occupational Therapy Association*, 59(3), 335–345.
- Pickard, H., Hirsch, C., Simonoff, E., & Happé, F. (2020). Exploring the cognitive, emotional and sensory correlates of social anxiety in autistic and neurotypical adolescents. *Journal of Child Psychology and Psychiatry*, 61, 1317–1327. <https://doi.org/10.1111/jcpp.13214>
- Rapee, R. M., Kennedy, S. J., Ingram, M., Edwards, S. L., & Sweeney, L. (2010). Altering the trajectory of anxiety in at-risk young children. *American Journal of Psychiatry*, 167(12), 1518–1525. <https://doi.org/10.1176/appi.ajp.2010.09111619>
- Rodgers, J., Herrema, R., Honey, E., & Freeston, M. (2018). Towards a treatment for intolerance of uncertainty for autistic adults: A single case experimental design study. *Journal of Autism and Developmental Disorders*, 48(8), 2832–2845. <https://doi.org/10.1007/s10803-018-3550-9>
- Rodgers, J., Hodgson, A., Shields, K., Wright, C., Honey, E., & Freeston, M. (2017). towards a treatment for intolerance of uncertainty in young people with autism spectrum disorder: Development of the Coping with Uncertainty in Everyday Situations (CUES[®]) Programme. *Journal of Autism and Developmental Disorders*, 47(12), 3959–3966. <https://doi.org/10.1007/s10803-016-2924-0>
- Russell, G., Mandy, W., Elliott, D., White, R., Pittwood, T., & Ford, T. (2019). Selection bias on intellectual ability in

- autism research: A cross-sectional review and meta-analysis. *Molecular Autism*, 10(1), 9. <https://doi.org/10.1186/s13229-019-0260-x>
- Sanchez, A. L., Cornacchio, D., Chou, T., Leyfer, O., Coxe, S., Pincus, D., & Comer, J. S. (2017). Development of a scale to evaluate young children's responses to uncertainty and low environmental structure. *Journal of Anxiety Disorders*, 45, 17–23. <https://doi.org/10.1016/j.janxdis.2016.11.006>
- Schaaf, R. C., Dumont, R. L., Arbesman, M., & May-Benson, T. A. (2017). Efficacy of occupational therapy using Ayres Sensory Integration®: A systematic review. *The American Journal of Occupational Therapy: Official Publication of the American Occupational Therapy Association*, 72(1), 7201190010p1–7201190010p10. <https://doi.org/10.5014/ajot.2018.028431>
- Schoen, S. A., Lane, S. J., Mailloux, Z., May-Benson, T., Parham, L. D., Smith Roley, S., & Schaaf, R. C. (2019). A systematic review of Ayres sensory integration intervention for children with autism. *Autism Research*, 12(1), 6–19. <https://doi.org/10.1002/aur.2046>
- Schoen, S. A., Miller, L. J., & Sullivan, J. (2017). The development and psychometric properties of the Sensory Processing Scale Inventory: A report measure of sensory modulation. *Journal of Intellectual & Developmental Disability*, 42(1), 12–21. <https://doi.org/10.3109/13668250.2016.1195490>
- Simonoff, E., Pickles, A., Charman, T., Chandler, S., Loucas, T., & Baird, G. (2008). Psychiatric disorders in children with autism spectrum disorders: Prevalence, comorbidity, and associated factors in a population-derived sample. *Journal of the American Academy of Child & Adolescent Psychiatry*, 47(8), 921–929. <https://doi.org/10.1097/CHI.0b013e318179964f>
- Siper, P. M., Kolevzon, A., Wang, A. T., Buxbaum, J. D., & Tavassoli, T. (2017). A clinician-administered observation and corresponding caregiver interview capturing DSM-5 sensory reactivity symptoms in children with ASD: Sensory assessment for neurodevelopmental disorders. *Autism Research*, 10(6), 1133–1140. <https://doi.org/10.1002/aur.1750>
- Song, M.-K., Lin, F.-C., Ward, S. E., & Fine, J. P. (2013). Composite variables: When and how. *Nursing Research*, 62(1), 45–49. <https://doi.org/10.1097/NNR.0b013e3182741948>
- South, M., & Rodgers, J. (2017). Sensory, emotional and cognitive contributions to anxiety in autism spectrum disorders. *Frontiers in Human Neuroscience*, 11, 20. <https://doi.org/10.3389/fnhum.2017.00020>
- Spain, D., Sin, J., Linder, K. B., McMahon, J., & Happé, F. (2018). Social anxiety in autism spectrum disorder: A systematic review. *Research in Autism Spectrum Disorders*, 52, 51–68. <https://doi.org/10.1016/j.rasd.2018.04.007>
- Spence, S. H. (1998). A measure of anxiety symptoms among children. *Behaviour Research and Therapy*, 36(5), 545–566. [https://doi.org/10.1016/S0005-7967\(98\)00034-5](https://doi.org/10.1016/S0005-7967(98)00034-5)
- Tavassoli, T., Bellesheim, K., Siper, P. M., Wang, A. T., Halpern, D., Gorenstein, M., Grodberg, D., Kolevzon, A., & Buxbaum, J. D. (2016). Measuring sensory reactivity in autism spectrum disorder: Application and simplification of a clinician-administered sensory observation scale. *Journal of Autism and Developmental Disorders*, 46(1), 287–293. <https://doi.org/10.1007/s10803-015-2578-3>
- Toscano, R., Baillie, A. J., Lynham, H. J., Kelly, A., Kidd, T., & Hudson, J. L. (2020). Assessment of anxiety in children and adolescents: A comparative study on the validity and reliability of the Spence Children's Anxiety Scale in children and adolescents with anxiety and Autism Spectrum Disorder. *Journal of Affective Disorders*, 260, 569–576. <https://doi.org/10.1016/j.jad.2019.09.055>
- Uljarević, M., Lane, A., Kelly, A., & Leekam, S. (2016). Sensory subtypes and anxiety in older children and adolescents with autism spectrum disorder: Sensory subtypes and anxiety in autism. *Autism Research*, 9(10), 1073–1078. <https://doi.org/10.1002/aur.1602>
- van Steensel, F. J. A., Bögels, S. M., & Perrin, S. (2011). Anxiety disorders in children and adolescents with autistic spectrum disorders: A meta-analysis. *Clinical Child and Family Psychology Review*, 14(3), 302–317. <https://doi.org/10.1007/s10567-011-0097-0>
- Vasa, R. A., Keefer, A., McDonald, R. G., Hunsche, M. C., & Kerns, C. M. (2020). A scoping review of anxiety in young children with autism spectrum disorder. *Autism Research*, 13, 2038–2057. <https://doi.org/10.1002/aur.2395>
- Vasa, R. A., Kreiser, N. L., Keefer, A., Singh, V., & Mostofsky, S. H. (2018). Relationships between autism spectrum disorder and intolerance of uncertainty: Relationships between IU and ASD. *Autism Research*, 11(4), 636–644. <https://doi.org/10.1002/aur.1916>
- Wechsler, D., & Naglieri, J. A. (2006). *Wechsler nonverbal scale of ability: WNV*. PsychCorp.
- White, S. W., Oswald, D., Ollendick, T., & Scahill, L. (2009). Anxiety in children and adolescents with autism spectrum disorders. *Clinical Psychology Review*, 29(3), 216–229. <https://doi.org/10.1016/j.cpr.2009.01.003>
- Wigham, S., Rodgers, J., South, M., McConachie, H., & Freeston, M. (2015). The interplay between sensory processing abnormalities, intolerance of uncertainty, anxiety and restricted and repetitive behaviours in autism spectrum disorder. *Journal of Autism and Developmental Disorders*, 45(4), 943–952. <https://doi.org/10.1007/s10803-014-2248-x>
- Yap, M. B. H., Morgan, A. J., Cairns, K., Jorm, A. F., Hetrick, S. E., & Merry, S. (2016). Parents in prevention: A meta-analysis of randomized controlled trials of parenting interventions to prevent internalizing problems in children from birth to age 18. *Clinical Psychology Review*, 50, 138–158. <https://doi.org/10.1016/j.cpr.2016.10.003>