

Patrolling the boundaries of social domains: neural activations to violations of expectations for romantic and work relationships

Article

Published Version

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Open Access

Bland, A. R., Zahn, R. R., Elliott, R., Taylor, J. R. and Hill, J. (2021) Patrolling the boundaries of social domains: neural activations to violations of expectations for romantic and work relationships. Social neuroscience, 16 (5). pp. 513-521. ISSN 1747-0927 doi: 10.1080/17470919.2021.1953134 Available at https://centaur.reading.ac.uk/99353/

It is advisable to refer to the publisher's version if you intend to cite from the work. See <u>Guidance on citing</u>.

To link to this article DOI: http://dx.doi.org/10.1080/17470919.2021.1953134

Publisher: Taylor & Francis

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the <u>End User Agreement</u>.



www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading Reading's research outputs online



Social Neuroscience Neuroscience

ne 9 Itsues 1-2 February-April 201

Social

ISSN: (Print) (Online) Journal homepage: https://www.tandfonline.com/loi/psns20

Patrolling the boundaries of social domains: Neural activations to violations of expectations for romantic and work relationships

A. R. Bland, R. Zahn, R. Elliott, J. R. Taylor & J. Hill

To cite this article: A. R. Bland, R. Zahn, R. Elliott, J. R. Taylor & J. Hill (2021): Patrolling the boundaries of social domains: Neural activations to violations of expectations for romantic and work relationships, Social Neuroscience, DOI: 10.1080/17470919.2021.1953134

To link to this article: https://doi.org/10.1080/17470919.2021.1953134

© 2021 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



0

Published online: 13 Jul 2021.

_	
	P
-	

Submit your article to this journal 🖸

Article views: 52



View related articles

View Crossmark data 🗹

Patrolling the boundaries of social domains: Neural activations to violations of expectations for romantic and work relationships

A. R. Bland ^[], R. Zahn^b, R. Elliott ^[], J. R. Taylor^d and J. Hill^e

^aDepartment of Psychology, Manchester Metropolitan University, Manchester, UK; ^bCentre for Affective Disorders, Institute of Psychiatry, Psychology and Neuroscience, Kings College London, UK; ^cNeuroscience and Psychiatry Unit, University of Manchester, UK; ^dDivision of Neuroscience and Experimental Psychology, School of Biological Sciences, University of Manchester, UK; ^eSchool of Psychology and Clinical Language Sciences, University of Reading, UK

ABSTRACT

According to the social domains hypothesis, we reduce the information-processing demands of complex social cues by classifying them into a limited number of domains, each with distinct sets of expectations. This requires rapid identification of violations of the boundaries between social domains. We hypothesized that these violations are likely to be associated with neural activation of the salience system. Using fMRI we compared responses of 20 adults to expected and unexpected everyday social scenarios in personal and work interactions. The vignettes exemplified different kinds of scenarios presented in the work setting, i.e., task-focused scenarios which are expected at work and scenarios with a personal focus, which are unexpected at work. The key contrast between task and personal focussed scenarios presented in the work setting was associated with fronto-insular activation. Perceived inappropriateness of the unexpected scenarios, and shorter response time to judgment of inappropriateness were also associated with fronto-insular activation, after controlling for unpleasantness. This study indicates specific neural responses to violations of expectations in different social situations. Our findings suggest that the fronto-insular region is implicated in rapid detection of behaviors that cross the boundaries of social domains, which are hypothesized to be necessary for efficient social information processing.

Introduction

Managing and respecting boundaries in social situations is a crucial skill for forming and maintaining different kinds of human relationships. For instance, a workrelated question asked by a colleague falls within the expected boundaries of the social situation, while a deeply personal question is likely to feel inappropriate and uncomfortable. This, we have proposed, is a particular instance of the general way in which humans classify social experiences into social domains, such as work and romantic settings, in order to guide the interpretation of social cues and the range of possible actions in each domain (Hill et al., 2010). It reflects the broader human capability to select from, and summarize, complex sensory inputs in order to underpin appropriate and timely actions (Bolton and Hill 2004, Gigerenzer, 2008, Kahneman et al. 1982). Social domains provide a rapid and reasonably accurate basis for social interactions, provided that others' social behaviors are monitored to detect whether they conform to the expectations of the domain, or whether they cross the boundary. Behaviors that cross the boundary such as a colleague interacting **ARTICLE HISTORY**

OPEN ACCESS Check for updates

Received 27 January 2021 Revised 1 June 2021 Published online 24 july 2021

Routledge

Faylor & Francis Group

KEYWORDS

-fMRI; fronto-insular; salience; socialinappropriateness; socialinteraction

in a very personal way, will give rise to uncertainty in how to respond and will require greater attentional and information processing resources while behaviors are evaluated and the identity of the domain reappraised. We hypothesized that this ability will be underpinned by neural structures implicated in evaluating salience and by neurones found in species with more complex social organizations (Allman et al., 2010). In this study we tested, using fMRI, the prediction that social stimuli that cross social domains boundaries will be associated with amygdala, dorsal anterior cingulate, and frontoinsular activation.

It seems likely that there is overlap between brain regions implicated in responding to salient social and nonsocial events. Indeed, some studies have found that those responses to social stimuli involve the same salient network structures as nonsocial stimuli (Luo et al., 2014). The potential importance of the salience network to accurate social responding was illustrated in a study of adolescents, where unexpected mismatches between facial emotional expressions were associated with correct AIC and dACC activations specifically in early

CONTACT A. R. Bland a.bland@mmu.ac.uk Department of Psychology, Manchester Metropolitan University, Brooks Building, 53 Bonsall St, Hulme, Manchester M15 6GXUK

© 2021 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (http://creativecommons.org/licenses/by-ncnd/4.0/), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.

adolescence, a period of increasing social sensitivity (Rosen et al., 2017). Furthermore, better social functioning was associated with the right AIC and the right DACC activation. However, social salience may not be unitary, with different aspects of social stimuli activating different brain regions. For example, in a study of responses to emotionally arousing pictures, amygdala activation was associated with participants' ratings of their emotional intensity, but not their ratings of how much they related to the pictures (Phan et al., 2004). Personal relatedness, however, was associated with dorsal medial frontal and insula activation. Furthermore, social stimuli used in experiments rarely refer to day-to-day social experiences and so we may need to look for more indirect evidence in generating hypotheses for social salience. One likely informative route is to identify regions and locations of neurons that are confined to animals with complex social organizations or even distinct to humans. In relation to the insula, humans and apes can be distinguished from monkeys and small mammals by the presence of Von Economo Neurones (VEN) in the ventral anterior (agranular) insula, with a particular abundance of these neurons in humans (Rose, 1928). Comparative studies have demonstrated VEN phylogenetic restriction to humans and other large-brained mammals with complex sociality, including great apes, cetaceans, and elephants (Allman et al., 2010; Butti et al., 2009; Hakeem et al., 2009; Hof & Van der Gucht, 2007; Nimchinsky et al., 1999). Contemporary functional imaging, lesion-deficit, and anatomical studies link the VEN-rich regions of the anterior cingulate and fronto-insular cortices to a host of social-emotional functions (Craig, 2010; Critchley 2005; Heimer & Van Hoesen, 2006; Seeley et al., 2007; Williamson & Allman, 2011). In the light of this evidence, the fronto-insular region as well as the anterior insula may be a candidate for processing social salience.

The salience of many social stimuli depends not only on their physical characteristics, but also on their "priority" in relation to high-level representations such as goals or expectations (Ptak, 2012). These expectations may be formulated as more or less explicit rules for interactions, with violations of the rules likely to activate the salience system. Neural correlates of responses to violations of social norms have been investigated using monetary exchange paradigms, where norms are established between "investors" and "trustees" based on offers that are likely to benefit both participants. Offers by investors that make mutual benefits less likely therefore violate the expected norm. Detection of this violation by trustees is essential to their ability to find ways of restoring cooperation. Several studies have demonstrated AIC activation in trustees in response to these violations by investors, suggesting its key role in detecting violations of social norms (Rilling & Sanfey, 2011; Spitzer et al., 2007). The question then arises whether the same mechanism is employed in more naturalistic social exchanges.

We have argued that information processing in dayto-day social exchanges is made manageable in humans by our ability to partition social interactions into different social domains. These "algorithms of social life" provide implicit rules that guide the interpretation of social events and social responding (Bugental, 2000; I. Hill et al., 2010; J. Hill et al., 2008). For example, the implicit rules of an educational or work environment include that each person has a task focused role and that expressions of emotions can generally be interpreted in relation to that role. By contrast, according to the implicit rules of a romantic relationship emotions need to be interpreted in relation to a much broader set of possibilities. These domains are efficient vehicles for social interaction, ensuring, for example, that, at work, people can cooperate efficiently immediately after meeting, while romantic relationships can provide emotional resources not available at work but rely on timeconsuming building of relationships. Social domain functioning will only be efficient if domains are implemented jointly by social participants. This is ensured to some degree by social structures such as employment contracts and marriage, but these are unlikely to guarantee shared moment-to-moment implementation of domains. This, we suggest, is maintained by constant monitoring of whether social behaviors conform to the expectations of the domain, or whether they cross the boundary. Behaviors that cross the boundary will require greater attentional and information processing resources, while they are evaluated and the identity of the domain reappraised. The implication is that unexpected, domain incongruent behaviors will be highly salient. In this study, we set out to examine whether brain regions of the salience network are activated by social domain incongruent stimuli.

A key contrast for the study of the neuroscience of domains functioning is between social cues that are congruent to a setting where a narrow range of emotions and behaviors can be accommodated, and those that are domain incongruent, because of their focus or intensity. For the present study, this is a contrast between different kinds of social cues presented in the work setting, on the one hand, task-focused cues that are domain congruent, and, on the other hand, personalfocused cues that are domain incongruent. Equally, it is predicted that this contrast of task versus personal focus will not have the same implications within a marital or partner relationship where a wide range of emotions and behaviors is expected and congruent. Thus, we include contrasts not only between different kinds of social cues within work but also with the same set of scenarios depicted within a partner relationship.

It is plausible that mechanisms in detecting and responding to domain incongruent social cues are both unconscious, in order to underpin rapid responses, and conscious, in order to facilitate reappraisal of the assumed rules for interaction and action. According to some recent hypotheses, the AIC is involved in initial unconscious processing that determines priorities for higher level attentional resources in networks associated with consciousness (Michel, 2017). We therefore sought to evaluate whether salience network activation is associated with reaction time (a likely index of initial unconscious processes) to make a judgment about a stimulus and with conscious evaluation of its inappropriateness.

In this fMRI study, we first tested the hypothesis that social cues that are congruent in marital or partner relationships will be associated with ratings of domain incongruence when displayed within a work setting and will therefore be associated with greater activation in the amygdala, dorsal anterior cingulate and fronto-insular regions. Second, we predicted that shorter reaction times from the appearance of domain incongruent stimuli to making the judgment of their inappropriateness will be associated with greater salient network activation in the domain incongruent, compared to domain congruent conditions. Third, the extent to which domain incongruent stimuli are judged as inappropriate will be associated with greater activation in the amygdala, dorsal anterior cingulate, and fronto-insular regions of incongruent stimuli. As inappropriateness may be associated with unpleasantness, likelihood, and uncertainty, these were examined as potential confounds.

We also examined frontopolar and anterior temporal activations, which have been shown to represent different aspects of knowledge relevant for understanding the rules underpinning social behavior. There is evidence to support the role of the frontopolar cortex as representing sequential aspects (Zahn et al., 2017) and the anterior temporal cortex as representing abstract conceptual aspects (Zahn et al., 2009). We predicted that there would be no selective activations in these regions for our domain-incongruent contrasts because activations of representations of social rules are entailed in both domain-congruent and domain-incongruent conditions.

Materials and methods

Participants

Twenty healthy volunteers (11 females) took part in this study. Participants were included if they met the

following criteria: 18-50 years old, no self-reported previous or current psychiatric disorders including depression, anxiety, eating disorders and drug/alcohol addiction, no neurological disorders, no significant head injury resulting in unconsciousness, no current use of medication known to affect mood or cognition, no first-degree relatives suffering from any psychiatric disturbances, smoking less than 5 cigarettes per day, drinking less than the UK government guidelines for weekly alcohol intake (males: 3-4 units per day; females: 2-3 units per day) and fluent in English. Participants were further interviewed using the M.I.N.I. International Neuropsychiatric Interview (Sheehan et al., 1998) and were excluded if they met the criteria for a current psychiatric diagnosis. All participants included in the study had at least 3-months work experience and had at least 3-months romantic experience. At the time of scanning, 8 participants were single and 12 in a relationship varying between 6 months - 7.5 years. At the time of scanning, 9 participants were in full-time employment, two were in part-time employment, and nine were in fulltime University education. A structural MRI scan was taken before fMRI scanning to aid analysis and to exclude data from participants who showed any evidence of structural abnormality. No participants were excluded from the subsequent analysis on this basis.

Procedure

Prior to their visit for scanning, participants read the information sheet, gave informed consent, and completed questionnaires online. On arrival, they were briefed about the scanning process, and then completed a series of practice trials in order to familiarize themselves with the task. The task was then presented in the scanner, and again out of the scanner together with selfreport questions about their responses to the task items. The study was approved by the University of Manchester Research Ethics Committees and all participants provided written informed consent after the study procedures were explained. Participants were reimbursed for their time and travel expenses.

Measures

Social domains assessment

The task was displayed on a Dell XT3 laptop using PsychoPy software (Pierce, 2007). Participants were presented in the scanner with short vignettes describing social interactions that differed by their congruence with the social domain, task focus being congruent with both work and partner relationships, and personal focus only to partner relationships. The vignettes were designed to exemplify four conditions: work congruent behaviors seen in work (WORK-TASK FOCUS) and in a partner

relationship (PARTNER-TASK FOCUS), and behaviors that are congruent in a partner relationship (PARTNER-PERSONAL FOCUS) but not in work (WORK-PERSONAL FOCUS). An example of WORK-TASK FOCUS was "A work colleague gets angry with you for leaving the light on" and of PARTNER-TASK FOCUS "Your partner gets angry with you for leaving the light on". An example of PARTNER-PERSONAL FOCUS was "Your partner gets angry with you for not noticing that they have changed their hairstyle" and of WORK-PERSONAL FOCUS "A work colleague gets angry with you for not noticing that they have changed their hair style". The WORK-PERSONAL FOCUS vs WORK-TASK FOCUS contrasts were used to test the hypotheses, with the PARTNER-TASK FOCUS vs PARTNER-PERSONAL FOCUS contrasts controlling for differences of focus that did not introduce domain incongruence, and the PARTNER–PERSONAL FOCUS controlling for simple effects of personal focus.

Participants were asked to report how unpleasant they would feel ("mildly" or "very") by pressing a button within 6 seconds. If they responded before the end of the 6 seconds their response remained on the screen and the alternative disappeared. This was followed by a jittered intertrial interval with a mean duration of 4 seconds. This was jittered in nine steps of 500 ms around the mean interval with equal distribution of intertrial intervals across different stimulus types. There were 26 vignettes for each of the four conditions that had been selected from a battery of 60 vignettes (total 240) to provide stimuli that were matched as closely as possible on degree of unpleasantness as rated by a panel of 20 volunteers.

Self-report responses to the task

After the imaging session, participants provided selfreport ratings for each trial. Perceived inappropriateness was rated for each social scenario on a Likert scale (0 = not at all inappropriate, 100 = highly inappropriate), and reaction times from display of the social scenarios to judgment of inappropriateness were recorded. Likert scales included as potential confounds were of: unpleasantness (0 = not at all unpleasant, 100 = extremely unpleasant), likelihood of its happening (0 = very unlikely, 100 = very likely), and how certain they would be of how to respond (0 = very uncertain, 100 = very uncertain). Reaction times for making these judgments were recorded.

MRI scanning

MRI data were acquired on a 3 T Philips Achieva scanner (Philips Medical Systems, the Netherlands) with an eight-channel coil. A series of echo-planar images (EPI) was acquired for each participant using a sequence optimized for detecting ventral frontal signals (240 volumes; 40 axial slices; 3 mm slice thickness; ascending sequential acquisition; repetition time: 2000 ms; echo time: 22 ms; field of view: 240 \times 240 \times 120 mm; acquisition matrix: 80 \times 80 voxels; reconstructed voxel size: 3 mm³; flip angle: 90°). A three-dimensional T1-weighted magnetizationprepared rapid-acquisition gradient-echo (MPRAGE) structural image was also acquired for each participant (160 axial slices; 0.9 mm slice thickness; repetition time: 8.4 ms; echo time: 3.9 ms; field of view: $240 \times 191 \times 144$ mm; acquisition matrix: 256×163 voxels; reconstructed voxel size: $0.94 \times 0.94 \times 0.9$ mm; flip angle: 8°). In order to rule out clinically significant neurological abnormalities, T2-weighted structural images were also acquired. Statistical Parametric Mapping (SPM12: https://www.fil.ion.ucl.ac.uk/spm/soft ware/spm12/) and the general linear model were used for image analysis (Friston, Frith, Turner, & Frackowiak, 1995; Worsley & Friston, 1995).

Statistical analyses

Functional images were realigned, unwarped, coregistered to the participant's T1-weighted images and normalized to the [SPM12] Wellcome Trust Centre for Neuroimaging [Internet] (2014) template using the transformation parameters for the T1-weighted image. A smoothing kernel of 8-mm full-width at half maximum was then applied. At the first level, the four conditions; PARTNER-PERSONAL, PARTNER-TASK FOCUS, WORK-PERSONAL FOCUS, and WORK-TASK FOCUS were modeled as t-contrasts for each subject. For the second level, we then set up an ANOVA model with the four conditions as the within-subject factor. In order to test the first hypothesis, within-group differences in WORK-PERSONAL FOCUS vs WORK-TASK FOCUS were examined. A single bilateral combined a priori region of interest (ROI) mask was defined for small volume multiple comparison correction at a voxel-based FWE-corrected level of p < 0.05 which was composed of the Automated Anatomical Labeling (AAL) atlas-defined anterior cingulate cortex (ACC), bilateral amygdala, bilateral insula, frontal pole (BA10, Talairach Daemon atlas as implemented into the WFU Pick-Atlas version 2.4) and temporal poles. We report all activations which either survived a voxel-based familywise error corrected p < 0.05 over the whole brain or the ROI mask. All data underwent registration and 12-parameter affine normalization and were transformed into standard MNI space using a $2 \times 2 \times 2$ isotropic resolution. Data were smoothed using an 8 mm full width at half maximum Gaussian kernel. In a further analysis, we confirmed that the resulting clusters were not driven by the PARTNER context by exclusively masking them with PARTNER-PERSONAL FOCUS and PARTNER-TASK FOCUS at p = 0.001 uncorrected.

In further confirmatory analyses, we extracted the cluster averages for regions surviving multiple comparison correction (voxel-based few p < 0.05 as described above) for analyses in SPSS. This allowed us to test the second hypothesis by regressing the difference between WORK-TASK FOCUS and WORK-PERSONAL FOCUS activations in each cluster on the differences in reaction times (recorded outside of the scanner) to judgment of inappropriateness across the conditions. For the third hypothesis the difference in activations was regressed on to perceived inappropriateness, and for the fourth it was regressed on to the contrast in expressions of anger or distress across the conditions. All analyses are controlled for perceived unpleasantness, uncertainty, or likelihood. In view of the sample size separate models were examined for each of the three confounds.

Results

Self-report measures

Table 1 shows the mean scores for perceived inappro-priateness of the four conditions, and reaction times toself-report of domain inappropriateness, together withpotentiallyconfoundingself-reportvariables.Respondentsreportedthatscenariosdepicting

 Table 1. Mean self-report ratings of the social scenarios and reaction times to report of domain inappropriateness.

	Work personal focus	Work task focus	Partner personal focus	Partner task focus
Inappropriateness	70.35 (10.19)	43.80 ^c (10.08)	24.80 (11.58) ***	28.20 (11.32) ***
Reaction time	2.76 (1.18)	3.01 (1.32)	2.66 (1.07)	2.59 (1.20)
Unpleasantness	62.45 (8.84)	50.90 (11.43) ***	54.75 (12.58)	44.30 (15.80) ***
Likelihood	34.80 (18.63)	51.25 (15.92) ***	47.35 (12.55) ***	54.35 (19.20) ***
Certainty	40.25 (12.50)	52.35 (15.8) ***	55.55 (13.35) ***	61.55 (17.00) ***

Reaction times were times to make the judgment of inappropriateness when the stimuli were viewed outside of the scanner.

Means and standard deviations, in parenthesis are shown. Means were compared using repeated measures ANOVA. Models for unpleasantness, likelihood, certainty, and inappropriateness were significant at p < 0.001, and for reaction time at p = 0.010. Superscripts indicate significance of comparisons with the domain incongruent WORK – PERSONAL FOCUS as the reference condition.

$$p^* = p < 0.05^{**} = p < 0.01^{***} = p < 0.001$$

disclosure of personal information or strong emotions in the work setting (WORK-PERSONAL FOCUS) were more inappropriate, less likely, and would create less certainty as to how to respond, than WORK-TASK FOCUS and PARTNER-PERSONAL FOCUS scenarios (p < 0.01). WORK-PERSONAL FOCUS scenarios were reported to be more unpleasant than WORK-TASK FOCUS scenarios (p < 0.001). Reaction times to the judgment of appropriateness of WORK-PERSONAL did not differ from reaction times to the other types of scenarios.

fMRI results consistent across participants

The contrast between WORK-PERSONAL FOCUS vs WORK-TASK FOCUS was associated with a selective activation in the bilateral fronto-insular cortex (Figure 1) with the highest level of activation on the right hemisphere (Table 2, Figure 2). Further analyses showed that this cluster remained significant, when exclusively masked with activations associated with PARTNER-PERSONAL FOCUS and PARTNER-TASK FOCUS, showing that this region was not activated by personal focus per se, or by the differences in focus in general.

In confirmatory data analyses using the extracted cluster averages, there was a significant main effect of



Figure 1. Shows an axial slice through the bilateral frontoinsular clusters of activation for the contrast of work: Personal focus vs. Work: Task focus. Maps were thresholded at p = 0.001uncorrected with no voxel threshold and inclusively masked with the a priori ROI. MRIcron (Rorden et al., 2007) was used for visual display.

	Table 2.	fMRI	comparison o	f work context:	Personal	focus vs.	work context:	Task focus.
--	----------	------	--------------	-----------------	----------	-----------	---------------	-------------

	Peak MNI Coordinates								
Hemisphere	Region	Х	Y	Z	Peak z score	Cluster Size	FWE-Corrected p Value		
Right	Fronto-insular cortex	45	26	-3	4.69	88	0.003*		
	Dorsal anterior cingulate cortex	15	23	28	4.24	37	0.019		
Left	Fronto-insular cortex	-27	23	-10	4.38	62	0.011*		

Results were thresholded at p = 0.001 uncorrected, only regions surviving additional Familywise error (FWE) Small Volume Correction at the voxel-level <.05 over our combined apriori ROI mask were reported. No significant voxels remained when correcting over the volume of the whole brain.

* Remained significant at *p* = .05 voxel-wise FWE correction after additional exclusive masking by Work Context: Personal Focus vs. Partner Context: Work Focus and Work Context: Personal Focus vs. Work Context: Work Focus.



Figure 2. Plot to show right fronto-insular cortex activation across each of the conditions. Error bars represent one standard error of the mean.

condition [*F*(3, 57) = 2.96, *p* < 0.04, η_p^2 = 0.14] on right fronto-insular activation illustrated in Figure 2, and in post hoc analyses, only WORK-TASK FOCUS and WORK-PERSONAL FOCUS differed significantly [*t*(1,19) = 5.41, *p* < 0.001]. In similar analyses, hypothesized effects on dACC activations were not found, and nor were there effects on the amygdala, frontopolar, and anterior temporal regions.

Individual difference effects on fMRI results and the role of potential confounders

Table 3 shows associations between scores for the differences between WORK-PERSONAL FOCUS and WORK-TASK FOCUS activations in the right fronto-insular and differences between the same conditions in self-report and reaction times for judgment of inappropriateness. Right front-insular activation was associated with greater perceived inappropriateness of WORK-PERSONAL FOCUS, and lower reaction times for judgment of its inappropriateness. Perceived inappropriateness of the WORK-PERSONAL FOCUS over WORK-TASK FOCUS was associated with lower likelihood, while greater unpleasantness of WORK-PERSONAL FOCUS was associated with shorter reaction time to judgment of inappropriateness.

In multiple linear regression, greater perceived inappropriateness of the WORK-PERSONAL FOCUS compared to the WORK-TASK FOCUS scenarios, predicted right fronto-insular activation for the same comparison, controlling for unpleasantness (Table 4), and in separate models controlling for likelihood (inappropriateness $\beta = 0.55$, p = 0.04) and uncertainty (inappropriateness $\beta = 0.56$, p = 0.03).

Shorter reaction times also predicted right frontoinsular controlling for perceived unpleasantness (Table 5), and controlling for unlikelihood ($\beta = -0.67$, p < .01) and uncertainty

-0.22

0.18

0.54*

(p = 0.012)

ontrasting WORK-PERSONAL FOCUS and WORK-TASK FOCUS conditions.								
	Right fronto-insular activation	Inappropriate-ness	Reaction time (RT)	Unpleasant-ness	Unlikeli-hood			
Inappropriateness	0.52^*							
RT	(p = 0.02) 0.62**	-0.21						

-0.43 (p = 0.06)

0.22

0.05

0.41

0.47

0.39

(p = 0.07)

(p = 0.04)

 Table 3. Bivariate associations between self-report and reaction time differences and right fronto-insular activation differences, contrasting WORK-PERSONAL FOCUS and WORK-TASK FOCUS conditions.

Values shown are Pearson's r.

Unpleasantness

Unlikelihood

Uncertainty

Values of p < 0.1 are shown in parentheses. * = $p < 0.05^{**} = p < 0.01$

(p = 0.007)

0.33

0.19

0.09

Reaction times (RT) were times to make the judgment of inappropriateness when the stimuli were viewed outside of the scanner.

Table 4. Summary of multiple linear regression models predicting right fronto-insular activation from self-report domain inappropriateness.

	$\square R^2$	р		β	р
Model 1	0.10	0.19	Unpleasantness	0.31	0.19
Model 2	0.19	0.05	Unpleasantness	0.14	

In separate regression analyses the effect of inappropriateness controlling for unlikelihood was β = 0.55, p = 0.04, and controlling for uncertainty β = 0.56, p = 0.03

For all models, the scores were the differences between self-report or right fronto-insular activations during WORK – PERSONAL FOCUS and WORK – TASK FOCUS scenarios.

Table 5. Summary of multiple linear regression models predicting right fronto-insular activation from reaction time to the judgment of domain inappropriateness.

	$\square R^2$	р		β	р
Model 1	0.10	0.19	Unpleasantness	0.31	0.19
Model 2	0.26	0.02	Unpleasantness	0.05	0.84
			Reaction time	-0.58	0.02

In separate regression analyses the effect of inappropriateness controlling for unlikelihood was $\beta = -0.67$, p = 0.003, and for uncertainty $\beta = -0.60$, p = 0.008.

For all models, the scores were the differences between reaction time, selfreport, or right fronto-insular activations during WORK – PERSONAL FOCUS and WORK – TASK FOCUS scenarios

Table 6. Summary of multiple linear regression models predicting right fronto-insular activation from rating of domain inappropriateness and reaction time to the indication of inappropriateness.

	$\square R^2$	р		β	р
Model 1 Model 2	0.27 0.24	0.02 0.01	Inappropriateness Inappropriateness Reaction time	0.52 0.41 –0.50	0.02 0.04 0.01

For all models, the scores were the differences between reaction time, selfreport, or right fronto-insular activations during WORK – PERSONAL FOCUS and WORK – TASK FOCUS scenarios.

 $(\beta = -0.60, p < 0.01)$. When examined jointly, without confounds, both perceived inappropriateness and shorter reaction times to indication of inappropriateness made independent contributions to right fronto-insular activation in the WORK-PERSONAL FOCUS vs WORK-TASK FOCUS comparison (Table 6).

Discussion

Social scenarios depicting interactions with work colleagues with a personal focus, predicted to be inappropriate to the domain, and hence in need of prioritizing for reappraisal and action, were associated with increased fronto-insular activation, consistent with activation of the salience network. This selectivity was not seen in the IdACC region. Reported inappropriateness of the WORK-PERSONAL FOCUS compared to the WORK-TASK FOCUS stimuli, and shorter reaction time to indication of inappropriateness across these conditions, were independently associated with elevated fronto-insular activation.

We are aware that this is the first study to examine whether violations of expectations involving complex social stimuli typical of those encountered in everyday life are associated with activation of the salience system. Our findings are consistent with the hypothesis that everyday interactions are underpinned by shared rules regarding expected behaviors and that departures from these expectations are rapidly detected. Salience depends not only on behavior but also on the domain in which it occurs. Importantly, scenarios with personal focus did not activate the right fronto-insular when portrayed in the context of a partner relationship, only in interaction with a work colleague. Furthermore, the finding that both shorter reaction time to indicating inappropriateness and the judgment of inappropriateness were independently associated with fronto-insular activation is consistent with its role in unconscious and conscious mapping of social domains incongruence.

Our findings are also relevant to understanding psychopathology. We have previously reported that limitations in the demarcation of social domains, likely to be associated with impaired "patrolling" of domains boundaries, "domain disorganization", is elevated in adults with Borderline Personality Disorder (BPD) compared to psychiatric patients with other personality disorders and no personality disorder (J. Hill et al., 2008; Morse et al., 2009). In a study comparing adults with Borderline Personality Disorder with healthy controls while playing an economic exchange game, violations of expectations of fair exchange were associated with anterior insula activation in the controls but not in those with BPD (King-Casas et al., 2008). BPD's patients therefore may fail to activate neural mechanisms associated with detection of norm violations, including norms associated with domain congruence.

The study had a number of strengths that enabled us to examine the specificity of the findings. While the key comparison was between behaviors with personal focus seen in partner relationships where they are to be expected, and in work relationships where they are not, the further comparison of task focussed behaviors between the domains allowed us to control for general effects of domain and focus. Furthermore, we were able to account for the extent to which any differences arose from the perceived unpleasantness of the behaviors. By obtaining self-report and reaction time data we were able to examine whether the observed activation was associated with both conscious awareness of appropriateness and performance. The main limitation of the study was the modest sample size, which while adequate for examination of limited numbers of possible confounds, was

not sufficiently large to include all those of interest in the same model. In addition, although unpleasantness (i.e., valence) is associated with arousal, we did not specifically gather participant's ratings of arousal.

The theoretical framework on which this study was based has features in common with other current theories regarding social decision-making (Rilling & Sanfey, 2011). These hypothesize that interpretations of others' behaviors are based on expectations derived from social norms, which as outlined earlier speeds up social information processing. Neural mechanisms that track violations alerts the person to reassess their expectations. Our findings are consistent with previous studies that have identified fronto-insular activation in response to violations, but different from others that have for other regions, for example, the anterior cingulate cortex (Chang & Sanfey, 2013). However, they have to be evaluated in the light of the differences, in theoretical framework and experimental tasks, compared to previous studies of social decision-making. While other theoretical approaches assume that qualities such as fairness and trust apply generally across social interactions, the domains framework proposes that there are classes of social interaction each with their distinctive and different norms and values. Consequently, the task that we employed was tested, not for violations of rules of exchange, but for violations of theoretically predicted demarcations of social domains. In spite of the differences from previous findings, those described here are consistent with the indication that there may be general mechanisms for responding to norm violations that are employed to patrol the boundaries of social domains.

Acknowledgments

The authors thank Emily Pye and staff at the NIHR/Wellcome Trust Clinical Research Facility, Manchester, UK.

Disclosure statement

No potential conflict of interest was reported by the author(s).

ORCID

A. R. Bland () http://orcid.org/0000-0002-7170-1491 R. Elliott () http://orcid.org/0000-0002-7602-010X

References

Allman, J. M., Tetreault, N. A., Hakeem, A. Y., Manaye, K. F., Semendeferi, K., Erwin, J. M., Park, S., Goubert, V., & Hof, P. R. (2010). The von Economo neurons in frontoinsular and anterior cingulate cortex in great apes and humans. Brain Structure & Function, 214(5–6), 495–517. https://doi. org/10.1007/s00429-010-0254-0

- Bolton D. and Hill J. (2004). *Mind, Meaning, and Mental Disorder: The Nature of Causal Explanation in Psychology and Psychiatry* (Oxford: Oxford University Press).
- Bugental, D. B. (2000). Acquisition of the algorithms of social life: A domain-based approach. *Psychological Bulletin*, 126(2), 187. https://doi.org/10.1037/0033-2909.126.2.187
- Butti, C., Sherwood, C. C., Hakeem, A. Y., Allman, J. M., & Hof, P. R. (2009). Total number and volume of Von Economo neurons in the cerebral cortex of cetaceans. *Journal of Comparative Neurology*, *515*(2), 243–259. https://doi.org/10.1002/cne.22055
- Chang, L. J., & Sanfey, A. G. (2013). Great expectations: Neural computations underlying the use of social norms in decision-making. *Social Cognitive and Affective Neuroscience*, 8(3), 277–284. https://doi.org/10.1093/scan/ nsr094
- Craig, A. D. (2010). The sentient self. *Brain Structure & Function*, 214(5–6), 563–577. https://doi.org/10.1007/s00429-010-0248-y
- Critchley, H.D. (2005). Neural mechanisms of autonomic, affective, and cognitive integration. J. Comp. *Neurol.*, 493: 154– 166. https://doi.org/10.1002/cne.20749
- Friston KJ, Frith CD, Turner R, Frackowiak RS. (1995). Characterizing evoked hemodynamics with fMRI Neuroimage. Jun;2(2):157–65. doi:10.1006/nimg.1995.1018. PMID: 9343598.
- Gigerenzer, G. (2008). 'Why Heuristics Work', Perspectives on Psychological Science 3: 20–29.
- Hakeem, A. Y., Sherwood, C. C., Bonar, C. J., Butti, C., Hof, P. R., & Allman, J. M. (2009). Von Economo neurons in the elephant brain. *The Anatomical Record: Advances in Integrative Anatomy and Evolutionary Biology: Advances in Integrative Anatomy and Evolutionary Biology, 292*(2), 242–248. https:// doi.org/10.1002/ar.20829
- Heimer, L., & Van Hoesen, G. W. (2006). The limbic lobe and its output channels: Implications for emotional functions and adaptive behavior. *Neuroscience and Biobehavioral Reviews*, 30(2), 126–147. https://doi.org/10.1016/j.neubiorev.2005.06. 006
- Hill, I., Pilkonis, P. A., & Bear, I. (2010). Social domains, personality, and interpersonal functioning. In L. M. Horowitz & S. Strack (Eds.), *Handbook of interpersonal psychology: Theory, research, assessment, and therapeutic interventions* (pp. 281). John Wiley & Sons.
- Hill, J., Pilkonis, P., Morse, J., Feske, U., Reynolds, S., Hope, H., ... Broyden, N. (2008). Social domain dysfunction and disorganization in borderline personality disorder. *Psychological Medicine*, 38(1), 135–146. https://doi.org/10.1017/ S0033291707001626
- Hof, P. R., & Van der Gucht, E. (2007). Structure of the cerebral cortex of the humpback whale, Megaptera novaeangliae (Cetacea, Mysticeti, Balaenopteridae). *The Anatomical Record: Advances in Integrative Anatomy and Evolutionary Biology: Advances in Integrative Anatomy and Evolutionary Biology, 290*(1), 1–31. https://doi.org/10.1002/ar.20407
- Kahneman, D., Slovic, P., & Tversky, A. (1982). Judgment Under Uncertainty: Heuristics and Biases (New York: Cambridge University Press).
- King-Casas, B., Sharp, C., Lomax-Bream, L., Lohrenz, T., Fonagy, P., & Montague, P. R. (2008). The rupture and repair

of cooperation in borderline personality disorder. *science*, *321*(5890), 806–810. https://doi.org/10.1126/science. 1156902

- Luo, Y., Qin, S., Fernandez, G., Zhang, Y., Klumpers, F., & Li, H. (2014). Emotion perception and executive control interact in the salience network during emotionally charged working memory processing. *Human Brain Mapping*, 35(11), 5606–5616. https://doi.org/10.1002/hbm.22573
- Michel, M. (2017). A role for the anterior insular cortex in the global neuronal workspace model of consciousness. *Consciousness and Cognition*, *49*, 333–346. https://doi.org/ 10.1016/j.concog.2017.02.004
- Morse, J. Q., Hill, J., Pilkonis, P. A., Yaggi, K., Broyden, N., Stepp, S., Reed, L. I., & Feske, U. (2009). Anger, preoccupied attachment, and domain disorganization in borderline personality disorder. *Journal of Personality Disorders*, 23(3), 240–257. https://doi.org/10.1521/pedi.2009.23.3.240
- Nimchinsky, E. A., Gilissen, E., Allman, J. M., Perl, D. P., Erwin, J. M., & Hof, P. R. (1999). A neuronal morphologic type unique to humans and great apes. *Proceedings of the National Academy of Sciences*, 96(9), 5268–5273. https://doi. org/10.1073/pnas.96.9.5268
- Phan, K. L., Taylor, S. F., Welsh, R. C., Ho, S. H., Britton, J. C., & Liberzon, I. (2004). Neural correlates of individual ratings of emotional salience: A trial-related fMRI study. *Neuroimage*, 21(2), 768–780. https://doi.org/10.1016/j.neuroimage.2003. 09.072
- Pierce, J. W. (2007). PsychoPy—psychophysics software in Python. Journal of Neuroscience Methods, 162(1), 8–13. https://doi.org/10.1016/j.jneumeth.2006.11.017
- Ptak, R. (2012). The frontoparietal attention network of the human brain: Action, saliency, and a priority map of the environment. *The Neuroscientist*, *18*(5), 502–515. https://doi. org/10.1177/1073858411409051
- Rilling, J. K., & Sanfey, A. G. (2011). The neuroscience of social decision-making. *Annual Review of Psychology*, 62(1), 23–48. https://doi.org/10.1146/annurev.psych.121208. 131647
- Rorden, C., Karnath, H. O., & Bonilha, L. (2007). Improving lesion-symptom mapping. *Journal of cognitive neuroscience*, 19(7), 1081–1088. https://doi.org/10.1162/jocn.2007.19.7. 1081

- Rose, M. (1928). Die Inselrinde des Menschen und der Tiere. Journal of Neurology and Psychology, 37, 467–624.
- Rosen, M. L., Sheridan, M. A., Sambrook, K. A., Dennison, M. J., Jenness, J. L., Askren, M. K., ... McLaughlin, K. A. (2017). Salience network response to changes in emotional expressions of others is heightened during early adolescence: Relevance for social functioning. *Developmental Science*. 21. https://doi.org/10.1111/desc.12571
- Seeley, W. W., Allman, J. M., Carlin, D. A., Crawford, R. K., Macedo, M. N., Greicius, M. D., DeArmond, S. J., & Miller, B. L. (2007). Divergent social functioning in behavioral variant frontotemporal dementia and Alzheimer disease: Reciprocal networks and neuronal evolution. *Alzheimer Disease & Associated Disorders*, 21(4), S50–S57. https://doi. org/10.1097/WAD.0b013e31815c0f14
- Sheehan, D. V., Lecrubier, Y., Sheehan, K. H., Amorim, P., Janavs, J., Weiller, E., Hergueta, T., Baker, R., & Dunbar, G. C. (1998). The Mini-International Neuropsychiatric Interview (MINI): The development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *Journal* of Clinical Psychiatry, 59(suppl 20), 22–33.
- Spitzer, M., Fischbacher, U., Herrnberger, B., Grön, G., & Fehr, E. (2007). The neural signature of social norm compliance. *Neuron*, *56*(1), 185–196. https://doi.org/10.1016/j.neuron. 2007.09.011
- [SPM12] Wellcome Trust Centre for Neuroimaging [Internet]. (2014). http://www.fil.ion.ucl.ac.uk/spm/software/spm12/
- Williamson, P., & Allman, J. M. (2011). The human illnesses: Neuropsychiatric disorders and the nature of the human brain. Oxford University Press.
- Worsley, K. J., & Friston, K. J. (1995). Analysis of fMRI time-series revisited-again. *Neuroimage*, 2(3), 173–181.
- Zahn, R., Green, S., Beaumont, H., Burns, A., Moll, J., Caine, D., ... Ralph, M. A. L. (2017). Frontotemporal lobar degeneration and social behaviour: Dissociation between the knowledge of its consequences and its conceptual meaning. *Cortex*, 93, 107–118. https://doi.org/10.1016/j.cortex.2017.05.009
- Zahn, R., Moll, J., Iyengar, V., Huey, E. D., Tierney, M., Krueger, F., & Grafman, J. (2009). Social conceptual impairments in frontotemporal lobar degeneration with right anterior temporal hypometabolism. *Brain*, 132(3), 604–616. https://doi.org/10. 1093/brain/awn343