

On the inverse source identification problem in L^∞ for fully nonlinear elliptic PDE

Article

Published Version

Creative Commons: Attribution 4.0 (CC-BY)

Open Access

Ayanbayev, B. and Katzourakis, N. ORCID:
<https://orcid.org/0000-0001-5292-270X> (2021) On the inverse source identification problem in L^∞ for fully nonlinear elliptic PDE. Vietnam Journal of Mathematics, 49 (3). pp. 815-829. ISSN 2305-2228 doi: <https://doi.org/10.1007/s10013-021-00515-6> Available at <https://centaur.reading.ac.uk/99741/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1007/s10013-021-00515-6>

Publisher: Springer Singapore

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online



On the Inverse Source Identification Problem in L^∞ for Fully Nonlinear Elliptic PDE

Birzhan Ayanbayev¹ · Nikos Katzourakis²

Received: 20 May 2020 / Accepted: 13 April 2021 / Published online: 22 July 2021
© The Author(s) 2021

Abstract

In this paper we generalise the results proved in N. Katzourakis (SIAM J. Math. Anal. **51**, 1349–1370, 2019) by studying the ill-posed problem of identifying the source of a fully nonlinear elliptic equation. We assume Dirichlet data and some partial noisy information for the solution on a compact set through a fully nonlinear observation operator. We deal with the highly nonlinear nonconvex nature of the problem and the lack of weak continuity by introducing a two-parameter Tychonov regularisation with a higher order L^2 “viscosity term” for the L^∞ minimisation problem which allows to approximate by weakly lower semicontinuous cost functionals.

Keywords Regularisation strategy · Tychonov regularisation · Inverse source identification problem · Fully nonlinear elliptic equations · Calculus of Variations in L^∞

Mathematics Subject Classification (2010) 35R25 · 35R30 · 35J60 · 35J70

1 Introduction

Let $n, k \in \mathbb{N}$ with $k, n \geq 2$ and let $\Omega \subseteq \mathbb{R}^n$ be a bounded connected domain with $C^{1,1}$ regular boundary $\partial\Omega$. Let also

$$F : \Omega \times \mathbb{R} \times \mathbb{R}^n \times \mathbb{R}_s^{n \otimes 2} \longrightarrow \mathbb{R}$$

be a Carathéodory function, namely $x \mapsto F(x, r, p, X)$ is Lebesgue measurable for all $(r, p, X) \in \mathbb{R} \times \mathbb{R}^n \times \mathbb{R}_s^{n \otimes 2}$ and $(r, p, X) \mapsto F(x, r, p, X)$ is continuous for a.e. $x \in \Omega$.

Dedicated to Enrique Zuazua on the occasion of his 60th birthday.

✉ Nikos Katzourakis
n.katzourakis@reading.ac.uk

Birzhan Ayanbayev
bayanbayev@zedat.fu-berlin.de

¹ Freie Universität Berlin, Arnimallee 6, 14195 Berlin, Germany

² Department of Mathematics and Statistics, University of Reading, Whiteknights, PO Box 220, Reading RG6 6AX, UK

In this paper, the notation $\mathbb{R}_s^{n^{\otimes k}}$ stands for the vector space of fully symmetric k -th order tensors in $\mathbb{R}^n \otimes \dots \otimes \mathbb{R}^n$ (k -times). Given $g \in W^{2,\infty}(\Omega)$, consider the Dirichlet problem

$$\begin{cases} F[u] = f & \text{in } \Omega, \\ u = g & \text{on } \partial\Omega, \end{cases} \tag{1.1}$$

for some appropriate source $f : \Omega \rightarrow \mathbb{R}$. Here $F[u]$ denotes the induced fully nonlinear 2nd order differential operator, defined on smooth functions u as

$$F[u] := F(\cdot, u, Du, D^2u). \tag{1.2}$$

Evidently, we are employing the standard symbolisations $Du = (D_i u)_{i=1,\dots,n}$, $D^2u = (D_{ij}^2 u)_{i,j=1,\dots,n}$ and $D_i \equiv \partial/\partial x_i$. The above direct Dirichlet problem for F asks to determine u , given a source f and boundary data g . (In fact the source f is obsolete and can be absorbed into F , but for the problem we are interested in this paper it is more convenient to write it in this separated form). This is a semi-classical problem which is essentially standard material, see e.g. [22]. In particular, it is known that under various sets of assumptions on F that (1.1) is well-posed and, given $f \in L^\infty(\Omega)$ and $g \in W^{2,\infty}(\Omega)$, for any $p > n$ there exists a unique solution u in the locally convex (Fréchet) space

$$\mathcal{W}_g^{2,\infty}(\Omega) := \bigcap_{1 < p < \infty} (W^{2,p} \cap W_g^{1,p})(\Omega).$$

In general, the solution u is not in the smaller space $(W^{2,\infty} \cap W_g^{1,\infty})(\Omega)$ (not even locally), due to the failure of the $W^{2,p}$ estimates for $p = \infty$, which happens even in the linear case (see e.g. [21]). Additionally, (1.1) satisfies for any $p > n$ the fully nonlinear L^p global estimate

$$\|F(\cdot, v, Dv, D^2v)\|_{L^p(\Omega)} \geq C_1 \|v\|_{W^{2,p}(\Omega)} - C_1 \|g\|_{W^{2,p}(\Omega)} - C_2 \tag{1.3}$$

for some constants $C_1, C_2 > 0$ depending only on the parameters and any $v \in (W^{2,p} \cap W_g^{1,p})(\Omega)$. For sufficient conditions on F which guarantee the satisfaction of solvability of (1.1) in the strong sense and of the uniform estimate (1.3) we refer to [12, 13, 16, 28, 30].

Note that the above problem contains as a special case the archetypal instance of divergence operators with C^1 matrix coefficient A , as well as the non-divergence linear case with continuous coefficient:

$$\begin{cases} L_1[u] = \operatorname{div}(ADu) + b \cdot Du + cu, \\ L_2[u] = A : D^2u + b \cdot Du + cu. \end{cases} \tag{1.4}$$

In the above, the notations “ \cdot ” and “ $:$ ” symbolise the Euclidean inner products in the space of symmetric matrices $\mathbb{R}_s^{n^{\otimes 2}}$ and in \mathbb{R}^n respectively. More generally, the inner product of two tensors $T, S \in \mathbb{R}_s^{n^{\otimes k}}$ will also be denoted by “ $:$ ”, that is

$$T : S := \sum_{1 \leq a_1, \dots, a_k \leq n} T_{a_1 \dots a_k} S_{a_1 \dots a_k}.$$

The *inverse problem* relating to (1.1) asks the question of perhaps determining f , given the boundary data g and some other *partial information on the solution u , typically some approximate experimental measurements of some function of it known only up to some error*. The inverse problem is *severely ill-posed* even in the linear case of the Laplacian operator $F = \Delta$, as the noisy data measured on a subset of Ω might either not be compatible with any exact solution, or they may not suffice to determine a unique f even if compatibility holds true.

The above type of inverse problems are especially crucial for various applications, even in the model case of the Poisson equation, see e.g. [1, 8, 17, 23, 31, 32, 34, 36–40]. In this paper we will assume that the approximate information on u takes the form

$$K[u] = k^\gamma \quad \text{on } \mathcal{K}, \tag{1.5}$$

where K is an *observation operator*, taken to be a first order fully nonlinear differential operator of the form

$$K[u] := K(\cdot, u, Du), \tag{1.6}$$

where K and its partial derivatives K_r, K_p satisfy

$$K, K_r \in C(\mathcal{K} \times \mathbb{R} \times \mathbb{R}^n), \quad K_p \in C(\mathcal{K} \times \mathbb{R} \times \mathbb{R}^n; \mathbb{R}^n). \tag{1.7}$$

In (1.5) and (1.7), \mathcal{K} symbolises the set on which we take measurements, which will be assumed to satisfy

$$\mathcal{K} \subseteq \overline{\Omega} \text{ is compact and exists } \kappa \in [0, n] : \mathcal{H}^\kappa(\mathcal{K}) < \infty. \tag{1.8}$$

In the above, \mathcal{H}^κ denotes the Hausdorff measure of dimension κ . Our general measure and functional notation will be either standard or self-explanatory, e.g. as in [15, 18, 27]. Finally, $k^\gamma \in L^\infty(\mathcal{K}, \mathcal{H}^\kappa)$ is the function of approximate (deterministic) measurements taken on \mathcal{K} , at noise level at most $\gamma > 0$:

$$\|k^\gamma - k^0\|_{L^\infty(\mathcal{K}, \mathcal{H}^\kappa)} \leq \gamma, \tag{1.9}$$

where $k^0 = K[u^0]$ corresponds to ideal error-free measurements of an exact solution to (1.1) with source $f = F[u^0]$.

To recapitulate, in this paper we study the following ill-posed inverse source identification problem for fully nonlinear elliptic PDEs:

$$\begin{cases} F[u] = f & \text{in } \Omega, \\ u = g & \text{on } \partial\Omega, \\ K[u] = k^\gamma & \text{on } \mathcal{K}. \end{cases} \tag{1.10}$$

This means that we are searching for a selection process of a suitable approximation for f from the data k^γ on \mathcal{K} through the observation $K[u]$ of the solution u . To the best of our knowledge, (1.10) has not been studied before, at least in this generality. Our approach does not exclude the extreme cases of $\mathcal{K} = \overline{\Omega}$ (full information) or of $\mathcal{K} = \emptyset$ (no information), although trivial changes are required in the proofs. Sadly, an exact solution may not exist as the constraint may be incompatible with the solution of (1.1), owing to the errors in measurements. On the other hand, it is not possible to have a uniquely determined source on the constraint-free region $\Omega \setminus \mathcal{K}$. Instead, our goal is a strategy to determine an optimally fitting u^γ (and respective source $f^\gamma := F[u^\gamma]$) to the ill-posed problem (1.10). A popular choice of operator K in the literature (when $L = \Delta$) consist of some term of the separation of variables formula, as e.g. in [39].

Herein, we follow an approach based on recent advances in Calculus of Variations in the space L^∞ (see e.g. [26] and references therein) developed for functionals involving higher order derivatives, which has already been applied to the special case of the inverse source problem for linear PDEs (1.4) in [25]. This relatively new field was pioneered by

Gunnar Aronsson in the 1960s (see e.g. [3–7]) and is still a very active area of research; for a review of the by-now classical theory involving scalar first order functionals we refer to [24].

Following [25], we aim at providing a *regularisation strategy* inspired by the classical Tykhonov regularisation strategy in L^2 (see e.g. [29, 33]). As a first possible step, consider the next putative L^∞ “error” functional:

$$E_{\infty,\alpha}(u) := \|K[u] - k^\gamma\|_{L^\infty(\mathcal{K}, \mathcal{H}^\kappa)} + \alpha \|F[u]\|_{L^\infty(\Omega)}, \quad u \in \mathcal{W}_g^{2,\infty}(\Omega), \quad (1.11)$$

for some fixed parameter $\alpha > 0$. The advantage of searching for a best fitting solution in L^∞ is evident: we can keep the error term $|K[u] - k^\gamma|$ uniformly small and not merely small on average, as would happen if one chose to minimise some integral of a power of the error instead. As in [25], the goal would be to minimise $E_{\infty,\alpha}$ over $\mathcal{W}_g^{2,\infty}(\Omega)$, and then any minimiser of (1.11) would provide a candidate solution for our problem. Then, for any fixed α , this would be the best fitting solution with the least possible uniform error, namely $F[u] \cong f$ uniformly on Ω and $K[u] \cong k^\gamma$ uniformly on \mathcal{K} . Unfortunately, even if one momentarily ignores the problem of lack of regularity for (1.11) and the fact that $\mathcal{W}_g^{2,\infty}(\Omega)$ is not a Banach space, the main problem is that in general minimisers do not exist in the genuine fully nonlinear case of operator F (namely when $X \mapsto F(x, r, p, X)$ is nonlinear) as (1.11) is not weakly lower semicontinuous in the Fréchet space $\mathcal{W}_g^{2,\infty}(\Omega)$, as the highest order term may be nonconvex/non-quasiconvex. In the special linear case of [25], this problem was not present as the linearity of the differential operator was implying the desired weak lower semi-continuity.

In this work we resolve the problem explained above by proposing a double approximation method (or rather triple, as we will see shortly) which involves an additional Tykhonov or “viscosity” term which effectively is a weakly lower-semicontinuous approximation of (1.11). Hence, we will consider instead

$$\begin{cases} E_{\infty,\alpha,\beta}(u) := \|K[u] - k^\gamma\|_{L^\infty(\mathcal{K}, \mathcal{H}^\kappa)} + \alpha \|F[u]\|_{L^\infty(\Omega)} + \frac{\beta}{2} \|D^{\bar{n}}u\|_{L^2(\Omega)}^2, \\ u \in \left(W^{\bar{n},2} \cap W_g^{1,2}\right)(\Omega), \end{cases} \quad (1.12)$$

where $\bar{n} := [n/2] + 3$. In the above $\beta > 0$ is a fixed parameter, $[\cdot]$ symbolises the integer part and $D^{\bar{n}}u$ is the \bar{n} -th order weak derivative of u .

It is well known in the Calculus of Variations in L^∞ that (global) minimisers of supremal functionals, although usually simple to obtain with a standard direct minimisation [15, 19], they are not genuinely minimal as they do not share the nice “local” optimisation properties of minimisers of their integral counterparts (see e.g. [10, 35]). The case of (1.12) studied herein is no exception to this rule. A relatively standard method is bypass these obstructions is to employ minimisers of L^p approximating functionals as $p \rightarrow \infty$, establishing appropriate convergence of L^p minimisers to a limit L^∞ minimiser. The idea underlying this approximation technique is based on the simple measure theory fact that the L^p norm (of a function in $L^1 \cap L^\infty$) converges to the L^∞ norm as $p \rightarrow \infty$. This method is quite standard in the field and furnishes a selection principle of L^∞ minimisers with additional desirable properties (see e.g. [9, 11, 14, 20, 26]). In this fashion one is also able to bypass the lack of differentiability of supremal functionals and derive necessary PDE conditions satisfied by L^∞ extrema. This is indeed the method that is employed in this work as well, along the lines of [25].

We now present the main results to be established in this paper. As already explained, we will obtain *special* minimisers of (1.12) as limits of minimisers of

$$\left\{ \begin{aligned} E_{p,\alpha,\beta}(u) &:= \| |K[u] - k^\gamma |_{(p)} \|_{L^p(\mathcal{K}, \mathcal{H}^\kappa)} + \alpha \| |F[u]|_{(p)} \|_{L^p(\Omega)} + \frac{\beta}{2} \| D^{\bar{n}} u \|_{L^2(\Omega)}^2, \\ u &\in (W^{\bar{n},2} \cap W_g^{1,2})(\Omega). \end{aligned} \right. \tag{1.13}$$

In (1.13) we have used the normalised L^p norms

$$\|f\|_{L^p(\mathcal{K}, \mathcal{H}^\kappa)} := \left(\int_{\mathcal{K}} |f|^p d\mathcal{H}^\kappa \right)^{1/p}, \quad \|f\|_{L^p(\Omega)} := \left(\int_{\Omega} |f|^p d\mathcal{L}^n \right)^{1/p}$$

and the integral signs with slashes symbolise the average with respect to the Hausdorff measure \mathcal{H}^κ and the Lebesgue measure \mathcal{L}^n , respectively. Further, $|\cdot|_{(p)}$ symbolises the next p -regularisation of the absolute value away from zero:

$$|a|_{(p)} := \sqrt{|a|^2 + p^{-2}}.$$

Let us also note that, due to our L^p -approximation method, as an auxiliary result we also provide an L^p regularisation strategy for finite p as well, which is of independent interest. For the proof we will need to assume that the Dirichlet problem (1.1) for the fully nonlinear operator F satisfies the $W^{2,p}$ elliptic estimates (1.3) for all large enough (finite) $p > n$, as well the following: F and its partial derivatives F_r, F_p, F_X satisfy

$$\left\{ \begin{aligned} F, F_r &\in C\left(\bar{\Omega} \times \mathbb{R} \times \mathbb{R}^n \times \mathbb{R}_s^{n \otimes 2}\right), \\ F_p &\in C\left(\bar{\Omega} \times \mathbb{R} \times \mathbb{R}^n \times \mathbb{R}_s^{n \otimes 2}, \mathbb{R}^n\right), \\ F_X &\in C\left(\bar{\Omega} \times \mathbb{R} \times \mathbb{R}^n \times \mathbb{R}_s^{n \otimes 2}; \mathbb{R}_s^{n \otimes 2}\right). \end{aligned} \right. \tag{1.14}$$

We note that sufficient general conditions of when such operators satisfy $W^{2,p}$ elliptic estimates can be found for instance in the papers [12, 13, 16, 28, 30].

Theorem 1 *Let $\Omega \subseteq \mathbb{R}^n$ be a bounded $C^{1,1}$ domain and $g \in W^{\bar{n},2}(\Omega)$. Suppose also the operators (1.2) and (1.6) are given, satisfying the assumptions (1.3), (1.7), (1.8) and (1.14). Suppose further a function $k^\gamma \in L^\infty(\mathcal{K}, \mathcal{H}^\kappa)$ is given which satisfies (1.9) for $\gamma > 0$. Let finally $\alpha, \beta > 0$ be fixed. Then, we have the following results for the inverse problem associated to (1.10):*

- (i) *There exists a global minimiser $u_\infty \equiv u_\infty^{\alpha,\beta,\gamma} \in (W^{\bar{n},2} \cap W_g^{1,2})(\Omega)$ of the functional $E_{\infty,\alpha,\beta}$ defined in (1.12). In particular, we have $E_\infty(u_\infty) \leq E_\infty(v)$ for all $v \in (W^{\bar{n},2} \cap W_g^{1,2})(\Omega)$ and*

$$f_\infty \equiv f_\infty^{\alpha,\beta,\gamma} := F[u_\infty^{\alpha,\beta,\gamma}] \in L^\infty(\Omega).$$

Further, there exist signed Radon measures

$$\mu_\infty \equiv \mu_\infty^{\alpha,\beta,\gamma} \in \mathcal{M}(\Omega), \quad \nu_\infty \equiv \nu_\infty^{\alpha,\beta,\gamma} \in \mathcal{M}(\mathcal{K})$$

such that the nonlinear divergence PDE

$$K_r[u_\infty] \nu_\infty - \operatorname{div} (K_p[u_\infty] \nu_\infty) + \alpha ((dF)_{u_\infty})^* [\mu_\infty] + \beta (-1)^{\bar{n}} (D^{\bar{n}} : D^{\bar{n}} u_\infty) = 0, \tag{1.15}$$

is satisfied by the triplet $(u_\infty, \mu_\infty, v_\infty)$ in the sense of distributions (see (2.2)). In (1.15), the operator $((dF)_{u_\infty})^*$ is the formal adjoint of the linearisation of F at u_∞ , defined via duality as

$$((dF)_{u_\infty})^*[v] := \operatorname{div}(\operatorname{div}(F_X[u_\infty]v)) - \operatorname{div}(F_p[u_\infty]v) + F_r[u_\infty]v,$$

F_r, K_r, F_p, K_p, F_X denote the partial derivatives of F, K with respect to the respective variables and $F_r[v], K_r[v], F_p[v], K_p[v], F_X[v]$ denote the respective differential operators $F_r(\cdot, v, Dv, D^2v), K_r(\cdot, v, Dv), \dots$ etc. Additionally, the error measure v_∞ is supported in the closure of the subset of \mathcal{K} of maximum error, namely

$$\operatorname{supp}(v_\infty) \subseteq \left\{ |K[u_\infty] - k^\gamma|^\star = \|K[u_\infty] - k^\gamma\|_{L^\infty(\mathcal{K}, \mathcal{H}^\kappa)} \right\}, \tag{1.16}$$

where “ $(\cdot)^\star$ ” denotes the “essential limsup” with respect to $\mathcal{H}^\kappa \llcorner \mathcal{K}$ on \mathcal{K} (see Definition 3 that follows). If additionally the data function k^γ is continuous on \mathcal{K} , (1.16) can be improved to

$$\operatorname{supp}(v_\infty) \subseteq \{ |K[u_\infty] - k^\gamma| = \|K[u_\infty] - k^\gamma\|_{L^\infty(\mathcal{K}, \mathcal{H}^\kappa)} \}. \tag{1.17}$$

- (ii) For any $\alpha, \beta, \gamma > 0$, the minimiser u_∞ can be approximated by a family of minimisers $(u_p)_{p>n} \equiv (u_p^{\alpha, \beta, \gamma})_{p>n}$ of the respective L^p functionals (1.13) and the pair of measures $(\mu_\infty, v_\infty) \in \mathcal{M}(\Omega) \times \mathcal{M}(\mathcal{K})$ can be approximated by respective signed measures $(\mu_p, v_p)_{p>n} \equiv (\mu_p^{\alpha, \beta, \gamma}, v_p^{\alpha, \beta, \gamma})_{p>n}$, as follows:

For any $p > n$, (1.13) has a global minimiser $u_p \equiv u_p^{\alpha, \beta, \gamma}$ in $(W^{\bar{n}, 2} \cap W_g^{1, 2})(\Omega)$ and there exists a sequence $(p_j)_1^\infty$, such that

$$\begin{cases} u_p \longrightarrow u_\infty & \text{in } C^2(\bar{\Omega}), \\ D^k u_p \longrightarrow D^k u_\infty & \text{in } L^2(\Omega, \mathbb{R}_s^{\otimes k}), \text{ for all } k \in \{3, \dots, \bar{n} - 1\}, \\ D^{\bar{n}} u_p \rightharpoonup D^{\bar{n}} u_\infty & \text{in } L^2(\Omega, \mathbb{R}_s^{\otimes \bar{n}}), \end{cases} \tag{1.18}$$

as $p_j \rightarrow \infty$. Moreover, we have

$$\begin{cases} v_p := \frac{|K[u_p] - k^\gamma|_{(p)}^{p-2} (K[u_p] - k^\gamma)}{\mathcal{H}^\kappa(\mathcal{K}) \| |K[u_p] - k^\gamma|_{(p)} \|_{L^p(\mathcal{K}, \mathcal{H}^\kappa)}^{p-1}} \mathcal{H}^\kappa \llcorner \mathcal{K} \xrightarrow{*} v_\infty & \text{in } \mathcal{M}(\mathcal{K}), \\ \mu_p := \frac{|F[u_p]|_{(p)}^{p-2} F[u_p]}{\mathcal{L}^n(\Omega) \| |F[u_p]|_{(p)} \|_{L^p(\Omega)}^{p-1}} \mathcal{L}^n \llcorner \Omega \xrightarrow{*} \mu_\infty & \text{in } \mathcal{M}(\Omega), \end{cases} \tag{1.19}$$

as $p_j \rightarrow \infty$. Further, for each $p > n$, the triplet (u_p, μ_p, v_p) solves the PDE

$$K_r[u_p]v_p - \operatorname{div}(K_p[u_p]v_p) + \alpha((dF)_{u_p})^*[\mu_p] + \beta(-1)^{\bar{n}}(D^{\bar{n}} : D^{\bar{n}}u_p) = 0, \tag{1.20}$$

in the sense of distributions (see (2.1)).

- (iii) For any exact solution $u^0 \in (W^{\bar{n}, 2} \cap W_g^{1, 2})(\Omega)$ of (1.10) (with $f = F[u^0]$ and $K[u^0] = k^0$) corresponding to measurements with zero error, we have the estimate:

$$\|K[u_\infty^{\alpha, \beta, \gamma}] - K[u^0]\|_{L^\infty(\mathcal{K}, \mathcal{H}^\kappa)} \leq 2\gamma + \alpha \|F[u^0]\|_{L^\infty(\Omega)} + \frac{\beta}{2} \|D^{\bar{n}}u^0\|_{L^2(\Omega)}^2, \tag{1.21}$$

for any $\alpha, \beta, \gamma > 0$.

- (iv) For any exact solution $u^0 \in (W^{\bar{n}, 2} \cap W_g^{1, 2})(\Omega)$ of (1.10) (with $f = F[u^0]$ and $F[u^0] = k^0$) corresponding to measurements with zero error and for $p > n$, we have the estimate:

$$\|K[u_p^{\alpha, \beta, \gamma}] - K[u^0]\|_{L^p(\mathcal{K}, \mathcal{H}^\kappa)} \leq 2\gamma + \alpha \|F[u^0]\|_{L^p(\Omega)} + \frac{\beta}{2} \|D^{\bar{n}}u^0\|_{L^2(\Omega)}^2, \tag{1.22}$$

for any $\alpha, \beta, \gamma > 0$.

We note that the estimate in part (iv) is useful if we have merely that $F[u^0] \in L^p(\Omega)$ for $p < \infty$ (namely, when $F[u^0] \notin L^\infty(\Omega)$). We close this section by noting that the reader may find in [25] various comments and counter-examples regarding the optimality of Theorem 1 (therein stated for the case of a linear differential operator F).

2 Discussion, Auxiliary Results and Proofs

We begin with some clarifications on Theorem 1.

Remark 2 In index form, the definition of the formal adjoint $((dF)_{u_\infty})^*$ of the linearisation of F at u_∞ can be written as

$$((dF)_{u_\infty})^*[v] = \sum_{i,j=1}^n D_{ij}^2 (F_{X_{ij}}[u_\infty]v) - \sum_{k=1}^n D_k (F_{p_k}[u_\infty]v) + F_r[u_\infty]v$$

and its distributional interpretation via duality reads

$$\langle ((dF)_{u_\infty})^*[v], \phi \rangle = \int_\Omega \left\{ \sum_{i,j=1}^n (D_{ij}^2 \phi) F_{X_{ij}}[u_\infty] + \sum_{k=1}^n (D_k \phi) F_{p_k}[u_\infty] + \phi F_r[u_\infty] \right\} v d\mathcal{L}^n,$$

for all $\phi \in C_c^\bar{n}(\Omega)$. Hence, by taking into account the definitions of the measures μ_p, ν_p in (1.19), the distributional interpretation of (1.20) is

$$\begin{aligned} & \int_{\mathcal{K}} (K_r[u_p]\phi + K_p[u_p] \cdot D\phi) \frac{|K[u_p] - k^\gamma|_{(p)}^{p-2} (K[u_p] - k^\gamma)}{\| |K[u_p] - k^\gamma|_{(p)} \|_{L^p(\mathcal{K}, \mathcal{H}^k)}^{p-1}} d\mathcal{H}^k \\ & + \alpha \int_\Omega (F_r[u_p]\phi + F_p[u_p] \cdot D\phi + F_X[u_p] : D^2\phi) \frac{|F[u_p]|_{(p)}^{p-2} F[u_p]}{\| |F[u_p]|_{(p)} \|_{L^p(\Omega)}^{p-1}} d\mathcal{L}^n \\ & + \beta \int_\Omega D^{\bar{n}} u_p : D^{\bar{n}} \phi d\mathcal{L}^n = 0, \end{aligned} \tag{2.1}$$

for all $\phi \in C_c^\bar{n}(\Omega)$. Similarly, the distributional interpretation of (1.15) is

$$\begin{aligned} & \int_{\mathcal{K}} (K_r[u_\infty]\phi + K_p[u_\infty] \cdot D\phi) d\nu_\infty \\ & + \alpha \int_\Omega (F_r[u_\infty]\phi + F_p[u_\infty] \cdot D\phi + F_X[u_\infty] : D^2\phi) d\mu_\infty \\ & + \beta \int_\Omega D^{\bar{n}} u_\infty : D^{\bar{n}} \phi d\mathcal{L}^n = 0, \end{aligned} \tag{2.2}$$

for all $\phi \in C_c^\bar{n}(\Omega)$.

Further, we note that the physical interpretation of the measures μ_∞ and ν_∞ arising in the PDE is that they essentially “charge” the sets whereon $|F[u_\infty]|$ and $|K[u_\infty] - k^\gamma|$ are maximised over Ω and \mathcal{K} , respectively.

We now state a definition and a result taken from [25] which are required for our proofs.

Definition 3 (The essential limsup, [25]) Let $X \subseteq \mathbb{R}^n$ be a Borel set and let $\nu \in \mathcal{M}(X)$ be a finite positive Radon measure on X . Given $f \in L^\infty(X, \nu)$, we define $f^\star \in L^\infty(X, \nu)$ by

$$f^\star(x) := \lim_{\varepsilon \rightarrow 0} \left(\nu - \operatorname{ess\,sup}_{y \in \mathbb{B}_\varepsilon(x)} f(y) \right), \quad \forall x \in X,$$

and call f^\star the ν -essential limsup of f . Here, $\mathbb{B}_\varepsilon(x)$ denotes the open ball of radius ε centred at $x \in X$.

The following result studies what we call ‘‘concentration measures’’ of the approximate L^p minimisation problems as $p \rightarrow \infty$.

Proposition 4 (L^p concentration measures as $p \rightarrow \infty$) Let X be a compact metric space, endowed with a positive finite Borel measure ν which gives positive values to any open subset of X except \emptyset . Consider $(f_p)_1^\infty \subseteq L^\infty(X, \nu)$ and the sequence of signed Radon measures $(\nu_p)_1^\infty \subseteq \mathcal{M}(X)$, given by:

$$\nu_p := \frac{1}{\nu(X)} \frac{(|f_p|_{(p)})^{p-2} f_p}{\| |f_p|_{(p)} \|_{L^p(X, \nu)}^{p-1}} \nu, \quad p \in \mathbb{N},$$

where $| \cdot |_{(p)} = (|\cdot|^2 + p^{-2})^{1/2}$. Then:

- (i) There exists a subsequence $(p_i)_1^\infty$ and a limit measure $\nu_\infty \in \mathcal{M}(X)$ such that

$$\nu_{p_i} \xrightarrow{*} \nu_\infty \quad \text{in } \mathcal{M}(X),$$

as $p_i \rightarrow \infty$.

- (ii) If there exists $f_\infty \in L^\infty(X, \nu) \setminus \{0\}$ such that

$$\sup_X |f_{p_i} - f_\infty| \rightarrow 0 \quad \text{as } p \rightarrow \infty,$$

then ν_∞ is supported in the set where $|f_\infty|$ is maximised:

$$\operatorname{supp}(\nu_\infty) \subseteq \left\{ |f_\infty|^\star = \|f_\infty\|_{L^\infty(X, \nu)} \right\}.$$

- (iii) If additionally to (ii) the modulus $|f_\infty|$ of f_∞ is continuous on X , then the following stronger assertion holds:

$$\operatorname{supp}(\nu_\infty) \subseteq \left\{ |f_\infty| = \|f_\infty\|_{L^\infty(X, \nu)} \right\}.$$

Now we establish Theorem 1. The proof consists of several lemmas. We note that some details might be quite well known to the experts of Calculus of Variations, but we chose to give most of the niceties for the convenience of the readers and for the sake of completeness of the exposition.

Lemma 5 For any $p > n$ and $\alpha, \beta, \gamma > 0$, the functional (1.13) has a minimiser $u_p \in (W^{\bar{n}, 2} \cap W_g^{1, 2})(\Omega)$:

$$E_{p, \alpha, \beta}(u_p) = \inf \left\{ E_{p, \alpha, \beta}(v) : v \in (W^{\bar{n}, 2} \cap W_g^{1, 2})(\Omega) \right\}.$$

Proof Since $g \in W^{2,\infty}(\Omega)$

$$\begin{aligned} E_{p,\alpha,\beta}(g) &\leq E_{\infty,\alpha,\beta}(g) \\ &\leq \|k^\gamma\|_{L^\infty(\mathcal{K}, \mathcal{H}^k)} + \|K(\cdot, g, Dg)\|_{L^\infty(\mathcal{K}, \mathcal{H}^k)} \\ &\quad + \alpha \|F(\cdot, g, Dg, D^2g)\|_{L^\infty(\Omega)} + \frac{\beta}{2} \|D^{\bar{n}}g\|_{L^2(\Omega)}^2 \\ &< \infty. \end{aligned}$$

Hence,

$$0 \leq \inf \left\{ E_{p,\alpha,\beta}(v) : v \in (W^{\bar{n},2} \cap W_g^{1,2})(\Omega) \right\} \leq E_{\infty,\alpha,\beta}(g) < \infty.$$

Further, $E_{p,\alpha,\beta}$ is coercive in the space $(W^{\bar{n},2} \cap W_g^{1,2})(\Omega)$. Indeed, by our assumption (1.3), Hölder’s inequality and that $p > n \geq 2$, for any $v \in (W^{\bar{n},2} \cap W_g^{1,2})(\Omega)$ we have

$$\begin{aligned} E_{p,\alpha,\beta}(v) &\geq \alpha \|F[v]\|_{L^p(\Omega)} + \frac{\beta}{2} \|D^{\bar{n}}v\|_{L^2(\Omega)}^2 \\ &\geq \alpha (C_1 \|v\|_{W^{2,p}(\Omega)} - C_1 \|g\|_{W^{2,p}(\Omega)} - C_2) + \frac{\beta}{2} \|D^{\bar{n}}v\|_{L^2(\Omega)}^2 \\ &\geq \alpha C_1 \|v\|_{W^{2,2}(\Omega)} + \frac{\beta}{2} \|D^{\bar{n}}v\|_{L^2(\Omega)}^2 - \alpha C_1 \|g\|_{W^{2,p}(\Omega)} - \alpha C_2 \end{aligned}$$

which implies

$$\alpha C_1 \|v\|_{W^{2,2}(\Omega)} + \frac{\beta}{2} \|D^{\bar{n}}v\|_{L^2(\Omega)}^2 \leq E_{\infty,\alpha,\beta}(g) + \alpha C_1 \|g\|_{W^{2,p}(\Omega)} + \alpha C_2,$$

for any $v \in (W^{\bar{n},2} \cap W_g^{1,2})(\Omega)$. Now, by Poincaré inequality in $W_g^{1,2}(\Omega)$ we have

$$\|v\|_{L^2(\Omega)} \leq C (\|Dv\|_{L^2(\Omega)} + \|g\|_{W^{1,2}(\Omega)})$$

for some $C > 0$, and by the interpolation inequalities in the Sobolev space $W^{\bar{n},2}(\Omega)$, we have

$$\|D^k v\|_{L^2(\Omega)} \leq C \left(\|Dv\|_{L^2(\Omega)} + \|D^{\bar{n}}v\|_{L^2(\Omega)} \right),$$

for some $C > 0$ and any $k \in \{1, \dots, \bar{n}\}$. By putting the last three estimates together, we conclude that

$$\|v\|_{W^{\bar{n},2}(\Omega)} \leq C,$$

where the constant $C > 0$ in general depends on p but is uniform for $v \in (W^{\bar{n},2} \cap W_g^{1,2})(\Omega)$.

Let now $(u_p^m)_1^\infty$ be a minimising sequence of $E_{p,\alpha,\beta}$:

$$E_{p,\alpha,\beta}(u_p^m) \longrightarrow \inf \left\{ E_p(v) : v \in (W^{\bar{n},2} \cap W_g^{1,2})(\Omega) \right\},$$

as $m \rightarrow \infty$. Then, by the coercivity estimate, we have the uniform bound

$$\|u_p^m\|_{W^{\bar{n},2}(\Omega)} \leq C$$

for some $C > 0$ independent of $m \in \mathbb{N}$. By standard weak and strong compactness arguments in Sobolev and Hölder spaces, together with the Morrey estimate

$$\|v\|_{C^{k-[n/2]-1,\sigma}(\bar{\Omega})} \leq C \|v\|_{W^{k,2}(\Omega)},$$

applied to $k = \bar{n}$, there exists a subsequence $(u_p^{m_k})_1^\infty$ and $u_p \in (W^{\bar{n},2} \cap W_g^{1,2})(\Omega)$ such that, along this subsequence we have

$$\begin{cases} u_p^m \rightarrow u_p & \text{in } C^2(\bar{\Omega}), \\ D^k u_p^m \rightarrow D^k u_p & \text{in } L^2(\Omega, \mathbb{R}_s^{n \otimes k}), \text{ for all } k \in \{3, \dots, \bar{n} - 1\}, \\ D^{\bar{n}} u_p^m \rightarrow D^{\bar{n}} u_p & \text{in } L^2(\Omega, \mathbb{R}_s^{n \otimes \bar{n}}), \end{cases}$$

as $m_k \rightarrow \infty$. The above modes of convergence and the continuity of the function K defining the operator K imply that $K[u_p^m] \rightarrow K[u_p]$ uniformly on \mathcal{K} as $m_k \rightarrow \infty$. Therefore,

$$\left\| |K[u_p^m] - k^\gamma|_{(p)} \right\|_{L^p(\mathcal{K}, \mathcal{H}^k)} \rightarrow \left\| |K[u_p] - k^\gamma|_{(p)} \right\|_{L^p(\mathcal{K}, \mathcal{H}^k)}$$

as $m_k \rightarrow \infty$. Additionally, by the continuity of the function F defining the operator F and the uniform convergence of the minimising sequence up to second order derivatives, we have

$$F[u_p^m] \rightarrow F[u_p] \quad \text{in } C(\bar{\Omega}),$$

as $m_k \rightarrow \infty$. Finally, by weak lower semi-continuity in L^2 we have

$$\left\| D^{\bar{n}} u_p \right\|_{L^2(\Omega)} \leq \liminf_{k \rightarrow \infty} \left\| D^{\bar{n}} u_p^{m_k} \right\|_{L^2(\Omega)}.$$

By putting all the above together, we infer that

$$E_{p,\alpha,\beta}(u_p) \leq \liminf_{k \rightarrow \infty} E_{p,\alpha,\beta}(u_p^{m_k}) \leq \inf \left\{ E_{p,\alpha,\beta}(v) : v \in (W^{\bar{n},2} \cap W_g^{1,2})(\Omega) \right\},$$

which concludes the proof. □

Note that the proof above reveals the fact that $E_{p,\alpha,\beta}$ is weakly lower semi-continuous on the space $(W^{\bar{n},2} \cap W_g^{1,2})(\Omega)$, even though it is not explicitly stated.

Lemma 6 *For any $\alpha, \beta, \gamma > 0$, there exists a (global) minimiser u_∞ of $E_{\infty,\alpha,\beta}$ in the space $(W^{\bar{n},2} \cap W_g^{1,2})(\Omega)$, as well as a sequence of minimisers $(u_{p_i})_1^\infty$ of the respective $E_{p,\alpha,\beta}$ -functionals from Lemma 5, such that (1.18) holds true.*

Proof For each $p > n$, let $u_p \in (W^{\bar{n},2} \cap W_g^{1,2})(\Omega)$ be the minimiser of $E_{p,\alpha,\beta}$ given by Lemma 5. (We will follow a similar method and utilise the estimates appearing therein). For any fixed $q \in (n, \infty)$ and $p \geq q$, Hölder’s inequality and minimality yield

$$E_{q,\alpha,\beta}(u_p) \leq E_{p,\alpha,\beta}(u_p) \leq E_{p,\alpha,\beta}(g) \leq E_{\infty,\alpha,\beta}(g) < \infty.$$

By employing again the coercivity of $E_{q,\alpha,\beta}$, we have the estimate

$$E_{q,\alpha,\beta}(u_p) \geq \alpha \left(C_1 \|u_p\|_{W^{2,q}(\Omega)} - C_1 \|g\|_{W^{2,q}(\Omega)} - C_2 \right) + \frac{\beta}{2} \|D^{\bar{n}} u_p\|_{L^2(\Omega)}^2,$$

which by Poincaré’s inequality and the interpolation inequalities in $W^{\bar{n},2}(\Omega)$, yield

$$\sup_{p \geq q} \|u_p\|_{W^{\bar{n},2}(\Omega)} \leq C$$

for some $C > 0$ depending on q and all the parameters, but independent of p . By a standard diagonal argument, for any sequence $(p_i)_1^\infty$ with $p_i \rightarrow \infty$ as $i \rightarrow \infty$, there exists a function

$$u_\infty \in (W^{\bar{n},2} \cap W_g^{1,2})(\Omega)$$

and a subsequence (denoted again by $(p_i)_1^\infty$) along which (1.18) holds true. It remains to show that u_∞ is in fact a minimiser of E_∞ over the same space. To this end, note that for any fixed $q \in (n, \infty)$ and $p \geq q$, we have

$$E_{q,\alpha,\beta}(u_p) \leq E_{p,\alpha,\beta}(u_p) \leq E_{p,\alpha,\beta}(v) \leq E_{\infty,\alpha,\beta}(v)$$

for any $v \in (W^{\bar{n},2} \cap W_g^{1,2})(\Omega)$. By the weak lower semi-continuity of $E_{q,\alpha,\beta}$ in the space $(W^{\bar{n},2} \cap W_g^{1,2})(\Omega)$ demonstrated in Lemma 5, we have

$$E_{q,\alpha,\beta}(u_\infty) \leq \liminf_{i \rightarrow \infty} E_{q,\alpha,\beta}(u_{p_i}) \leq E_{\infty,\alpha,\beta}(v),$$

for any $v \in (W^{\bar{n},2} \cap W_g^{1,2})(\Omega)$. By letting $q \rightarrow \infty$ in the estimate above, we deduce that

$$E_{\infty,\alpha,\beta}(u_\infty) \leq \inf \left\{ E_{\infty,\alpha,\beta}(v) : v \in (W^{\bar{n},2} \cap W_g^{1,2})(\Omega) \right\},$$

as desired. □

Lemma 7 *For any $\alpha, \beta, \gamma > 0$ and $p > n$, consider the minimiser u_p of the functional $E_{p,\alpha,\beta}$ over $(W^{2,p} \cap W_g^{1,p})(\Omega)$ constructed in Lemma 5. Consider also the signed Radon measures $\mu_p \in \mathcal{M}(\Omega)$ and $\nu_p \in \mathcal{M}(\mathcal{K})$, defined in (1.19). Then, the triplet (u_p, μ_p, ν_p) satisfies the PDE (1.20) in the distributional sense, namely (2.1) holds true for all test functions $\phi \in C_c^\infty(\Omega)$.*

Proof We involve a standard Gateaux differentiability argument. Let us begin by checking that μ_p, ν_p as defined in (1.19) are uniformly bounded Radon measures when $u_p \in W^{\bar{n},2}(\Omega)$. Since by the regularity of F, K, u_p they obviously define absolutely continuous measures, it suffices to check that by Hölder inequality's, we have the total variation estimates

$$\begin{aligned} \|\nu_p\|(\mathcal{K}) &\leq \left(\| |K[u_p] - k^\gamma|_{(p)} \|_{L^p(\mathcal{K}, \mathcal{H}^\kappa)} \right)^{1-p} \int_{\mathcal{K}} |K[u_p] - k^\gamma|_{(p)}^{p-1} d\mathcal{H}^\kappa \\ &\leq \left(\| |K[u_p] - k^\gamma|_{(p)} \|_{L^p(\mathcal{K}, \mathcal{H}^\kappa)} \right)^{1-p} \left(\int_{\mathcal{K}} |K[u_p] - k^\gamma|_{(p)}^p d\mathcal{H}^\kappa \right)^{\frac{p-1}{p}} \\ &= 1 \end{aligned}$$

and similarly

$$\begin{aligned} \|\mu_p\|(\Omega) &\leq \left(\| |F[u_p]|_{(p)} \|_{L^p(\Omega)} \right)^{1-p} \int_{\Omega} |F[u_p]|_{(p)}^{p-1} d\mathcal{L}^n \\ &\leq \left(\| |F[u_p]|_{(p)} \|_{L^p(\Omega)} \right)^{1-p} \left(\int_{\Omega} |F[u_p]|_{(p)}^p d\mathcal{L}^n \right)^{\frac{p-1}{p}} \\ &= 1. \end{aligned}$$

Next, fix $\phi \in C_c^{\bar{n}}(\Omega)$. By using the regularity assumptions on F, K , we formally compute, recalling the abbreviations $F[v] = F(\cdot, v, Dv, D^2v)$ and $K[v] = K(\cdot, v, Dv)$:

$$\begin{aligned} & \left. \frac{d}{d\varepsilon} \right|_{\varepsilon=0} E_{p,\alpha,\beta}(u_p + \varepsilon\phi) \\ &= \left(\int_{\mathcal{K}} |K[u_p] - k^\gamma|_{(p)}^p d\mathcal{H}^k \right)^{\frac{1}{p}-1} \int_{\mathcal{K}} |K[u_p] - k^\gamma|_{(p)}^{p-2} (K[u_p] - k^\gamma) \cdot \\ & \quad \cdot [K_r[u_p]\phi + K_p[u_p] \cdot D\phi] d\mathcal{H}^k \\ & + \alpha \left(\int_{\Omega} |F[u_p]|_{(p)}^p d\mathcal{L}^n \right)^{\frac{1}{p}-1} \int_{\Omega} |F[u_p]|_{(p)}^{p-2} F[u_p] \cdot \\ & \quad \cdot [F_r[u_p]\phi + F_p[u_p] \cdot D\phi + F_X[u_p] : D^2\phi] d\mathcal{L}^n \\ & + \beta \int_{\Omega} D^{\bar{n}}u_p : D^{\bar{n}}\phi d\mathcal{L}^n. \end{aligned}$$

By invoking the definitions of μ_p, ν_p , the above yields that

$$\begin{aligned} \left. \frac{d}{d\varepsilon} \right|_{\varepsilon=0} E_{p,\alpha,\beta}(u_p + \varepsilon\phi) &= \int_{\mathcal{K}} [K_r[u_p]\phi + K_p[u_p] \cdot D\phi] d\nu_p \\ & + \alpha \int_{\Omega} [F_r[u_p]\phi + F_p[u_p] \cdot D\phi + F_X[u_p] : D^2\phi] d\mu_p \\ & + \beta \int_{\Omega} D^{\bar{n}}u_p : D^{\bar{n}}\phi d\mathcal{L}^n. \end{aligned}$$

Since u_p is the minimiser of $E_{p,\alpha,\beta}$, we have that $E_{p,\alpha,\beta}(u_p) \leq E_{p,\alpha,\beta}(u_p + \varepsilon\phi)$ for all $\varepsilon \in \mathbb{R}$ and all $\phi \in C_c^{\bar{n}}(\Omega)$. Hence, our computation implies that the PDE (1.20) is indeed satisfied as claimed, once we confirm that the formal computation is rigorous, and that therefore E_p is Gateaux differentiable at the minimiser u_p for any direction $\phi \in C_c^{\bar{n}}(\Omega)$ because by the continuity of F, K and the fact that $u_p \in (C^2 \cap W^{\bar{n},2})(\Omega)$, $F[u_p] \in C(\overline{\Omega})$ and $K[u_p] - k^\gamma \in L^\infty(\mathcal{K}, \mathcal{H}^k)$, Hölder’s inequality implies that

$$F_r[u_p]\phi + F_p[u_p] \cdot D\phi + F_X[u_p] : D^2\phi \in C(\overline{\Omega})$$

and

$$K_r(\cdot, u_p, Du_p)\phi + K_p(\cdot, u_p, Du_p) \cdot D\phi \in C(\mathcal{K})$$

for any $\phi \in C_c^{\bar{n}}(\Omega)$. Finally, $D^{\bar{n}}u_p : D^{\bar{n}}\phi \in L^1(\Omega)$ since $D^{\bar{n}}u_p \in L^2(\Omega, \mathbb{R}_s^{\otimes \bar{n}})$. □

Lemma 8 *For any $\alpha, \beta, \gamma > 0$, consider the minimiser u_∞ of $E_{\infty,\alpha,\beta}$ constructed in Lemma 6 and the minimisers $(u_p)_{p>n}$ of the functionals $(E_{p,\alpha,\beta})_{p>n}$. Then, there exist signed Radon measures $\mu_\infty \in \mathcal{M}(\Omega)$ and $\nu_\infty \in \mathcal{M}(\mathcal{K})$ such that the triplet $(u_\infty, \mu_\infty, \nu_\infty)$ satisfies the PDE (1.15) in the distributional sense, that is (2.2) holds true. Additionally, there exists a further subsequence along which the weak* modes of convergence of (1.19) hold true as $p \rightarrow \infty$.*

Proof By the proof of Lemma 7, we have the uniform in p total variation bounds $\|\mu_p\|(\Omega) \leq 1$ and $\|\nu_p\|(\mathcal{K}) \leq 1$. Hence, by the sequential weak* compactness of the spaces of Radon measures

$$\mathcal{M}(\Omega) = (C_0(\Omega))^*, \quad \mathcal{M}(\mathcal{K}) = (C(\mathcal{K}))^*, \tag{2.3}$$

there exists a further subsequence denoted again by $(p_i)_1^\infty$ such that $\mu_p \xrightarrow{*} \mu_\infty$ in $\mathcal{M}(\Omega)$ and $v_p \xrightarrow{*} v_\infty$ in $\mathcal{M}(\mathcal{K})$, as $p_i \rightarrow \infty$. Fix now $\phi \in C_c^{\bar{n}}(\Omega)$. By Lemma 7, we have that the triplet (u_p, μ_p, v_p) satisfies (1.20), that is (2.1) holds true for any fixed test function. Since $u_p \rightarrow u_\infty$ in $C^2(\bar{\Omega})$ as $p_i \rightarrow \infty$, $F, F_r, F_p, F_X, K_r, K_p$ are all continuous up to the boundary, we have as $p_i \rightarrow \infty$ that

$$\begin{aligned} K_r[u_p]\phi + K_p[u_p] \cdot D\phi &\rightarrow K_r K_r[u_\infty]\phi + K_p[u_\infty] \cdot D\phi, \\ F_r[u_p]\phi + F_p[u_p] \cdot D\phi + F_X[u_p] : D^2\phi &\rightarrow F_r[u_\infty]\phi + F_p[u_\infty] \cdot D\phi + F_X[u_\infty] : D^2\phi, \end{aligned}$$

uniformly on \mathcal{K} and on $\bar{\Omega}$ respectively, for any fixed $\phi \in C_c^{\bar{n}}(\Omega)$. Further, we have $\mu_p \xrightarrow{*} \mu_\infty$ and $v_p \xrightarrow{*} v_\infty$ in $\mathcal{M}(\Omega)$ and in $\mathcal{M}(\mathcal{K})$ respectively as $p_j \rightarrow \infty$. By standard properties of the weak*-strong continuity of the duality pairings (2.3), we have that the first two terms of (2.2) converge to the respective first two terms of (2.1), as $p_i \rightarrow \infty$. Finally, since $D^{\bar{n}}u_p \rightharpoonup D^{\bar{n}}u_p$ in $L^2(\Omega, \mathbb{R}_s^{n \otimes \bar{n}})$ and $D^{\bar{n}}\phi \in L^2(\Omega, \mathbb{R}_s^{n \otimes \bar{n}})$, we also have that the last term of (2.2) converges to the respective last term of (2.1), as $p_i \rightarrow \infty$. Hence, we have indeed obtained (1.15) by passing to the limit as $p_i \rightarrow \infty$ in (1.20). \square

Lemma 9 For any $\alpha, \beta, \gamma > 0, p > n$ and $u^0 \in (W^{2,p} \cap W_g^{1,p})(\Omega)$ such that

$$\|k^\gamma - K[u^0]\|_{L^\infty(\mathcal{K}, \mathcal{H}^\kappa)} \leq \gamma,$$

the (α, β, γ) -dependent minimiser u_p of E_p (constructed in Lemmas 5–8), satisfies the error bounds (1.22). If additionally $u^0 \in (W^{\bar{n},2} \cap W_g^{1,2})(\Omega)$, then the (α, β, γ) -dependent minimiser u_∞ of E_∞ (constructed in Lemmas 5–8), satisfies the error bounds (1.21).

Proof Let us denote $k^0 := K[u^0]$, noting that $k^0 \in C(\mathcal{K})$ and that

$$\|k^\gamma - k^0\|_{L^\infty(\mathcal{K}, \mathcal{H}^\kappa)} \leq \gamma.$$

Recall that for any $p > n, u_p$ is a global minimiser of $E_{p,\alpha,\beta}$ in $(W^{\bar{n},2} \cap W_g^{1,2})(\Omega)$. Therefore,

$$E_{p,\alpha,\beta}(u_p) \leq E_{p,\alpha,\beta}(u^0).$$

This implies

$$\begin{aligned} &\|K[u_p] - k^\gamma\|_{L^p(\mathcal{K}, \mathcal{H}^\kappa)} + \alpha \|F[u_p]\|_{L^p(\Omega)} + \frac{\beta}{2} \|D^{\bar{n}}u_p\|_{L^2(\Omega)}^2 \\ &\leq \|K[u^0] - k^\gamma\|_{L^p(\mathcal{K}, \mathcal{H}^\kappa)} + \alpha \|F[u^0]\|_{L^p(\Omega)} + \frac{\beta}{2} \|D^{\bar{n}}u^0\|_{L^2(\Omega)}^2. \end{aligned}$$

The above together with Minkowski’s and Hölder’s inequalities yields

$$\begin{aligned} \|K[u_p] - K[u^0]\|_{L^p(\mathcal{K}, \mathcal{H}^\kappa)} &\leq \|K[u^0] - k^\gamma\|_{L^p(\mathcal{K}, \mathcal{H}^\kappa)} + \|K[u^0] - k^\gamma\|_{L^p(\mathcal{K}, \mathcal{H}^\kappa)} \\ &\quad + \alpha \|F[u^0]\|_{L^p(\Omega)} + \frac{\beta}{2} \|D^{\bar{n}}u^0\|_{L^2(\Omega)}^2 \\ &= 2\|k^\gamma - k^0\|_{L^p(\mathcal{K}, \mathcal{H}^\kappa)} + \alpha \|F[u^0]\|_{L^p(\Omega)} + \frac{\beta}{2} \|D^{\bar{n}}u^0\|_{L^2(\Omega)}^2 \\ &\leq 2\gamma + \alpha \|F[u^0]\|_{L^p(\Omega)} + \frac{\beta}{2} \|D^{\bar{n}}u^0\|_{L^2(\Omega)}^2, \end{aligned}$$

as claimed. To obtain the corresponding estimate for u_∞ in the case that additionally $u^0 \in (W^{\bar{n},2} \cap W_g^{1,2})(\Omega)$, we pass to the limit as $p_i \rightarrow \infty$ in the estimate above and the conclusion

follows by letting $p_i \rightarrow \infty$ due to the strong convergence $u_{p_i} \rightarrow u_\infty$ in $C^2(\overline{\Omega})$ as $p_i \rightarrow \infty$. The lemma ensues. \square

The proof is now complete by noting that the statements (1.16)–(1.17) in Theorem 1 follow from Proposition 4 and the established modes of convergence.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Alves, C.J.S., Abdallah, J.B., Jaoua, M.: Recovery of cracks using a point-source reciprocity gap function. *Inverse Probl. Sci. Eng.* **12**, 519–574 (2004)
2. Ambrosio, L., Fusco, N., Pallara, D.: *Functions of Bounded Variation and Free Discontinuity Problems*. Oxford University Press, Oxford (2000)
3. Aronsson, G.: Minimization problems for the functional $\sup_x \mathcal{F}(x, f(x), f'(x))$. *Arkiv Mat.* **6**, 33–53 (1965)
4. Aronsson, G.: Minimization problems for the functional $\sup_x \mathcal{F}(x, f(x), f'(x))$ II. *Arkiv Mat.* **6**, 409–431 (1966)
5. Aronsson, G.: Extension of functions satisfying Lipschitz conditions. *Arkiv Mat.* **6**, 551–561 (1967)
6. Aronsson, G.: On certain minimax problems and Pontryagin's maximum principle. *Calc. Var. PDE* **37**, 99–109 (2010)
7. Aronsson, G., Barron, E.N.: L^∞ , variational problems with running costs and constraints. *Appl. Math. Optim.* **65**, 53–90 (2012)
8. El Badia, A., Ha Duong, T.: Some remarks on the problem of source identification from boundary measurements. *Inverse Probl.* **14**, 883 (1998)
9. Barron, E.N., Jensen, R.R., Wang, C.Y.: The Euler equation and absolute minimizers of L^∞ functionals. *Arch. Rational Mech. Anal.* **157**, 255–283 (2001)
10. Bocea, M., Nesi, V.: Γ -convergence of power-law functionals, variational principles in L^∞ , and applications. *SIAM J. Math. Anal.* **39**, 1550–1576 (2008)
11. Bocea, M., Popovici, C.: Variational principles in L^∞ with applications to antiplane shear and plane stress plasticity. *J. Convex Anal.* **18**, 403–416 (2011)
12. Caffarelli, L.A.: Interior a priori estimates for solutions of fully non-linear equations. *Ann. Math.* **130**, 189–213 (1989)
13. Caffarelli, L.A., Cabré, X.: *Fully Nonlinear Elliptic Equations*. American Mathematical Society, Providence (1995)
14. Champion, T., De Pascale, L., Prinari, F.: Γ -convergence and absolute minimizers for supremal functionals. *ESAIM: COCV* **10**, 14–27 (2004)
15. Dacorogna, B.: *Direct Methods in the Calculus of Variations*, 2nd edn. Applied Mathematical Sciences, vol. 78. Springer, Berlin (2008)
16. Dong, H., Krylov, N.V., Li, X.: On fully nonlinear elliptic and parabolic equations with VMO coefficients in domains. *St. Petersburg Math. J.* **24**, 39–69 (2013)
17. Engl, H.W., Hanke, M., Neubauer, A.: *Regularization of Inverse Problems*. Springer, Netherlands (1996)
18. Evans, L.C.: *Weak Convergence Methods for Nonlinear Partial Differential Equations*. Regional Conference Series in Mathematics, vol. 74. American Mathematical Society, Providence (1990)
19. Fonseca, I., Leoni, G.: *Modern Methods in the Calculus of Variations: L^p Spaces*. Springer Monographs in Mathematics. Springer, New York (2007)
20. Garroni, A., Nesi, V., Ponsiglione, M.: Dielectric breakdown: optimal bounds. *Proc. R. Soc. A* **457**, 2317–2335 (2001)

21. Giaquinta, M., Martinazzi, L.: An Introduction to the Regularity Theory for Elliptic Systems, Harmonic Maps and Minimal Graphs. Publications of the Scuola Normale Superiore. Edizioni della Normale Pisa (2012)
22. Gilbarg, D., Trudinger, N.: Elliptic Partial Differential Equations of Second Order. Classics in Mathematics, vol. 224 (Reprint of the 1998 edition). Springer, Berlin (2001)
23. Isakov, V.: Inverse Source Problems. Mathematical Surveys and Monographs, vol. 34. American Mathematical Society, Providence (1990)
24. Katzourakis, N.: An Introduction to Viscosity Solutions for Fully Nonlinear PDE with Applications to Calculus of Variations in L^∞ . Springer Briefs in Mathematics. Springer, Cham (2015)
25. Katzourakis, N.: An L^∞ regularisation strategy to the inverse source identification problem for elliptic equations. *SIAM J. Math. Anal.* **51**, 1349–1370 (2019)
26. Katzourakis, N., Moser, R.: Existence, uniqueness and structure of second order absolute minimisers. *Arch. Rational Mech. Anal.* **231**, 1615–1634 (2019)
27. Katzourakis, N., Várdućá, E.: An Illustrative Introduction to Modern Analysis. CRC Press/Taylor & Francis, Boca Raton (2017)
28. Krylov, N.V.: On the existence of W_p^2 solutions for fully nonlinear elliptic equations under either relaxed or no convexity assumptions. *Commun. Contemp. Math.* **19**(06), 1750009 (2017)
29. Kirsch, A.: An Introduction to the Mathematical Theory of Inverse Problems, 2nd edn. Applied Mathematical Sciences, vol. 120. Springer, New York (2011)
30. Li, D., Zhang, K.: $W^{2,p}$ interior estimates of fully nonlinear elliptic equations. *Bull. Lond. Math. Soc.* **47**, 301–314 (2015)
31. Ling, L., Hon, Y.C., Yamamoto, M.: Inverse source identification for Poisson equation. *Inverse Probl. Sci. Eng.* **13**, 433–447 (2005)
32. Magnoli, N., Viano, G.A.: The source identification problem in electromagnetic theory. *J. Math. Phys.* **38**, 2366–2388 (1997)
33. Neubauer, A.: An a posteriori parameter choice for Tikhonov regularization in Hilbert scales leading to optimal convergence rates. *SIAM J. Numer. Anal.* **25**, 1313–1326 (1988)
34. Nara, T., Ando, S.: A projective method for an inverse source problem of the Poisson equation. *Inverse Probl.* **19**, 355 (2003)
35. Ribeiro, A.N., Zappale, E.: Existence of minimisers for nonlevel convex supremal functionals. *SIAM J. Control Optim.* **52**, 3341–3370 (2014)
36. Shigeta, T., Hon, Y.-C.: Numerical source identification for Poisson equation. In: Tanaka, M. (ed.) *Inverse Problems in Engineering Mechanics IV* (Nagano, Japan), pp. 137–175. Elsevier (2003)
37. Xie, O., Zhao, Z.: Identifying an unknown source in the Poisson equation by a modified Tikhonov regularization method. *Int. J. Math. Comput. Sci.* **6**, 86–90 (2012)
38. Yang, F.: The truncation method for identifying an unknown source in the Poisson equation. *Appl. Math. Comput.* **22**, 9334–9339 (2011)
39. Yang, F., Fu, C.-L.: The modified regularization method for identifying the unknown source on Poisson equation. *Appl. Math. Model.* **36**, 756–763 (2012)
40. Zhao, Z., Meng, Z., You, L., Xie, O.: Identifying an unknown source in the Poisson equation by the method of Tikhonov regularization in Hilbert scales. *Appl. Math. Model.* **38**, 4686–4693 (2014)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.