

Detectability of COVID-19 global emissions reductions in local CO₂ concentration measurements

Article

Published Version

Creative Commons: Attribution 4.0 (CC-BY)

Open Access

Dacre, H. F. ORCID: <https://orcid.org/0000-0003-4328-9126>, Western, L. M., Say, D., O'Doherty, S., Arnold, T., Rennick, C. and Hawkins, E. ORCID: <https://orcid.org/0000-0001-9477-3677> (2021) Detectability of COVID-19 global emissions reductions in local CO₂ concentration measurements. *Environmental Research Letters*, 16 (9). 094043. ISSN 1748-9326 doi: 10.1088/1748-9326/ac1eda Available at <https://centaur.reading.ac.uk/99849/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1088/1748-9326/ac1eda>

Publisher: Institute of Physics

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

LETTER • **OPEN ACCESS**

Detectability of COVID-19 global emissions reductions in local CO₂ concentration measurements

To cite this article: H F Dacre *et al* 2021 *Environ. Res. Lett.* **16** 094043

View the [article online](#) for updates and enhancements.

You may also like

- [Anthropogenic CO₂ emissions assessment of Nile Delta using XCO₂ and SIF data from OCO-2 satellite](#)
Ankit Shekhar, Jia Chen, Johannes C Paetzold et al.
- [Land cover change in low-warming scenarios may enhance the climate role of secondary organic aerosols](#)
Marianne T Lund, Alexandru Rap, Gunnar Myhre et al.
- [Inventories and scenarios of nitrous oxide emissions](#)
Eric A Davidson and David Kanter

ENVIRONMENTAL RESEARCH
LETTERS

LETTER

OPEN ACCESS

RECEIVED
17 March 2021REVISED
5 August 2021ACCEPTED FOR PUBLICATION
18 August 2021PUBLISHED
3 September 2021

Original Content from
this work may be used
under the terms of the
[Creative Commons
Attribution 4.0 licence](#).

Any further distribution
of this work must
maintain attribution to
the author(s) and the title
of the work, journal
citation and DOI.

Detectability of COVID-19 global emissions reductions in local
CO₂ concentration measurementsH F Dacre¹ , L M Western² , D Say² , S O'Doherty², T Arnold^{3,4}, C Rennick³ and E Hawkins⁵¹ University of Reading, Department of Meteorology, Reading, United Kingdom² School of Chemistry, University of Bristol, Bristol, United Kingdom³ National Physical Laboratory, Teddington, Middlesex, United Kingdom⁴ School of GeoSciences, University of Edinburgh, Edinburgh, United Kingdom⁵ National Centre for Atmospheric Science, Department of Meteorology, University of Reading, Reading, United KingdomE-mail: h.f.dacre@reading.ac.uk**Keywords:** CO₂ emissions policy, climate change, multiple linear regression

Abstract

It is estimated that global anthropogenic carbon dioxide (CO₂) emissions reduced by up to 12% at the start of 2020 compared to recent years due to the COVID-19 related downturn in economic activity. Despite the large decrease in CO₂ emissions, no reduction in the trend in background atmospheric CO₂ concentrations has been detected. So, how long would it take for sustained COVID-19 CO₂ emission reductions to be detected in daily and monthly averaged local CO₂ concentration measurements? CO₂ concentration measurements for five measurement sites in the UK and Ireland are combined with meteorological numerical weather prediction data to build statistical models that can predict future CO₂ concentrations. It is found that 75% of the observed daily variability can be explained by these simple models. Emission reduction scenario experiments using these simple models illustrate that large daily and seasonal variability in local CO₂ concentrations precludes the rapid emergence of a detectable signal. COVID-19 magnitude emissions reductions would only be detectable in the daily CO₂ concentrations after at least 38 months and in monthly CO₂ concentrations after 11 months of sustained reductions. For monthly CO₂ concentrations the time of emergence is similar for all sites since the seasonal variability is largely driven by non-local fluxes of CO₂ between the terrestrial biosphere and the atmosphere. The COVID-19 CO₂ anthropogenic emissions reductions are similar in magnitude to those that are required to meet the Paris Agreement target of keeping global temperatures below 2 °C. This study demonstrates that, using measurements alone, there will be a considerable lag between changes in global anthropogenic emissions and a detected signal in local CO₂ concentration trends. Thus, there is likely to be a delay of several years between changes in policy designed to meet CO₂ anthropogenic emissions targets and our ability to detect the impact of these policies on CO₂ concentrations using atmospheric measurements alone.

1. Introduction

Electricity production, transportation and industrial activity account for more than 80% of carbon dioxide (CO₂) emissions from fuel combustion (Quadrelli and Peterson 2007). Since the start of 2020, COVID-19 restrictions have significantly reduced these activities. Current estimates suggest that global fossil fuel CO₂ emissions in 2020 may have dropped by around 7%–8% (Friedlingstein *et al* 2020, Hale and Leduc

2020, Le Quéré *et al* 2020, Liu *et al* 2020). Andreoni (2021) estimates that in Europe more than 195 600 thousand tons of CO₂ have been avoided between January and June 2020, compared to the same period of the previous year, representing a –12.1% emissions change. A decline in annual CO₂ emissions of this size would exceed any decline since the end of World War II. The magnitude of these emissions reductions is similar to those required to meet the target of the Paris Agreement, which aims to keep

the global temperature rise below 2°C (hereafter ‘Paris Agreement magnitude emissions reductions’). To meet the Paris Agreement temperature target, emissions from energy production and transport will have to peak almost immediately in the developed world (Annex I countries) and decline at about 10% each year until net-zero emissions are reached around 2030 (IPCC 2018). Thus the COVID-19 crisis presents a test bed for understanding these longer-term climate change policies on a more immediate time-scale.

While the recent reductions in CO₂ emissions are indeed substantial, they do not immediately equate to similar reductions in the trend of atmospheric CO₂ concentrations. Background CO₂ concentration measurements have not, so far, shown any changes as a result of COVID-19 emissions reductions (Liu *et al* 2020). This is consistent with previous situations when reductions in CO₂ associated with economic downturns did not significantly change the trend in CO₂ concentrations (Granados *et al* 2012). The lack of sensitivity to emissions reductions is due to the long atmospheric lifetime of CO₂ (50–200 years) which makes any perturbation in emission rate small compared to the reservoir of CO₂ currently present in the atmosphere. In addition, the large daily and seasonal variability of CO₂ concentrations makes changes in global CO₂ emissions difficult to detect (Samset *et al* 2020). Thus if we cannot expect immediately measurable impacts, how long would we need to wait to detect a change in the CO₂ concentration trend due to COVID-19 emission reductions?

In the climate change literature, many studies have investigated the response of the climate system to changes in greenhouse gases (Taylor and Penner 1994, Stainforth *et al* 2005, Sitch *et al* 2015). These studies typically involve running experiments with coupled atmosphere-ocean climate models with greenhouse gas forcing running over 50–100 year time periods. They may also be coupled to models of other processes in the Earth’s atmosphere such as the carbon cycle, so as to better simulate climate feedbacks such as interaction with the terrestrial ecosystems or oceans. On decadal to centennial timescales changes in CO₂ emissions can alter the climate. Therefore, these long integrations are necessary so that the response of the climate to the changing greenhouse gas emissions can reach equilibrium (Tebaldi and Friedlingstein 2013). However, on shorter timescales (days to months) CO₂ behaves more like a passive tracer. The concentrations of CO₂ on these timescales is largely controlled by changes in the weather and terrestrial biospheric activity. Therefore complex climate models which represent interactions occurring over longer timescales (years to decades) are not needed to capture the near-term consequence of changes in CO₂ emissions on CO₂ concentrations.

The aim of this work is to determine how long it would take for COVID-19/Paris Agreement magnitude emissions reductions to be detected in local daily and monthly CO₂ concentration measurements. We have built multiple linear regression (MLR) models, similar to those used to predict short-lived air quality pollutants (e.g. Carslaw and Beevers 2005, Dacre *et al* 2020), to predict CO₂ concentrations using only meteorological data and recent local CO₂ measurements. These models will not capture the responses in the complex models because they are tuned using recent data and they do not include climate feedbacks. However, these reduced complexity models may nevertheless be used to gain insights into our ability to detect greenhouse gas emissions reductions. In addition, they are much less computationally expensive, making them very fast to run.

2. Data

2.1. CO₂ data

The hourly atmospheric CO₂ measurements used in this study were taken from five *in-situ* observatories situated across the UK and Ireland (figure 1). Four of these stations, Tacolneston, Ridge Hill, Bilsdale and Heathfield, form the UK-based part of the UK Deriving Emissions linked to Climate Change (DECC 2020) network (Stanley *et al* 2018, Stavert *et al* 2019). Each of these sites makes use of a tall telecommunications tower to sample air from multiple height inlets (ranging from 42 to 248 m above ground level (magl) across the network). At each UK DECC site, we use data from the highest inlet only (table 1). The fifth site, Mace Head, is situated on the west coast of Ireland. This station is ideally positioned to intercept northern hemispheric background air from the North Atlantic. CO₂ measurements from Mace Head are made by Laboratoire des Sciences du Climat et de l’Environnement as part of the integrated carbon observation system (ICOS) network, from a 23 magl sample inlet (Vardag *et al* 2014). Note that lower-frequency CO₂ measurements are available from Mace Head data prior to 2011.

Figure 1 shows the average footprint emissions sensitivity obtained from 30 day backwards simulations of the Met Office’s numerical atmospheric dispersion modelling environment (NAME) model for the five sites for March 2020. The sensitivity is defined as the contribution per unit emission to the mole fraction measurement (Manning *et al* 2011). These footprints provide an indication of the location of the local emissions contributing to the measurements at each site in March 2020. At all five sites, continuous *in-situ* CO₂ measurements are made using cavity ring-down spectrometers (Picarro G2301 or G2401). At UK DECC sites, ambient measurements are corrected for linear instrumental drift via daily measurements of a standard gas. A small non-linear




Table 1. Location, inlet height (metres above ground level, magl) and data availability for the 5 CO₂ measurement sites used in this study.

Site	Lat/Lon	Inlet height (magl)	Data availability
Mace Head	53.327° N, 9.904° W	23	2011–present
Tacolneston	52.518° N, 1.139° E	185	2012–present
Ridge Hill	51.998° N, 2.540° W	90	2012–present
Bilsdale	54.359° N, 1.150° W	248	2014–present
Heathfield	50.977° N, 0.231° E	100	2013–present

correction is applied based on monthly analyses of four calibration gases that span above and below the ambient mole fraction range (Stanley *et al* 2018). The calibration strategy differs slightly at Mace Head, where ambient measurements are assigned a mole fraction based on comparison to a linear fit of four calibration cylinders. Like the UK sites, these calibration gases span the complete ambient range. All

calibration cylinders are of natural composition and were assigned CO₂ mole fractions at the World Calibration Centre at Empa or the GasLab, Max Planck Institute for Biogeochemistry, Jena, linking them to the World Meteorological Organization X2007 CO₂ calibration scale.

2.2. Meteorological data

The meteorological data to build the statistical models in this study comes from the UK Met Office UKV model (Tang *et al* 2013). Hourly data covering the period 1 January 2015–31 May 2020 is used. Hourly 4D-Var assimilation allows the production of state-of-the-art weather forecasts for the UK, initialised every hour (Ballard *et al* 2016). The UKV has a high resolution inner domain (1.5 km grid spacing) over the UK, separated from a lower resolution grid (4 km grid spacing) near the boundaries by a variable resolution transition zone. The high resolution contains a better representation of land surface processes and orography than coarser resolution global models. Sub-grid scale processes such as, boundary layer turbulence, radiation, cloud,

Table 2. Regression coefficients (β_i) from the MLR models for each measurement site. The coefficients quantify how much the daily CO₂ concentration is expected to increase/decrease when each explanatory variable (x_i) increases by one, holding all the other variables constant. Explanatory variables are monthly averaged temperature (averaged over the preceding month), daily averaged easterly wind speed (u-wind), daily averaged northerly wind speed (v-wind) and daily averaged boundary layer depth (BLD). Coefficients are only included where they are significant at the 95% level.

Explanatory variable (x_i)		Ridge Hill	Tacolneston	Bilsdale	Mace Head	Heathfield	Average
Date	β_1	0.007	0.007	0.007	0.007	0.007	0.007
Temperature (K)	β_2	−1.1	−1.0	−1.2	−1.5	−1.2	−1.2
U-wind (ms ^{−1})	β_3	3.4	1.2	2.4	2.3	1.8	2.2
V-wind (ms ^{−1})	β_4	—	−1.1	−0.8	0.7	1.4	0.2
BLD (m)	β_5	−0.004	−0.007	—	−0.002	−0.008	−0.005

microphysics and orographic drag are represented by parameterizations. The UKV model has been shown to compare well against observations (Lean *et al* 2008, Roberts and Lean 2008, Clark *et al* 2016). Meteorological data for the 5-year period (1 January 2015–31 December 2019) is extracted from the UKV and interpolated to the location of the DECC sites to build the statistical models described in section 3. One advantage of using weather model output is that above surface variables, such as boundary layer height, can be extracted, although modelled boundary layer height has not been evaluated at many sites in the UK due to lack of climatological measurements (Harvey *et al* 2013, 2015). One disadvantage of using weather model output is that the resolution of the model data is 1.5 km², whereas the CO₂ concentration data are point measurements. If they were available, use of local meteorological measurements to build the statistical models would be more accurate.

3. MLR modelling

In order to predict daily and monthly average CO₂ concentrations during the ongoing COVID-19 restrictions, MLR models are built using up to 5-years of DECC and UKV data (2015–2019). MLR modelling is used since the concentration of CO₂ is likely to depend on more than one predictor variable. The technique enables the relative influences of the predictor variables to be analysed, which allows us to perform the simple CO₂ emission scenario experiments described in section 5. MLR modelling is commonly used for predicting the variability in short-lived pollutant concentrations such as NO₂ (Shi and Harrison 1997, Carslaw and Beevers 2005, Dacre *et al* 2020). The MLR models predict the daily average CO₂ concentrations we would expect, during the COVID-19 period, given no change in CO₂ emissions. Several meteorological and temporal explanatory variables (x_i) are used to predict CO₂ concentrations (y) at each DECC station. The regression coefficients (β_i) describe the size of the effect of the explanatory variable on the daily CO₂ concentrations and α is the value y is predicted to have when all the explanatory variables are equal to zero.

$$y = \sum_{i=1}^n \beta_i x_i + \alpha. \quad (1)$$

The Akaike information criterion (AIC) is used to determine which explanatory variables to include in the models. The model with the lowest AIC score is expected to have the best balance between its ability to fit the data set and its ability to avoid over-fitting the data set. The explanatory variables and regression coefficients used in this study are shown in table 2. Since wind direction is cyclic not linear (i.e. 0 and 360 degrees have the same direction) it is partitioned into its northerly (v-wind) and easterly (u-wind) components. Wind speed, wind direction and temperature are all extracted 10 m above ground level. Sensitivity studies using meteorological variables extracted at the height of the sample inlets for each site did not improve the MLR models.

Our aim in the design of the MLR models was to keep the number of explanatory variables to a minimum and to restrict the models to use local data only. This is desirable to ensure that others, with only local CO₂ concentration and meteorological measurements available to them, can build similar models for their site locations. Also, for simplicity, the same explanatory variables are used for each of five sites analysed. The importance of each variable in explaining the observed CO₂ concentrations varies for each site, but the variables in table 2 were found to contribute to a reduced AIC for all five sites.

4. Evaluation of predicted CO₂ concentrations

In this section the CO₂ concentrations predicted by the MLR models are compared to the observed CO₂ concentrations at all five DECC sites. The evaluation is performed for various temporal averaging periods. The aim is to determine whether the MLR models are a credible representation of reality and thus can be used to perform emission scenario experiments.

4.1. Annual and seasonal CO₂ concentration variability

Figure 2 shows the yearly averaged observed and predicted CO₂ concentrations between January 2015




Figure 2. CO₂ concentrations from 1 January 2015 to 30 May 2020 at (a) Ridge Hill, (b) Tacolneston, (c) Bilsdale, (d) Heathfield and (e) Mace Head. Daily averaged observed concentrations (blue) and predicted concentrations (red). 30-day averaged observed concentrations (cyan) and predicted concentrations (orange). Annually averaged observed concentrations (black) and predicted concentrations (grey). Note that there is 2 weeks of missing meteorological data in 2017.

and June 2020. At all sites there is a monotonic increase in yearly averaged CO₂ concentrations. CO₂ concentrations are primarily rising because of the increased amounts of fossil fuels that humans are

burning for energy. The predicted CO₂ concentrations capture this annual increase in CO₂ concentrations due to the inclusion of the date in the MLR models with a coefficient of $0.007 \text{ ppm day}^{-1}$

at all sites which is equivalent to $2.5 \text{ ppm year}^{-1}$ (table 2).

Figure 2 also shows the monthly averaged observed and predicted CO_2 concentrations between January 2015 and June 2020. At all sites there is a strong annual cycle in CO_2 concentrations, with highest CO_2 concentrations measured during the winter months and lowest CO_2 concentrations measured during the summer months. This annual cycle is the result of photosynthetic activity by plants. As plants begin to photosynthesize in the spring and summer, they absorb CO_2 from the atmosphere and eventually use it as a carbon source for growth and reproduction. Once winter arrives, plants save energy by decreasing photosynthesis. Without photosynthesis, the dominant process is the exhalation of CO_2 by the total ecosystem, including bacteria, plants, and animals. The modelled CO_2 concentrations capture the annual cycle in CO_2 concentrations fairly well due to the inclusion of monthly averaged temperature in the MLR models. The coefficients used in the MLR models for monthly averaged temperature are all negative indicating that CO_2 decreases as the temperature increases, and vice-versa, with an average coefficient of -1.2 ppm K^{-1} (table 2).

4.2. Daily CO_2 concentration variability

In this section we focus on the daily variability in CO_2 concentrations, typically caused by the movement of synoptic-scale high and low pressure systems. To illustrate this we compare the daily averaged observed and predicted CO_2 concentrations between January 2020 and June 2020 (figure 3). Note that no data from 2020 was used to build the MLR models. The daily variability is largely driven by transport and mixing of CO_2 in the atmospheric boundary layer. High CO_2 concentrations occur when the boundary layer is shallow. During these conditions mixing is suppressed and emissions do not disperse rapidly away from sources but are trapped within the boundary layer where they can accumulate. The MLR models quantify this negative relationship with an average coefficient of $-0.005 \text{ ppm m}^{-1}$. Since the daily averaged BLD can vary by several hundred metres this can result in CO_2 variability of $1\text{--}2 \text{ ppm day}^{-1}$. In addition, for certain wind directions, transport from regional CO_2 sources towards the measurement site occurs resulting in high CO_2 concentrations. The coefficients used in the MLR models for easterly wind speeds are all positive. This suggests that easterly winds, which advect air from mainland Europe, contain higher CO_2 concentrations than westerly winds which transport relatively low CO_2 concentration air from the North Atlantic. The coefficients for northerly wind speeds are more mixed. The Tacolneston and Bilsdale MLR models contain negative coefficients indicating that southerly winds increase CO_2 concentrations. This is consistent with their locations

which have a long fetch of sea to their north. Conversely, the Heathfield and Mace Head MLR models contain positive coefficients indicating that northerly winds increase CO_2 at these sites. Heathfield is located south of several large urban areas so is potentially influenced by CO_2 emitted locally. Finally, Ridge Hill has no significant correlation with northerly wind direction since there are sources of CO_2 to both the north and south. Thus the modelled CO_2 concentrations capture the daily variability in CO_2 concentrations due to the inclusion of wind speed, wind direction and boundary layer depth in the MLR models.

4.3. MLR model evaluation

Over the training period (January 2015–December 2019), the MLR models capture 75% of the observed variability in daily averaged CO_2 concentrations with a root mean square error (RMSE) of 3.71 ppm (table 3). The RMSE in daily average CO_2 concentrations is relatively large due to an underestimation of the spikes in the observed daily CO_2 concentrations which are likely to be due to local emissions of CO_2 occurring within a few km's surrounding the tall towers. The normalised mean bias (NMB) is close to zero for all sites. The highest correlations (R^2) and lowest RMSE are found at Mace Head and Bilsdale. These sites have relatively small daily variability compared to the other sites suggesting that they are influenced less by local sources of pollution. Over the 2015–2019 training period the models explain more of the variability in the Spring/Summer ($R^2 = 0.77$) than in the Autumn/Winter periods ($R^2 = 0.68$) and the RMSE is lower (3.36 and 4.01 ppm respectively) (table 3). This is due to spikes in the observed daily CO_2 concentrations which occur predominantly during the winter and can reach 440 ppm (figure 2).

During 2020 the correlations are lower than during the training period but the MLR models still explain on average 67% of the observed variability in daily CO_2 (table 3). After the 16 March 2020 (UK lockdown) the RMSE increases at four out of the five sites (figure 3). However, none of the MLR models systematically overestimate the observed CO_2 concentrations after the UK lockdown demonstrating that it will take longer than 2 months for any signal of reduced CO_2 emissions to be observed in the atmospheric CO_2 concentrations.

5. Global CO_2 emission scenarios

Since the MLR models describe so much of the observed daily CO_2 variability they provide a realistic substitute for the real world and thus can be used to perform emission scenario simulations. In particular, the MLR models are used in this section to determine how long it would take for




Figure 3. CO₂ concentrations from 1 January to 30 May 2020 at (a) Ridge Hill, (b) Tacolneston, (c) Bilsdale, (d) Ridge Hill and (e) Mace Head. Daily average observed CO₂ concentrations (blue) and modelled CO₂ concentrations (red). Dashed line indicates date of UK lockdown on 16 March 2020.

COVID-19/Paris Agreement magnitude CO₂ emissions reductions to be detected in daily and monthly averaged CO₂ measurements. Since CO₂ has a lifetime much longer than 5 years, simple global emission

scenario simulations can be performed by scaling the regression coefficient controlling the trend (i.e. the date) whilst maintaining the seasonal and daily variability. $\beta_1 = 0.007 \text{ ppm day}^{-1}$ represents 100% of the

Table 3. Correlation (R^2), RMSE and normalised mean bias (NMB) statistics for daily model prediction of CO₂ concentration at each measurement site. Statistics are calculated for the model training period (January 2015–December 2019), Autumn/Winter months (September–February) in the training period, Spring/Summer months (March–August) in the training period and for the prediction period (January–December 2020).

	Ridge Hill	Tacolneston	Bilsdale	Mace Head	Heathfield	Average
2015–2019						
R^2	0.75	0.71	0.79	0.78	0.73	0.75
RMSE (ppm)	3.77	3.95	3.32	3.21	4.28	3.71
NMB (%)	−0.02	0.01	0.01	−0.01	−0.003	0.00
Autumn/Winter						
R^2	0.71	0.57	0.71	0.76	0.63	0.68
RMSE (ppm)	3.94	4.53	3.55	3.10	4.94	4.01
NMB (%)	0.04	−0.05	0.00	0.10	−0.03	0.01
Spring/Summer						
R^2	0.76	0.74	0.79	0.79	0.75	0.77
RMSE (ppm)	3.43	3.36	3.11	3.23	3.68	3.36
NMB (%)	−0.07	0.08	−0.02	−0.09	0.04	−0.01
2020						
R^2	0.62	0.65	0.75	0.68	0.65	0.67
RMSE (ppm)	3.52	3.81	2.96	2.97	4.03	3.46
NMB (%)	−0.03	−0.15	0.05	−0.01	0.03	−0.02

annual CO₂ concentration increase due to increasing anthropogenic emissions and $\beta_1 = 0.0 \text{ ppm day}^{-1}$ represents net-zero anthropogenic emissions. Thus sensitivity to different global emission scenarios can be performed while keeping the seasonal and daily variability constant (i.e. the regression coefficients for wind speed and wind direction, boundary layer depth and monthly averaged temperature remain unchanged).

The variability in daily CO₂ concentrations (daily noise) is estimated by the standard deviation of observed CO₂ concentrations over 30-day moving windows (figure 4). The variability in monthly CO₂ concentrations (monthly noise) is estimated by the standard deviation of the 2015–2019 de-trended observed CO₂ concentrations over moving 3 month periods. The difference in simulated CO₂ concentration between the 100% emissions and reduced global emissions scenarios (signal) increases with time and is proportional to the magnitude of the global emission reduction. The signal-to-noise ratio thus determines how reductions in CO₂ concentrations resulting from the emissions reduction scenarios compare to the estimated variability in CO₂ concentrations. The time of emergence is defined as the earliest time that the signal-to-noise ratio exceeds a value of 1. Since the time of emergence may depend on the initial conditions it is calculated for simulations initialised at varying weekly intervals between January 2015 and January 2020 to give a range of emergence times for each emission scenario. Figure 4(a) shows the evolution of daily CO₂ concentrations and daily noise for Ridge Hill assuming 100% emissions (plotted every 7 days). Different emission scenarios are also shown. For the Ridge Hill simulation initialised on 15 January 2015 (figure 4(a)) the time of emergence for the

net-zero scenario simulation (−100%) occurs 10.3 months after the start of the simulation. The time of emergence for the −50%, −25% and −12% emissions scenarios occur 15.3, 27.3 and 50.6 months after the start of the simulation respectively. Figure 4(b) shows simulations initialised at the same time for the Mace Head site. The time of emergence for the net-zero scenario is similar to that at Ridge Hill, for this initialisation time, but the time of emergence for the −50%, −25% and −12% emissions scenarios occurs earlier than the respective emission scenarios at Ridge Hill.

Table 4 shows the range of time of emergence for multiple emission scenarios initialised at monthly intervals. If net-zero anthropogenic emissions are assumed (−100%) then a signal would be detectable in the daily CO₂ concentrations after an average of 8 months. The signal in daily CO₂ concentrations would likely emerge at Bilsdale and Mace Head 2–3 months earlier than the other DECC sites as the daily variability at these sites is smaller than at the other DECC measurement locations. The longest daily time of emergence for the net-zero emission scenario would likely be at the Heathfield site, which is a semi-rural UK site located 19 km south of Royal Tunbridge Wells (population 118 000), in East Sussex, UK. As the emission scenario reduces in magnitude, the daily time of emergence increases. If a 50% reduction in anthropogenic emissions is maintained indefinitely a reduction in the trend of daily CO₂ concentrations would be observable after an average of 15 months. For COVID-19/Paris Agreement like magnitude emissions reductions of −12% (Andreoni 2021) the daily time of emergence would be on average after 38 months. Thus we would be able to detect a reduction in daily CO₂ concentrations after 2–3 years




Figure 4. Example 24-hour averaged predicted CO₂ concentrations for the (a) Ridge Hill and (b) Mace Head site with 100% emissions (black), −12% emissions (orange), −25% emissions (red), −50% emissions (purple) and −100% emissions (blue). Grey shading shows 100% emissions simulations ± 1 standard deviation of the 24-hour averaged observed CO₂ concentrations for a centred 30-day window. Simulations initialised on 15 January 2015. Vertical lines show the time at which signal-to-noise ratio exceeds 1 (time of emergence) for the different emission scenarios.

Table 4. Time of emergence (ToE, months) of CO₂ concentration differences due to reduced emission scenarios. The ranges are the 25–75th percentile ToE estimated using different start dates and for two averaging periods (24 hours and 30 days).

	Ridge Hill	Tacolneston	Bilsdale	Mace Head	Heathfield	Average
Daily ToE						
−12% emissions	42–48	20–31	32–48	29–37	47–61	38
−25% emissions	24–30	17–28	18–26	17–20	37–41	24
−50% emissions	14–19	12–18	11–15	8–14	19–24	15
−100% emissions	8–12	7–10	6–9	6–8	10–13	8
Monthly ToE						
−12% emissions	8–13	9–15	6–14	8–13	7–15	11
−25% emissions	5–10	6–12	5–10	5–9	5–11	8
−50% emissions	4–8	5–9	4–8	4–8	4–8	6
−100% emissions	3–6	3–6	3–6	3–7	3–6	5

depending on the measurement site. Note that for the smallest emission reduction scenario (−12% emissions) there are large daily time of emergence inter-quartile ranges due to a decrease in the sample size.

If we average the daily data to calculate monthly CO₂ concentrations then we smooth out the daily variability. Thus we can detect a reduction in the monthly CO₂ concentration trend due to COVID-19 like magnitude emissions reductions earlier, after about 11 months. Therefore, if current global lockdown restrictions continue we might detect a reduction in monthly averaged CO₂ concentration trend some time in 2021 at the earliest. When averaging over a month, the differences in the time of emergence between the measurement sites reduces as the CO₂ concentrations are less dependent on local emissions and are largely driven by non-local biogenic emissions.

6. Conclusions

In this paper, analysis of 5 CO₂ monitoring sites in the UK and Ireland has shown that after several months of CO₂ emissions reductions there are no detectable decreases in CO₂ concentrations exceeding the natural variations in measured CO₂ concentrations. Furthermore, global emission reduction scenario experiments show that it would take around 3 years of sustained global emissions reductions before any such signal could be detected in the local daily CO₂ concentration trend and 1 year before a reduction in CO₂ concentration trend would be detectable in the monthly averaged local CO₂ concentration trend. Future work could include performing the linear regression modelling using the fossil fuel contribution of CO₂ calculated from the measured ¹⁴C content of CO₂, instead of using total CO₂

concentrations. Since the method used to create the MLR models is generalizable, similar MLR models could be built for other locations with only local CO₂ and meteorological measurements available. It would be interesting to perform a similar study at a remote location, such as the Mauna Loa observatory, which is not affected by local CO₂ emissions in order to determine if the time of emergence appears earlier or later than those estimated for the sites in the UK and Ireland. The models used to make these estimates do not include climate feedbacks or processes determining plant growth which may make any detection of any signal even more difficult, hence these results should be seen as a lower limit. The results of this study show that the growth rate of CO₂ in the atmosphere will not decrease unless there is a substantial and persistent reduction in emissions over many decades.

Since CO₂ emissions are projected to eventually return to business-as-usual levels, the overall impact of COVID-19 CO₂ emission reductions on CO₂ concentrations in the atmosphere and therefore on climate change is likely to be small in the long run Forster *et al* (2020). The COVID-19 CO₂ emission reductions are similar in magnitude to those that are necessary to mitigate the worst effects of climate change. The COVID-19 crisis thus offers insights into the substantial changes in behaviour and infrastructure that are necessary if we are to achieve the temperature targets set out by the Paris Agreement. However, the measures deployed in response to the COVID-19 pandemic are not suitable or sustainable in the long term. These results support the need to create policies for recovering from the current economic downturn that do not further increase CO₂ emissions but which provide sustainable growth such as those outlined by Hepburn *et al* (2020).

The simple linear regression models used in this study could be used in the future to detect global scale emissions changes. However, the results of this study demonstrate that, using local measurements alone, there will be a significant delay between changes in global emissions and a detected signal in the local CO₂ concentrations.

Data availability statement


CO₂ data from the UK DECC network are available from the Centre for Environmental Data Analysis (CEDA) data archive (<https://catalogue.ceda.ac.uk/uuid/a18f43456c364789aac726ed365e41d1>) DECC (2020). Atmospheric CO₂ data from Mace Head is available at the ICOS Carbon Portal (<https://www.icos-cp.eu/>): doi:10.18160/ere9-9d85 Ramonet *et al* (2020).

Acknowledgments

The UK DECC network operations were funded by the UK Department of Business, Energy and Industrial Strategy (BEIS) through contract 1537/06/2018 to the University of Bristol. The authors would like to thank Kieran Stanley, Dickon Young, Ann Stavert, Aoife Grant and Anita Ganesan at the University of Bristol for setting up and running the DECC measurement sites. We would also like to thank Dr Adam Wisher, the specialist technician at Tacolneston, for maintaining the instrument. The Mace Head data is provided by ICOS. Maintenance of the Heathfield site measurements is supported by UK National Measurement System funding to the National Physical Laboratory. H F Dacre is funded by the NERC Detection and Attribution of Regional greenhouse gas Emissions in the UK (DARE-UK) project NE/S004505/1.

ORCID iDs

H F Dacre  <https://orcid.org/0000-0003-4328-9126>

L M Western  <https://orcid.org/0000-0002-0043-711X>

D Say  <https://orcid.org/0000-0003-3615-7926>

References

- Andreoni V 2021 Estimating the European CO₂ emissions change due to COVID-19 restrictions *Sci. Total Environ.* **769** 145115
- Ballard S P, Li Z, Simonin D and Caron J F 2016 Performance of 4D-Var NWP-based nowcasting of precipitation at the Met Office for summer 2012 *Q. J. R. Meteorol. Soc.* **142** 472–87
- Carslaw D C and Beevers S D 2005 Development of an urban inventory for road transport emissions of NO₂ and comparison with estimates derived from ambient measurements *Atmos. Environ.* **39** 2049–59
- Clark P, Roberts N, Lean H, Ballard S P and Charlton-Perez C 2016 Convection-permitting models: a step-change in rainfall forecasting *Meteorol. Appl.* **23** 165–81
- Dacre H F, Mortimer A H and Neal L S 2020 How have surface NO₂ concentrations changed as a result of the UK's COVID-19 travel restrictions? *Environ. Res. Lett.* **15** 104089
- DECC 2020 Deriving Emissions linked to Climate Change (DECC Network (available at: <http://data.ceda.ac.uk/badc/ukdecc-network>) (accessed April 2020)
- Forster P M *et al* 2020 Current and future global climate impacts resulting from COVID-19 *Nat. Clim. Change* **10** 913–19
- Friedlstein P *et al* 2020 Global carbon budget 2020 *Earth Syst. Sci. Data* **12** 3269–340
- Granados J A T, Ionides E L and Carpintero o 2012 Climate change and the world economy: short-run determinants of atmospheric CO₂ *Environ. Sci. Policy* **21** 50–62
- Hale G and Leduc S 2020 COVID-19 and CO₂ *FRBSF Economic Lett.* **2020** 1–06
- Harvey N J, Hogan R J and Dacre H F 2013 A method to diagnose boundary-layer type using Doppler lidar *Q. J. R. Meteorol. Soc.* **139** 1681–93
- Harvey N J, Hogan R J and Dacre H F 2015 Evaluation of boundary-layer type in a weather forecast model utilizing long-term Doppler lidar observations *Q. J. R. Meteorol. Soc.* **141** 1345–53

- Hepburn C, O'Callaghan B, Stern N, Stiglitz J and Zenghelis D 2020 Will COVID-19 fiscal recovery packages accelerate or retard progress on climate change? *Oxford Rev. Economic Policy* 36 (<https://www.smithschool.ox.ac.uk/publications/wpapers/workingpaper20-02.pdf>)
- IPCC, 2018: Summary for policymakers In: *Global Warming of 1.5 °C. An IPCC Special Report on the impacts of global warming of 1.5 °C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty* [Masson-Delmotte V, et al (eds.)] World Meteorological Organization, Geneva, Switzerland, p 32
- Le Quéré C et al 2020 Temporary reduction in daily global CO₂ emissions during the COVID-19 forced confinement *Nat. Clim. Change* 10 647–53
- Lean H W, Clark P A, Dixon M, Roberts N M, Fitch A, Forbes R and Halliwell C 2008 Characteristics of high-resolution versions of the Met Office Unified Model for forecasting convection over the United Kingdom *Mon. Weather Rev.* 136 3408–24
- Liu Z, Deng Z, Ciais P, Lei R, Feng S, Davis S J, Wang Y, Yue X, Lei Y, Zhou H and Cai Z, 2020 2020 COVID-19 causes record decline in global CO₂ emissions *Preprint* (available at: <http://arxiv.org/abs/2004.13614>)
- Manning A J, O'Doherty S, Jones A R, Simmonds P G and Derwent R G 2011 Estimating UK methane and nitrous oxide emissions from 1990 to 2007 using an inversion modeling approach *J. Geophys. Res.: Atmos.* 116 D2
- Mitra A, Ray Chadhuri T, Mitra A, Pramanick P and Zaman S 2020 Impact of COVID-19 related shutdown on atmospheric carbon dioxide level in the city of Kolkata *Parana J. Sci. Edu.* 6 84–92
- Quadrelli R and Peterson S 2007 The energy–climate challenge: recent trends in CO₂ emissions from fuel combustion *Energy Policy* 35 5938–52
- Ramonet M 2020 The fingerprint of the summer 2018 drought in Europe on ground-based atmospheric CO₂ measurements *Philosophical Transactions of the Royal Society B* 375 20190513
- Roberts N M and Lean H W 2008 Scale-selective verification of rainfall accumulations from high-resolution forecasts of convective events *Mon. Weather Rev.* 136 78–97
- Samset B H, Fuglestad J S and Lund M T 2020 Delayed emergence of a global temperature response after emission mitigation *Nat. Commun.* 11 1–10
- Shi J P and Harrison R M 1997 Regression modelling of hourly NO_x and NO₂ concentrations in urban air in London *Atmos. Environ.* 31 4081–94
- Sitch S et al 2015 Recent trends and drivers of regional sources and sinks of carbon dioxide *Biogeosciences* 12 653–79
- Stainforth D A et al 2005 Uncertainty in predictions of the climate response to rising levels of greenhouse gases *Nature* 433 403–6
- Stanley K M et al 2018 Greenhouse gas measurements from a UK network of tall towers: technical description and first results *Atmos. Meas. Tech.* 11 1437–58
- Stavert A R, O'Doherty S, Stanley K, Young D, Manning A J, Lunt M F, Rennick C and Arnold T 2019 UK greenhouse gas measurements at two new tall towers for aiding emissions verification *Atmos. Meas. Tech.* 12 4495–518
- Tang Y, Lean H W and Bornemann J 2013 The benefits of the Met Office variable resolution NWP model for forecasting convection *Meteorol. Appl.* 20 417–26
- Taylor K E and Penner J E 1994 Response of the climate system to atmospheric aerosols and greenhouse gases *Nature* 369 734–7
- Tebaldi C and Friedlingstein P 2013 Delayed detection of climate mitigation benefits due to climate inertia and variability *Proc. Natl Acad. Sci.* 110 17229–34
- Vardag S N, Hammer S, O'Doherty S, Spain T G, Wastine B, Jordan A and Levin I 2014 Comparisons of continuous atmospheric CH₄, CO₂ and N₂O measurements—results from a travelling instrument campaign at Mace Head *Atmos. Chem. Phys.* 14 8403–18